arXiv:math/0510348v3 [math.RT] 17 Nov 2009

n-dimensional global correspondences of Langlands

C. Pierre

Institut de Mathématique pure et appliquée
Université de Louvain
Chemin du Cyclotron, 2
B-1348 Louvain-la-Neuve, Belgium
pierre@math.ucl.ac.be

Mathematics subject classification (1991): 11G18, 11R34, 11R37, 11R39.

Abstract

The program of Langlands is studied here on the basis of:

- new concepts of global class field theory related to the explicit construction of global class fields and of reciprocity laws;
- the representations of the reductive algebraic groups GL(n) constituting the n-dimensional representations of the associated global Weil-Deligne groups;
- a toroidal compactification of the conjugacy classes of these reductive algebraic groups whose analytic representations constitute the cuspidal representations of these groups GL(n) in the context of harmonic analysis.

This leads us to build two types of n-dimensional global bilinear correspondences of Langlands by taking into account the irreducibility or the reducibility of the representations of the considered bilinear algebraic semigroups.

The major outcome of this global approach is the generation of general algebraic symmetric structures, consisting of double symmetric towers of conjugacy class representatives of algebraic groups, so that the analytic toroidal representations of these conjugacy classes are the cuspidal conjugacy class representations of these algebraic groups.

Contents

Introduction **1**

0.1 Nonabelian global class field concepts 5

0.2 Langlands irreducible global bilinear correspondences 9

0.3 Langlands reducible global bilinear correspondences 13

1 Global class field concepts, motivic chain bicomplexes and pure Chow bimotives **16**

1.1 Global class field concepts 16

1.2 Suslin-Voevodsky motivic bicomplexes 26

1.3 Chow motives 28

2 Nonabelian global class field concepts based on the representation of an algebraic bilinear semigroup **33**

2.1 The general bilinear algebraic semigroup 33

2.2 Lattices and Hecke algebras 44

2.3 Lattice semispaces 50

2.4 Representations of global Weil groups 52

2.5 Trace formulas 61

3 Langlands global correspondences for irreducible representations of $\mathrm{GL}(n)$ **75**

Introductive keys 75

3.1 Compactifications of bilinear algebraic semigroups and global holomorphic correspondences 76

3.2 General bilinear cohomology theory 86

3.3 Borel-Serre toroidal compactification 91

3.4 Langlands global correspondence on the irreducible algebraic bilinear semigroup $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$ 97

3.5 Langlands global correspondence on the boundary $\partial\overline{X}^{(I\!R)}_{S_{R\times L}}$ of the irreducible compactified lattice bisemispace $\overline{X}_{S_{R\times L}}$ 110

4 Langlands global correspondences for reducible representations of $\mathrm{GL}(n)$ **118**

4.1 Reducibility of $\mathrm{GL}_{(2)n}(F_{\overline{\omega}} \times F_\omega)$ 118

4.2 Langlands global correspondences on the reducible compactified bisemispaces $\overline{X}^{(2)n}_{S_{R\times L}}$ and $\partial\overline{X}^{(2)n}_{S_{R\times L}}$ 122

Appendix **136**

1 Philosophy of bisemiobjects 136
2 Semistructures 136
3 Bisemistructures [Pie5] 138

References **142**

Introduction

After the pioneer works of G. Laumon, M. Rapoport, U. Stuhler and others, recent significant advances were realized these past few years in the Langlands program [Lan1], [Kna] by M. Harris, R. Taylor [H-T] and G. Henniart [Hen] with respect to the proof of the Langlands correspondences in the case of p-adic number fields and by L. Lafforgue [Laf], [Lau] in the case of function fields.

However, the approach of the Langlands program from a global point of view was relatively left sideways until now: it is the aim of this paper to try to fill up partially this gap by studying the Langlands correspondences over global number fields.

The global methods used in this context are based on:

- new concepts in global class field theory related to the explicit construction of global class fields and of reciprocity laws;
- the representations of the reductive algebraic groups GL(n) constituting the n-dimensional representations of the associated global Weil-Deligne groups;
- a toroidal compactification of the conjugacy classes of these algebraic groups whose analytic representations constitute the cuspidal representations of these groups GL(n) in the context of harmonic analysis. It will be shown that the global approach of the Langlands program consists in studying the algebraic groups and their representations to bridge the gap between analytic and algebraic problems as described in the diagram:



Algebraic number theory
• global class field theory
• Galois and Weil groups
Theory (of representations) of Algebraic groups GL(n)
Toroidal compactification
Analytic number theory Harmonic analysis
n-dimensional representations of the Weil-Deligne group
∼
Langlands correspondences
supercuspidal representations of GL(n)

The developments considered here will essentially concern symmetric objects in such a way that the envisaged mathematical objects will be cut into two symmetric semiobjects $\mathcal{O}_R$ and $\mathcal{O}_L$ localized respectively in (or referring to) the lower and in the upper half space.

The right semiobject $\mathcal{O}_R$ is then the dual of the left semiobject $\mathcal{O}_L$ and the interest of considering an object, decomposed into two dual symmetric semiobjects $\mathcal{O}_R$ and $\mathcal{O}_L$, is that the informations concerning the internal mathematical structure of the object " $\mathcal{O}$ "

can be obtained from the product $\mathcal{O}_R \times \mathcal{O}_L$ of the semiobjects $\mathcal{O}_R$ and $\mathcal{O}_L$. Indeed, every endomorphism E of " $\mathcal{O}$ " can be decomposed into the product $E_R \times E_L$ of a right endomorphism E_R acting on the right semiobject $\mathcal{O}_R$ by the "opposite" left endomorphism E_L acting on the corresponding left semiobject $\mathcal{O}_L$ in such a way that $E_R \simeq E_L^{-1}$.

The general existence of symmetric objects can be shown by the following considerations on function fields. Let k denote a global number field of characteristic zero. Let $k[x_1, \cdots, x_n]$ be a polynomial ring over k and let $R = R_R \cup R_L$ be a symmetric (closed) extension of k as introduced in the following.
Let $k[x_1, \cdots, x_n]$ be a polynomial ring at n indeterminates over k .
Let $I_L = \{P_\mu(x_1, \cdots, x_n) \mid P_\mu(V_L) = 0\}$ be the ideal of $k[x_n, \cdots, x_n]$ in such a way that:

a) $P_\mu(V_L)$ be the polynomial function in $k[V_L]$ represented by $P_\mu(x_1, \cdots, x_n)$.

b) $V_L \subset R_L$ be an affine semispace restricted to the upper half space.

Let $I_R = \{P_\mu(-x_1, \cdots, -x_n) \mid P_\mu(V_R) = 0\}$ be the symmetrical ideal of I_L obtained by the involution:

$$\tau : \quad P_\mu(x_1, \cdots, x_n) \longrightarrow P_\mu(-x_1, \cdots, -x_n)$$

in such a way that

a) $P_\mu(V_R)$ be the polynomial function in $k[V_R]$ represented by $P_\mu(-x_1, \cdots, -x_n)$.

b) $V_R \subset R_R$ be an affine semispace

- of dimension n .
- restricted to the lower half space.
- symmetric of V_L .
- disjoint of V_L or possibly connected to V_L on the symmetric axis, plane, ...

The quotient ring obtained modulo the ideal I_L (resp. I_R) is $Q_L = k[x_1, \cdots, x_n]/I_L$ (resp. $Q_R = k[x_1, \cdots, x_n]/I_R$) [Wat].
Q_L and Q_R are quotient algebras characterized by the corresponding homomorphisms:

$$\phi_L : \quad Q_L \longrightarrow R_L , \qquad\qquad \phi_R : \quad Q_R \longrightarrow R_R ,$$

where R_L and R_R are commutative (division) semirings localized respectively in (or referring to) the upper and the lower half space. (Commutative (division) semirings are recalled (or introduced) in the appendix).

So, the pair of homomorphisms ϕ_L and ϕ_R sends the “general” solution to a pair of symmetric solutions respectively in R_L and R_R .
On the other hand, let $\widetilde{F}=\widetilde{F}_R\cup\widetilde{F}_L$ be a symmetric “algebraic” finite extension of k .
Let $T_n(\widetilde{F}_L)$ (resp. $T_n^t(\widetilde{F}_R)$) $\subset \mathrm{GL}_n(\centerdot)$ be the (semi)group of matrices of dimension n over $\widetilde{F}_L$ (resp. $\widetilde{F}_R$) viewed as an operator sending $\widetilde{F}_L$ (resp. $\widetilde{F}_R$) into the affine semispace $T^{(n)}(\widetilde{F}_L)$ (resp. $T^{(n)}(\widetilde{F}_R)$) of dimension n :

$$\begin{aligned} T_n(\centerdot): \quad & \widetilde{F}_L \longrightarrow T^{(n)}(\widetilde{F}_L) \\ (\text{resp.} \quad T_n^t(\centerdot): \quad & \widetilde{F}_R \longrightarrow T^{(n)}(\widetilde{F}_R)\ ) \end{aligned}$$

in such a way that to the indeterminates $(x_1,\cdots,x_\ell,\cdots,x_n,\cdots,x_{1\ell},\cdots,x_{\ell n},\cdots)$ of Q_L (resp. $(-x_1,\cdots,-x_\ell,\cdots,-x_n,\cdots,-x_{1\ell},\cdots,-x_{\ell n},\cdots)$ of Q_R), $\forall\ x_{\ell n}=x_\ell\cdot x_n$, corresponds the homomorphism [Bor2]:

$$\begin{aligned} \phi'_L: \quad & Q_L \longrightarrow \widetilde{F}_L(x_1\to e_{11},\cdots,x_\ell\to e_{\ell\ell},\cdots,x_{1\ell}\to e_{1\ell},\cdots,x_{n\ell}\to e_{n\ell}) \\ (\text{resp.} \quad \phi'_R: \quad & Q_R \longrightarrow \widetilde{F}_R(-x_1\to -e_{11},\cdots,-x_\ell\to -e_{\ell\ell},\cdots, \\ & \qquad\qquad\qquad x_{\ell 1}\to e_{\ell 1},\cdots,x_{n\ell}\to e_{n\ell})\ ) \end{aligned}$$

where:

$$\begin{aligned} T_n(\widetilde{F}_L) &= \{e_L=(e_{\ell n})\in T_n(\widetilde{F}_L)\mid P_{T_\mu}(e_{\ell n})=0\} \\ (\text{resp.} \quad T_n^t(\widetilde{F}_R) &= \{e_R=(e_{\ell n})\in T_n^t(\widetilde{F}_R)\mid P_{T_\mu}(e_{\ell n})=0\}\ ) \end{aligned}$$

with the polynomials $P_{T_\mu}(e_{\ell n})\in k[x]$.
On the other hand, let X_L (resp. X_R) be a functor from the quotient ring Q_L (resp. Q_R) to the affine semispace V_L (resp. V_R) in such a way that the diagram:

$$\begin{tikzcd} Q_L \arrow[r, "\phi'_L"] \arrow[d, "X_L"] & \widetilde{F}_L(e_{11},\cdots,e_{nn},\cdots,e_{\ell n}) \arrow[d, "T_n(\centerdot)"] \\ V_L \arrow[r, "\psi_L"'] & T^{(n)}(\widetilde{F}_L) \end{tikzcd}$$

$$\begin{tikzcd} & Q_R \arrow[r, "\phi'_R"] \arrow[d, "X_R"] & \widetilde{F}_R(-e_{11},\cdots,-e_{nn},\cdots,e_{\ell n}) \arrow[d, "T_n^t(\centerdot)"] & \\ \Big(\text{resp} & {} & {} & \Big) \\ & V_R \arrow[r, "\psi_R"'] & T^{(n)}(\widetilde{F}_R) & \end{tikzcd}$$

commutes, leading to:

$$X_L=\psi_L^{-1}\circ T_n\circ\phi'_L \qquad (\text{resp.} \quad X_R=\psi_L^{-1}\circ T_n^t\circ\phi_R\ ).$$

Thus, if the elements $v_L \in V_L$ (resp. $v_R \in V_R$), resulting from $X_L \circ \phi_L^{'-1}(\widetilde{F}_L)$ (resp. $X_R \circ \phi_R^{'-1}(\widetilde{F}_R)$), correspond to solutions in $\widetilde{F}_L$ (resp. $\widetilde{F}_R$) of some family of equations, there is a k-algebra Q_L (resp. Q_R) and a natural correspondence between $X_L \circ \phi_L^{'-1}\widetilde{F}_L$ (resp. $X_R \circ \phi_R^{'-1}\widetilde{F}_R$) and $\mathrm{Hom}_k(Q_L, \widetilde{F}_L)$ (resp. $\mathrm{Hom}_k(Q_R, \widetilde{F}_R)$), taking into account the algebraic (semi)group of matrices $T_n(\widetilde{F}_L)$ (resp. $T_n^t(\widetilde{F}_R)$) at the condition that $\psi_L : V_L \to T^{(n)}(\widetilde{F}_L)$ (resp. $\psi_R : V_R \to T^{(n)}(\widetilde{F}_R)$) be a homeomorphism.

Such X_L (resp. X_R) is then called representable in the sense that a left (resp. right) affine semigroup scheme over k is a representable functor from the k-algebra Q_L (resp. Q_R) to the affine semispace V_L (resp. V_R), homeomorphic to $T^{(n)}(\widetilde{F}_L)$ (resp. $T^{(n)}(\widetilde{F}_R)$).

The left and right affine semigroup schemes X_L and X_R are said to be symmetric if every element $v_L \in V_L$, localized in the upper half space, is symmetric to every element $v_R \in V_R$, localized in the lower half space with respect to an element or a set of elements of symmetry.

Thus, the consideration of a sufficiently big polynomial ring as $k[x_1, \cdots, x_n]$ allows to consider objects as being generally symmetric and able to be cut into two symmetric semiobjects in one-to-one correspondence.

In this respect, as the endomorphisms $\mathrm{End}_{\widetilde{F}}(B)$ of a division $\widetilde{F}$-algebra B can be handled throughout its enveloping algebra $B^e = B \otimes_{\widetilde{F}} B^{\mathrm{op}}$, where B^{op} denotes the opposite algebra of B , because $B^e \simeq \mathrm{End}_{\widetilde{F}}(B)$ and as fundamental algebras, such as the algebra of automorphic forms, are essentially defined on half spaces, we shall work in a bilinear mathematical framework. So, enveloping semialgebras as well as bisemialgebras will be considered, which has the following advantages:

1. As the representation of automorphic forms is intrinsically defined in the upper half space, the bialgebra of automorphic biforms, associated to the algebra of automorphic forms, will be envisaged in such a way that the coalgebra will refer to dual automorphic forms localized in the lower half space. One of the interests of considering a bialgebra of automorphic biforms is that the endomorphism of this bialgebra leads to take into account the Hecke bialgebra of tensor products of Hecke bioperators having a nice matricial representation [Pie3].
2. The representation of a reducible general bilinear (semi)group of order $2n$ decomposes diagonally following the direct sum of irreducible bilinear representations and off diagonally following the direct sum of complementary irreducible bilinear representations (see section 0.3 of the introduction and chapter 4).

So, the nonorthogonal reducibility of the representation of a bilinear general Lie semigroup generating off diagonal complementary irreducible bilinear representations could lead to a new approach of:

- the endoscopy problem considered in trace formulas;
- the functoriality envisaged in automorphic representations of general "linear" groups [Lan2], [C-K-P-S].

3. Some conjectures of algebraic number theory proceed probably from the fact that the techniques of algebra endomorphisms could be more worked out, as for example, by considering judicious enveloping (semi)algebras [Pie3].

In this context, let us point out that the appendix of this paper introduces the used semistructures and bisemistructures [Pie5].

In a few words, let us say that the Langlands global program developed here is related to the generation of a reductive algebraic group with entries in a double tower of extensions of k which are characterized by increasing Galois extension degrees.

The conjugacy class representatives of this reductive algebraic group then appear in symmetric pairs in such a way that a double tower of conjugacy class representatives, characterized by increasing ranks, is generated symmetrically respectively in the upper and in the lower half space and constitutes the n-dimensional representation of a global Weil-Deligne group.

The Langlands global correspondences then consist in finding the toroidal analytic representations of these conjugacy class representatives by means of a suitable toroidal compactification of these in such a way that the sum of these toroidal analytical representatives of conjugacy classes constitutes the searched supercuspidal representation of the reductive algebraic group, i.e. the Fourier development of the n-dimensional automorphic form on this algebraic group.

So, the Langlands global program is directly based on the generation of a general algebraic symmetric structure being in one-to-one correspondence with its automorphic analytical counterpart.

0.1 Nonabelian global class field concepts

More concretely, the new concepts introduced in global class field theory are inspired by the factorization of a prime p into primes in the ring of integers $\mathcal{O}_E$ of a finite extension E of $\mathbb{Q}$ and by the Artin's reciprocity law which asserts that a Dirichlet character $\chi_\sigma : (\mathbb{Z}/N\,\mathbb{Z})^* \to \mathbb{C}^*$ exists such that $\sigma(\mathrm{Fr}\,p)=\chi_\sigma(p)$ where $\sigma : \mathrm{Gal}(E/\mathbb{Q}) \to \mathbb{C}^*$. This leads us to introduce pseudo-ramified completions, covered by the symmetric algebraic extensions of a global number field k of characteristic zero, at the infinite places such that their degrees or ranks are integers modulo N where N is the order of the global inertia subgroups. The subsets of pseudo-ramified completions are thus covered by finite subsets of Galois extensions and generate a double symmetric tower of one-dimensional k-semimodules characterized by increasing ranks. A character χ_{ω_j} is associated to each basic completion F_{ω_j} in an infinite or global place ω_j and the Dirichlet character $\chi_{\omega_{j,m_j}}$, corresponding to an equivalent completion $F_{\omega_{j,m_j}}$ in ω_j, can be obtained by the action of the decomposition group element $D_{\omega_{j,m_j}}$ on χ_{ω_j}. As we are concerned with pairs of symmetric (semi)structures, left algebraic extension semifields $\widetilde{F}_\omega=\{\widetilde{F}_{\omega_1},\dots,\widetilde{F}_{\omega_{j,m_j}},\dots,\widetilde{F}_{\omega_r}\}$ referring to the upper half space and right algebraic extension semifields $\widetilde{F}_{\overline{\omega}}=\{\widetilde{F}_{\overline{\omega}_1},\dots,\widetilde{F}_{\overline{\omega}_{j,m_j}},\dots,\widetilde{F}_{\overline{\omega}_r}\}$ referring to the lower half space are considered.

In correspondence with these left and right algebraic extension semifields, we have a left and a right tower $F_\omega=\{F_{\omega_1},\cdots,F_{\omega_{j,m_j}},\cdots,F_{\omega_r}\}$ and $F_{\overline{\omega}}=\{F_{\overline{\omega}_1},\cdots,F_{\overline{\omega}_{j,m_j}},\cdots,F_{\overline{\omega}_r}\}$ of packets of equivalent completions characterized by increasing ranks.

From these, we can consider:

a) their direct sums

$$F_{\omega_\oplus}=\bigoplus_j\bigoplus_{m_j}F_{\omega_{j,mj}} \qquad \text{and} \qquad F_{\overline{\omega}_\oplus}=\bigoplus_j\bigoplus_{m_j}F_{\overline{\omega}_{j,mj}}\ ,$$

b) and the products

$$\mathbb{A}^{\infty}_{F_\omega}=\prod_{j_p}F_{\omega_{j_p}} \qquad \text{and} \qquad \mathbb{A}^{\infty}_{F_{\overline{\omega}}}=\prod_{j_p}F_{\overline{\omega}_{j_p}}$$

of packets of primary completions $F_{\omega_{j_p}}$ and $F_{\overline{\omega}_{j_p}}$ where $\mathbb{A}^{\infty}_{F_\omega}$ and $\mathbb{A}^{\infty}_{F_{\overline{\omega}}}$ are infinite adele semirings.

This allows to introduce the Galois subgroups of these left and right algebraic extensions and the global Weil groups $W^{ab}_{F_L}$ and $W^{ab}_{F_R}$ referring respectively to (the sums of) the automorphisms of left and right algebraic extensions explicited subsequently.

The set of left (resp. right) pseudo-ramified extensions considered until now constitute a left (resp. right) affine semigroup $\mathbb{S}^1_L$ (resp. $\mathbb{S}^1_R$).

All that constitutes elementary concepts of abelian global class field theory. To enter into the n-dimensional global Langlands program, nonabelian global class field concepts have to be set up [Lan2]. The challenge then consists in building up the n-dimensional equivalent of the affine semigroup $\mathbb{S}^1_L$ (resp. $\mathbb{S}^1_R$), and from a bilinear point of view, the n-dimensional equivalent of the bilinear affine semigroup $\mathbb{S}^1_R \times \mathbb{S}^1_L$, which needs the introduction of the injective morphisms [Gel]:

$$\begin{array}{llcl}
\sigma_L : & W^{ab}_{F_L} & \longrightarrow & T_n(\widetilde{F}_{\omega_\oplus}) \ , \\[2ex]
\sigma_R : & W^{ab}_{F_R} & \longrightarrow & T^t_n(\widetilde{F}_{\overline{\omega}_\oplus}) \ , \\[2ex]
\sigma_{R\times L} : & W^{ab}_{F_R} \times W^{ab}_{F_L} & \longrightarrow & \mathrm{GL}_n(\widetilde{F}_{\overline{\omega}_\oplus} \times \widetilde{F}_{\omega_\oplus}) \equiv T^t_n(\widetilde{F}_{\overline{\omega}_\oplus}) \times T_n(\widetilde{F}_{\omega_\oplus}) \ .
\end{array}$$

The affine bilinear algebraic semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega) \equiv T^{(2n)}(\widetilde{F}_{\overline{\omega}}) \times T^{(2n)}(\widetilde{F}_\omega)$ results from the action of the bilinear algebraic semigroup of matrices $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega) \equiv T^t_n(\widetilde{F}_{\overline{\omega}}) \times T_n(\widetilde{F}_\omega)$ where $T^t_n(\widetilde{F}_{\overline{\omega}})$ is the group of lower triangular matrices with entries in $\widetilde{F}_{\overline{\omega}}$ while $T_n(\widetilde{F}_\omega)$ is the group of upper triangular matrices with entries in $\widetilde{F}_\omega$. The algebraic bilinear affine semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$ over $\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega$ is a $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$-bisemimodule $\widetilde{M}_R \otimes \widetilde{M}_L$. And, the algebraic representation of $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$ in $(\widetilde{M}_R \otimes \widetilde{M}_L)$ refers to an algebraic morphism from $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$ into $\mathrm{GL}(\widetilde{M}_R \otimes \widetilde{M}_L)$ which denotes the group of automorphisms of $(\widetilde{M}_R \otimes \widetilde{M}_L)$.

The reasons for considering bilinear algebraic (semi)groups $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$ instead of linear algebraic groups $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}-\omega})$ result from the facts that:

1. a bilinear algebraic semigroup covers the corresponding linear algebraic group (as it is proved in chapter 2) in the sense that the representation space of this linear algebraic group is a n^2-complex dimensional vector space W corresponding to the n^2-complex dimensional representation space $(\widetilde{M}_R \otimes \widetilde{M}_L)$ of the bilinear algebraic (semi)group $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$. So, we have: $W \simeq \widetilde{M}_R \otimes \widetilde{M}_L$;

2. a bilinear algebraic semigroup is directly connected to the enveloping algebras allowing to generate endomorphisms as it was explained above.

The bilinear semigroup $G^{(2n)}(F_{\overline{\omega}} \times F_\omega)$ over the product, right by left, of pseudo-ramified completions $F_{\overline{\omega}}$ and F_ω, is a complete "algebraic" bilinear semigroup or an abstract bisemivariety since the sets of completions $F_{\overline{\omega}}$ and F_ω are covered by the corresponding sets of algebraic extensions $\widetilde{F}_{\overline{\omega}}$ and $\widetilde{F}_\omega$. By concern of abbreviation, this complete "algebraic" bilinear semigroup $G^{(2n)}(F_{\overline{\omega}} \times F_\omega)$ will sometimes be simply called "algebraic" bilinear semigroup, taking into account a universal property between these bilinear semigroups.

Let $\widetilde{M}_{R_\oplus} \otimes \widetilde{M}_{L_\oplus}$ denote the representation space of $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}_\oplus} \times \widetilde{F}_{\omega_\oplus})$ in such a way that $\widetilde{M}_{R_\oplus} \otimes \widetilde{M}_{L_\oplus}$ decomposes into the direct sum of subbisemimodules $\widetilde{M}_{\overline{\omega}_{j,m_j}} \otimes \widetilde{M}_{\omega_{j,m_j}}$ being the conjugacy class representatives of $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$.

Then, $\mathrm{GL}(\widetilde{M}_{R_\oplus} \otimes \widetilde{M}_{L_\oplus})$ constitutes the n-dimensional complex equivalent of the product $W^{ab}_{F_R} \times W^{ab}_{F_L}$ of the global Weil groups. As $\mathrm{GL}(\widetilde{M}_{R_\oplus} \otimes \widetilde{M}_{L_\oplus})$ is isomorphic to $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}_\oplus} \times \widetilde{F}_{\omega_\oplus})$, the bilinear algebraic semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}_\oplus} \times \widetilde{F}_{\omega_\oplus})$ becomes the $2n$-dimensional (irreducible) complex representation (space) $(\mathrm{Irr})\,\mathrm{Rep}^{(2n)}_{W_{F_{R\times L}}}(W^{ab}_{F_R} \times W^{ab}_{F_L})$ of $W^{ab}_{F_R} \times W^{ab}_{F_L}$:

$$(\mathrm{Irr})\,\mathrm{Rep}^{(2n)}_{W_{F_{R\times L}}}(W^{ab}_{F_R} \times W^{ab}_{F_L}) : \quad \mathrm{GL}(\widetilde{M}_{R_\oplus} \otimes \widetilde{M}_{L_\oplus}) \longrightarrow G^{(2n)}(\widetilde{F}_{\overline{\omega}_\oplus} \times \widetilde{F}_{\omega_\oplus}) \ .$$

As the bilinear algebraic semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$ is built over $(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$, it is composed of r conjugacy classes, $1 \le j \le r \le \infty$, having multiplicities $m^{(r)}$ and associated with the r places of F_ω or of $F_{\overline{\omega}}$. Consequently, $(\mathrm{Irr})\,\mathrm{Rep}^{(2n)}_{W_{F_{R\times L}}}(W^{ab}_{F_R} \times W^{ab}_{F_L})$ also decomposes into r conjugacy classes with multiplicities $m^{(r)}$.

This decomposition of $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$ into conjugacy classes can also be obtained by considering the cutting of bilattices $\Lambda_{\overline{\omega}} \times \Lambda_\omega$ into subbilattices in $(\widetilde{M}_R \otimes \widetilde{M}_L)$ under the action of the product $T_R(n;r) \otimes T_L(n;r)$ of Hecke operators having as representation $\mathrm{GL}_{2n}((\mathbb{Z}/N\,\mathbb{Z})^2)$, $N \in \mathbb{N}$.

This leads to a one-to-one correspondence between the pseudo-ramified extensions $\widetilde{F}_{\omega_j}$ (and $\widetilde{F}_{\omega_{j,m_j}}$), the diagonal (and off-diagonal) conjugacy classes of the bilinear algebraic semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$ and the subbilattices of Hecke in the $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$-bisemimodule $\widetilde{M}_R \otimes \widetilde{M}_L$.

Let $G^{(2n)}(\widetilde{F}^+_{\overline{v}} \times \widetilde{F}^+_{v})$ denote the bilinear algebraic semigroup over the product of the symmetric sets $\widetilde{F}^+_{\overline{v}} = \{\widetilde{F}^+_{\overline{v}_{1_\delta}}, \cdots, \widetilde{F}^+_{\overline{v}_{j_\delta,m_{j_\delta}}}, \cdots, \widetilde{F}^+_{\overline{v}_{r_\delta}}\}$ and $\widetilde{F}^+_{v} = \{\widetilde{F}^+_{v_{1_\delta}}, \cdots, \widetilde{F}^+_{v_{j_\delta,m_{j_\delta}}}, \cdots, \widetilde{F}^+_{v_{r_\delta}}\}$ of real extensions and let $G^{(2n)}(F^+_{\overline{v}} \times F^+_{v})$ denote its compact equivalent. The compactification of $G^{(2n)}(\widetilde{F}^+_{\overline{v}} \times \widetilde{F}^+_{v})$ into $G^{(2n)}(F^+_{\overline{v}} \times F^+_{v})$ is realized by means of the Fulton-McPherson compactification by a set of successive blowups [F-M] as developed in chapter 3.

If the conjugacy class representatives $g^{(2n)}_{R\times L}[j,m_j]$ of $G^{(2n)}(F^+_{\overline{\omega}} \times F^+_{\omega})$ are glued together, the bifunction $(f_{\overline{\omega}}(z^*) \otimes f_\omega(z))$ on $G^{(2n)}(F^+_{\overline{\omega}} \times F^+_{\omega})$ has a holomorphic development in the multiple power series

$$f_{\overline{\omega}}(z^*) \otimes f_\omega(z) = \sum_{j=1}^{\infty} \sum_{m_j} c^*_{j,m_j}\ c_{j,m_j}\ (z^*_1\ z_1 - z^*_{01}\ z_{01})^j\ \cdots\ (z^*_n\ z_n - z^*_{0n}\ z'_{0n})^j$$

at the bipoint $(z^*_0 \times z_0)$, $z_0 \in \mathbb{C}^n$, such that each term of $f_{\overline{\omega}}(z^*) \otimes f_\omega(z)$ corresponds to a conjugacy class representative $g^{(2n)}_{R\times L}[j,m_j]$. And, the holomorphic bifunction $f_{\overline{\omega}}(z^*) \otimes f_\omega(z)$ constitutes an irreducible holomorphic representation $\mathrm{Irr}\,\mathrm{hol}(\mathrm{GL}_n(F_{\overline{\omega}} \times F_\omega))$ of $\mathrm{GL}_n(F_{\overline{\omega}} \times F_\omega) \approx G^{(2n)}(F_{\overline{\omega}} \times F^+_\omega)$.

0.2 Langlands irreducible global bilinear correspondences

Let $\widetilde{F}_L$ (resp. $\widetilde{F}_R$) denote a complex symmetric splitting semifield.
Then, the pseudo-ramified lattice bisemispace

$$X_{S_{R\times L}} = \mathrm{GL}_n(\widetilde{F}_R \times \widetilde{F}_L)\Big/ \mathrm{GL}_n((\mathbb{Z}\,/N\ \mathbb{Z}\,)^2) \approx G^{(2n)}(F_{\overline{\varpi}} \times F_{\omega})$$

was introduced such that its cosets are isomorphic to the conjugacy classes of $G^{(2n)}(\widetilde{F}_{\overline{\varpi}} \times \widetilde{F}_{\omega})$.
Let $\{\widetilde{F}_{\omega^1}\}$ (resp. $\{\widetilde{F}_{\overline{\varpi}^1}\}$) denote the set of irreducible extensions $\widetilde{F}_{\omega^1_j}$ (resp. $\widetilde{F}_{\overline{\varpi}^1_j}$) having a rank N . Then, the smallest pseudo-ramified normal bilinear algebraic subsemigroup $P^{(2n)}(\widetilde{F}_{\overline{\varpi}^1} \times \widetilde{F}_{\omega^1})$ of $G^{(2n)}(\widetilde{F}_{\omega} \times \widetilde{F}_{\omega})$ is introduced as being the complex bilinear equivalent of the minimal (not standard) parabolic subgroup: it corresponds to the irreducible n-dimensional complex representation of the product $I_{F_R} \times I_{F_L}$ of global inertia subgroups having an order equal to $(N \cdot m^{(j)})$ where $m^{(j)} = m_j + 1$ is the multiplicity of the j-th complex place.

Then, a toroidal compactification of $X_{S_{R\times L}}$ is envisaged by mapping $X_{S_{R\times L}}$ into the corresponding toroidal compactified lattice bisemispace

$$\overline{X}_{S_{R\times L}} = \mathrm{GL}_n(F^T_R \times F^T_L)\Big/ \mathrm{GL}_n((\mathbb{Z}\,/N\ \mathbb{Z}\,)^2) \approx G^{(2n)}(F^T_{\overline{\varpi}} \times F^T_{\omega})$$

in such a way that $X_{S_{R\times L}}$ may be viewed as the interior of $\overline{X}_{S_{R\times L}}$ in the sense of the Borel-Serre compactification. F^T_{ω} and $F^T_{\overline{\varpi}}$ are the sets of toroidal completions.
A double coset decomposition of the bilinear complete algebraic semigroup $\mathrm{GL}_n(F^T_R \times F^T_L)$ gives rise to the compactified bisemispace

$$\overline{S}^{P_n}_{K_n} = P_n(F^T_{\overline{\varpi}^1} \times F^T_{\omega^1}) \setminus \mathrm{GL}_n(F^T_R \times F^T_L)\Big/ \mathrm{GL}_n((\mathbb{Z}\,/N\ \mathbb{Z}\,)^2)\ .$$

A general bilinear cohomology is introduced in section 3.2 as a contravariant functor H^* from smooth abstract (resp. algebraic) bisemivarieties together with a natural transformation $n_{H^*\to H^{[*,*]}}$ from H^* to the associated the de Rham bilinear cohomology $H^{[*,*]}$.
This bilinear cohomology is:

- of general type in the sense that it is a motivic (or Weil) bilinear cohomology directly related to the singular, de Rham of Betti cohomologies,

- characterized by Hodge (bisemi)-cycles, a Künneth standard conjecture and a Künneth biisomorphism.

It results then that:

$$H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{R_\oplus}} \otimes M^{2i}_{T_{L_\oplus}}) = \mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus}))$$

where

- $M^{2i}_{T_{R_\oplus}} \otimes M^{2i}_{T_{L_\oplus}}$ is an irreducible $\mathrm{GL}_i(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus})$-subbisemimodule of $M^{2n}_{T_{R_\oplus}} \otimes M^{2n}_{T_{L_\oplus}}$, $i \leq n$ is a complex dimension; $2i$ is the real corresponding dimension ;
- $\mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus}))$ is the representation space of the complete bilinear semigroup of matrices $\mathrm{GL}_i(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus})$.

As the toroidal compactification of $X_{S_{R\times L}}$ is an isomorphism, we have that:

$$\begin{aligned}\mathrm{Irr}\,\mathrm{Rep}\,W^{(2i)}_{F_{R\times L}}(W^{ab}_{F_R} \times W^{ab}_{F_L}) &\simeq G^{(2i)}(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus}) \\ &\simeq \mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus}))\ .\end{aligned}$$

On the other hand, the bilinear cohomology $H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{R_\oplus}} \otimes M^{2i}_{T_{L_\oplus}})$ has a decomposition following the equivalent representatives $g^{2i}_{T_{R\times L}}[j,m_j]$ of the conjugacy classes of the complete bilinear semigroup $G^{2i}(F^T_{\overline{\omega}} \times F^T_{\omega})$ according to:

$$H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{R_\oplus}} \otimes M^{2i}_{T_{L_\oplus}}) = \bigoplus_{j=1}^{r} \bigoplus_{m_j} g^{2i}_{T_{R\times L}}[j,m_j]\ .$$

As a consequence, the bilinear cohomology $H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{R_\oplus}} \otimes M^{2i}_{T_{L_\oplus}})$ has an analytic development consisting in the product $\mathrm{EIS}_R(2i,j,m_j) \otimes \mathrm{EIS}_L(2i,j,m_j)$ of the (truncated) Fourier development of a cusp form of weight 2 by its left equivalent. The n-dimensional complex cusp biform $\mathrm{EIS}_R(2i,j,m_j) \otimes \mathrm{EIS}_L(2i,j,m_j)$, in one-to-one correspondence with the bilinear cohomology $H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{R_\oplus}} \otimes M^{2i}_{T_{L_\oplus}})$, is constructed according to a solvable way and corresponds to an eigenbifunction of the product $T_R(i;r) \otimes T_L(i;r)$, right by left, of the Hecke operators; so, $\mathrm{EIS}_R(2i,j,m_j) \otimes \mathrm{EIS}_L(2i,j,m_j)$ constitutes an irreducible supercuspidal representation $\mathrm{Irr}\,\mathrm{cusp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega}))$ of the general bilinear semigroup $\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})$.

This leads us to a Langlands irreducible global correspondence:

$$\begin{tikzcd}
\operatorname{Irr}\operatorname{Rep}^{(2i)}_{W_{F_{R\times L}}}(W^{ab}_{F_R}\times W^{ab}_{F_L}) \arrow[r] \arrow[d, equal] & \operatorname{Irr}\operatorname{cusp}(\mathrm{GL}_i(F^T_{\overline{\omega}}\times F^T_{\omega})) \arrow[d, equal] \\
G^{(2i)}(\widetilde{F}_{\overline{\omega}_\oplus}\times \widetilde{F}_{\omega_\oplus}) \arrow[d, "\wr"] & \mathrm{EIS}_{R\times L}(2i,j,m_j) \\
G^{(2i)}(F^T_{\overline{\omega}_\oplus}\times F^T_{\omega_\oplus}) \arrow[r] \arrow[d, equal] & \widehat{G}^{(2i)}(F^T_{\overline{\omega}}\times F^T_{\omega}) \arrow[u] \\
H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{R_\oplus}}\otimes M^{2i}_{T_{L_\oplus}}) &
\end{tikzcd}$$

- from the sum of the products, right by left, of the conjugacy classes (associated with the places of the semifields F^T_ω and $F^T_{\overline{\omega}}$) of the irreducible i-dimensional representation $\operatorname{Irr}\operatorname{Rep}^{(2i)}_{W_{F_{R\times L}}}(W^{ab}_{F_R}\times W^{ab}_{F_L})$ of the bilinear Weil-Deligne group given by $G^{(2i)}(F_{\overline{\omega}_\oplus}\times F_{\omega_\oplus}))$;
- to the sum of the products, right by left, of the conjugacy classes of the irreducible cuspidal representation $\operatorname{Irr}\operatorname{cusp}(\mathrm{GL}_i(F^T_{\overline{\omega}_\oplus}\times F^T_{\omega_\oplus}))$ of $\mathrm{GL}_i(F^T_{\overline{\omega}}\times F^T_\omega)$ given by $\mathrm{EIS}_R(2i,j,m_j)\otimes \mathrm{EIS}_L(2i,j,m_j)$

where $\widehat{G}^{(2i)}(F^T_{\overline{\omega}}\times F^T_\omega)$ is a (bisemi)sheaf over the complete bilinear semigroup $G^{(2i)}(F^T_{\overline{\omega}}\times F^T_\omega)$

A second kind of Langlands irreducible global correspondence can be reached by considering the real equivalent of the preceding correspondence: this can be realized by envisaging the boundary $\partial\overline{X}_{S_{R\times L}} = \mathrm{GL}_n(F^{+,T}_R\times F^{+,T}_L)\big/\mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)$ of the compactified lattice bisemispace $\overline{X}_{S_{R\times L}}$, where $F^{+,T}_L$ (resp. $F^{+,T}_R$) is a toroidal compact semifield. This inclusion morphism $\gamma^\delta_{R\times L}:\overline{X}_{S_{R\times L}}\to\partial\overline{X}_{S_{R\times L}}$ sends bijectively the (diagonal) "complex" class representatives into the corresponding "real" conjugacy class representatives covering them. This gives rise to a double coset decomposition

$$\partial\overline{S}^{P_n}_{K_n} = P_n(F^{+,T}_{\overline{v}^1}\times F^{+,T}_{v^1})\setminus \mathrm{GL}_n(F^{+,T}_R\times F^{+,T}_L)\Big/\mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)$$

of the general bilinear semigroup $\mathrm{GL}_n(F^{+,T}_R\times F^{+,T}_L)$.
$\partial\overline{S}^{P_n}_{K_n}$ is then the equivalent of a Shimura (bisemi)variety whose cohomology:

$$H^{2i}(\partial\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{\overline{v}_{R_\oplus}}}\otimes M^{2i}_{T_{v_\oplus L}}) = \mathrm{Repsp}(\mathrm{GL}_{2i}(F^{+,T}_{\overline{v}_\oplus}\times F^{+,T}_{v_\oplus}))$$

is the irreducible Eisenstein bilinear cohomology which has an analytic representation given by the product

$$\mathrm{ELLIP}_R(2i,j^\delta,m_{j^\delta})\otimes\mathrm{ELLIP}_L(2i,j^\delta,m_{j^\delta})$$

of a right $2i$-dimensional global elliptic semimodule $\mathrm{ELLIP}_R(2i,j^\delta,m_{j^\delta})$ by its left equivalent. $\mathrm{ELLIP}_R(2i,j^\delta,m_{j^\delta})\otimes\mathrm{ELLIP}_L(i,j^\delta,m_{j^\delta})$ is also a “solvable” (bi)series and is an eigenbifunction of the product of Hecke operators.
On the irreducible bisemivariey $\partial\overline{S}^{P_n}_{K_n}$, there is a Langlands irreducible global correspondence:

$$\begin{tikzcd}
\mathrm{Irr}\,\mathrm{Rep}^{(2i)}_{W_{F^+_{R\times L}}}(W^{ab}_{F^+_{\overline{v}}}\times W^{ab}_{F^+_v}) \arrow[r] \arrow[d, equal] & \mathrm{Irr}\,\mathrm{ELLIP}(\mathrm{GL}_{2i}(F^{+,T}_{\overline{v}}\times F^{+,T}_v)) \arrow[d, equal] \\
G^{(2i)}(\widetilde{F}^+_{\overline{v}_\oplus}\times\widetilde{F}^+_{v_\oplus}) \arrow[r] \arrow[d, "\wr"] & \mathrm{ELLIP}_{R\times L}(2i,j_\delta,m_{j_\delta}) \\
G^{(2i)}(F^{+,T}_{\overline{v}_\oplus}\times F^{+,T}_{v_\oplus}) \arrow[r] \arrow[d, equal] & \widehat{G}^{(2i)}(F^{+,T}_{\overline{v}}\times F^{+,T}_v) \arrow[u] \\
H^{2i}(\overline{S}^{P_n}_{K_n},M^{2i}_{T_{v_{R_\oplus}}}\otimes M^{2i}_{T_{v_{L_\oplus}}}) &
\end{tikzcd}$$

- from the sum of the products, right by left, of the conjugacy classes of the irreducible n-dimensional representation $\mathrm{Irr}\,\mathrm{Rep}^{(2i)}_{W_{F^+_{R\times L}}}(W^{ab}_{F^+_{\overline{v}}}\times W^{ab}_{F^+_v})$ of the bilinear global Weil group given by the complete bilinear real semigroup $G^{(2i)}(F^+_{\overline{v}_\oplus}\times F^+_{v_\oplus})$;
- to the sum of the products, right by left, of the conjugacy classes of the irreducible cuspidal (and elliptic) representation $\mathrm{Irr}\,\mathrm{ELLIP}(\mathrm{GL}_{2i}(F^{+,T}_{\overline{v}}\times F^{+,T}_v))$ of $\mathrm{GL}_{2i}(F^{+T}_{\overline{v}}\times F^{+T}_v)$ given by the “solvable” global elliptic bisemimodule $\mathrm{ELLIP}_R(2i,j^\delta,m_{j^\delta})$ $\otimes\,\mathrm{ELLIP}_L(2i,j^\delta,m_{j^\delta})$.

Two kinds of trace formulas have been considered. The first type of trace formula is an adaptation of the Arthur-Selberg trace formula to the operator associated with the bilinear parabolic semigroup $P^{(2n)}(F_{\overline{\varpi}^1}\times F_{\omega^1})$ envisaged in the present context as the unitary representation of the complete bilinear semigroup $\mathrm{GL}_n(F_{\overline{\omega}}\times F_\omega)$. This operator acts by convolution on the bialgebra $L^{1-1}_{R\times L}(G^{(2n)}(F^{nr}_{\overline{\omega}}\times F^{nr}_\omega))$ of smooth continuous bifunctions on the pseudo-unramified bilinear complete semigroup $G^{(2n)}(F^{nr}_{\overline{\omega}}\times F^{nr}_\omega)$ and decomposes according to the unitary conjugacy classes of the pseudo-ramified bilinear complete semigroup $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$. The resulting trace formula occurs in the bialgebra $L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}}\times F_\omega))$ of bifunctions on $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$ and relies on Lefschetz trace formula. Remark that the complete trace formula, referring to the set of irreducible representations of $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$ must be envisaged in the frame of chapter 4.

The second kind of trace formula, occurring directly in the bialgebra $L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}}\times F_\omega))$ of bifunctions on the pseudo-ramified algebraic complete semigroup $G^{(2n)}(F_{\overline{\omega}}\times F_\omega))$, leads to the Plancherel formula and corresponds to the first type of trace formula.

0.3 Langlands reducible global bilinear correspondences

While the bilinear global correspondences of Langlands deal essentially with the irreducible representations of algebraic general bilinear semigroups, reducible global bilinear correspondences will be constructed with respect to the reducibility of the representations of the considered general bilinear semigroups. Three kinds of reducibility will be envisaged:

1. the representation of the complete bilinear semigroup $\mathrm{GL}_n(F_{\overline{\omega}} \times F_\omega)$ will be said to be **partially reducible** if it decomposes according to the direct sum of irreducible bilinear representations $\mathrm{Rep}(\mathrm{GL}_{n_\ell}(F_{\overline{\omega}} \times F_\omega))$ taking into account the partition $n = n_1 + \cdots + n_\ell + \cdots + n_s$ of n .
2. the representation of $\mathrm{GL}_{2n}(F_{\overline{\omega}} \times F_\omega)$ will be said to be **orthogonally completely reducible** if it decomposes "diagonally" following the direct sum of irreducible bilinear representations $\mathrm{Rep}(\mathrm{GL}_{2_\ell}(F_{\overline{\omega}} \times F_\omega))$, $1 \le \ell \le n$.
3. the representation of $\mathrm{GL}_{2n}(F_{\overline{\omega}} \times F_\omega)$ will be said to be **nonorthogonally completely reducible** if it decomposes diagonally following the direct sum of irreducible bilinear representations $\mathrm{Rep}(\mathrm{GL}_{2_\ell}(F_{\overline{\omega}} \times F_\omega))$ and off diagonally following the direct sum of irreducible bilinear representations $\mathrm{Rep}(T^t_{2_{k_R}}(F_{\overline{\omega}}) \times T_{2_{\ell_L}}(F_\omega))$.

Let then $\overline{X}^{n=n_1+\cdots+n_s}_{S_{R\times L}}$, $\overline{X}^{2n=2_1+\cdots+2_n}_{S_{R\times L}}$ and $\overline{X}^{2n_R\times 2n_L}_{S_{R\times L}}$ be the three kinds of reducible compactified lattice bisemispaces and let $\overline{S}^{P_{n=n_1+\cdots+n_s}}_{K_{n=n_1+\cdots+n_s}}$, $\overline{S}^{P_{2n=2_1+\cdots+2_n}}_{K_{2n=2_1+\cdots+2_n}}$ and $\overline{S}^{P_{2n_R\times 2n_L}}_{K_{2n_R\times 2n_L}}$ be their corresponding double coset decompositions.
Their bilinear cohomologies then decompose according to:

- $$\begin{aligned} H^*(\overline{S}^{P_{n=n_1+\cdots+n_s}}_{K_{n=n_1+\cdots+n_s}}, \widehat{M}^{2n}_{T_{R_\oplus}} \otimes \widehat{M}^{2n}_{T_{L_\oplus}}) &= \bigoplus_{n_\ell} H^{2n_\ell}(\overline{S}^{P_n}_{K_n}, \widehat{M}^{2n_\ell}_{T_{R_\oplus}} \otimes \widehat{M}^{2n_\ell}_{T_{L_\oplus}}) \\ &= \bigoplus_{n_\ell} \big(\mathrm{EIS}_R(2n_\ell, j_R, m_{j_R}) \otimes \mathrm{EIS}_L(2n_\ell, j_L, m_{j_L})\big) \end{aligned}$$
- $$\begin{aligned} H^{2n}(\overline{S}^{P_{2n=2_1+\cdots+2_n}}_{K_{2n=2_1+\cdots+2_n}}, \widehat{M}^{2n^*}_{T_{R_\oplus}} \otimes \widehat{M}^{2n^*}_{T_{L_\oplus}}) &= \bigoplus_{\ell} H^{2_\ell}(\overline{S}^{P_{2n}}_{K_{2n}}, \widehat{M}^{2_\ell}_{T_{R_\oplus}} \otimes \widehat{M}^{2_\ell}_{T_{L_\oplus}}) \\ &= \bigoplus_{\ell} \big(\mathrm{EIS}_R(2_\ell, j_R, m_{j_R}) \otimes \mathrm{EIS}_R(2_\ell, j_L, m_{j_L})\big) \end{aligned}$$

- $$\begin{aligned} H^{2n}(\overline{S}^{P_{2n_R\times 2n_L}}_{K_{2n_R\times 2n_L}},\widehat{M}^{2n_R}_{T_{R_\oplus}}\otimes\widehat{M}^{2n_L}_{T_{L_\oplus}}) &= \underset{\ell}{\oplus}\, H^{2_\ell}(\overline{S}^{P_{2n_R\times 2n_L}}_{K_{2n_R\times 2n_L}},\widehat{M}^{2_\ell}_{T_{R_\oplus}}\otimes\widehat{M}^{2_\ell}_{T_{L_\oplus}}) \\ &\quad \underset{k_R\neq\ell_L}{\oplus} H^{2_{k_R},\ell_L}(\overline{S}^{P_{2n_R\times 2n_L}}_{K_{2n_R\times 2n_L}},\widehat{M}^{2_{k_R}}_{T_{R_\oplus}}\otimes\widehat{M}^{2_{\ell_L}}_{T_{L_\oplus}}) \\ &= \underset{\ell}{\oplus}\big(\mathrm{EIS}_R(2_\ell,j_R,m_{j_R})\otimes\mathrm{EIS}_L(2_\ell,j_L,m_{j_L})\big) \\ &\quad \underset{k_R\neq\ell_L}{\oplus}\big(\mathrm{EIS}_R(2_{k_R},j_R,m_{j_R})\otimes\mathrm{EIS}_L(2_{\ell_L},j_L,m_{j_L})\big) \end{aligned}$$

where $\widehat{M}^{2n_\ell}_{T_{R_\oplus}} \equiv \widehat{G}^{(2n_\ell)}(F^T_{\overline{\omega}_\oplus})$ (resp. $\widehat{M}^{2n_\ell}_{T_{L_\oplus}} \equiv \widehat{G}^{(2n_\ell)}(F^T_{\omega_\oplus})$) is a right (resp. left) semisheaf over the right (resp. left) linear semigroup $G^{(2n_\ell)}(F^T_{\overline{\omega}_\oplus})$ (resp. $G^{(2n_\ell)}(F^T_{\omega_\oplus})$) ,
and are in one-to-one correspondence with the sums of the products of the (truncated) Fourier developments of the cusp forms of weight two. This leads to evident reducible global bilinear correspondences of which the partially reducible case is explicitly developed here:

$$\begin{tikzcd} \mathrm{Red}(\mathrm{Rep}^{(2n)}_{W_{F_{R\times L}}}(W^{2n=2n_1+\cdots+2n_s}_{F_{\overline{\omega}}}\times W^{2n=2n_1+\cdots+2n_s}_{F_{\omega}})) \arrow[rr] \arrow[dr] & & \mathrm{Red\,cusp}(\mathrm{GL}_{n=n_1+\cdots+n_s}(F^T_{\overline{\omega}}\times F^T_{\omega})) \\ & \mathrm{CY}^{2n=2n_1+\cdots+2n_s}_T(\overline{X}_R)\times\mathrm{CY}^{2n=2n_1+\cdots+2n_s}_T(\overline{X}_L) \arrow[ur] & \end{tikzcd}$$

where:

- $\mathrm{Red}(\mathrm{Rep}^{(2n)}_{W_{F_{R\times L}}}(W^{2n=2n_1+\cdots+2n_s}_{F_{\overline{\omega}}}\times W^{2n=2n_1+\cdots+2n_s}_{F_{\omega}}))$ is given by $\widehat{G}^{(2n=2n_1+\cdots+2n_s)}(F_{\overline{\omega}_\oplus}\times F_{\omega_\oplus})$;

- $\mathrm{Red\,cusp}(\mathrm{GL}_{n=n_1+\cdots+n_s}(F^T_{\overline{\omega}}\times F^T_{\omega}))$ is given by $\underset{n_\ell}{\oplus} H^{2n_\ell}(\overline{S}^{P_{n=n_1+\cdots+n_s}}_{K_{n_1+\cdots+n_s}},\widehat{M}^{2n_\ell}_{T_{R_\oplus}}\otimes\widehat{M}^{2n_\ell}_{T_{L_\oplus}}) = \underset{n_\ell}{\oplus}(\mathrm{EIS}_R(2n_\ell,j_R,m_{j_R})\otimes\mathrm{EIS}_L(2n_\ell,j_L,m_{j_L}))$;

- $\mathrm{CY}^{2n=2n_1+\cdots+2n_s}_T(\overline{X}_L) = \underset{2n_\ell}{\oplus}\mathrm{CY}^{2n_\ell}_T(\overline{X}_L)$ with $\mathrm{CY}^{2n_\ell}_T(\overline{X}_L)$ a $2n_\ell$-dimensional toroidal compactified semi "cycle" over the (compactified) semischeme $\overline{X}_L$.

As in the irreducible case, a second kind of reducible bilinear global correspondences of Langlands has been developed in section 4.2 on the equivalents of the reducible Shimura bisemivarieties.
This last version of the paper was undertaken in order to precise and define the bilinear cohomology theory in section 3.2.

Finally, I would like to thank J. Arthur, P. Cartier, H. Carayol, P. Deligne, M. Harris, G. Henniart, H. Kim, L. Lafforgue, R. Langlands, G. Laumon and many others for helpful advices and conversations.

1 Global class field concepts, motivic chain bicomplexes and pure Chow bimotives

1.1 Global class field concepts

1.1.1 Classical background

The two main challenges of class field theory are the explicit construction of class fields and of reciprocity laws. Historically, this program proceeds essentially from the fundamental problem in algebraic number theory consisting in describing how an ordinary prime p factorizes into "primes" in the ring of integers $\mathcal{O}_E$ of finite extension E of $\mathbb{Q}$. If we set that $\mathcal{O}_E=\mathbb{Z}\,(i)$, then we obtain the following well known:

Theorem: *Suppose p is an odd prime. Then p can be written following $p=n^2+m^2=(n+im)(n-im)$, $n,m\in\mathbb{Z}$ if and only if $p\equiv 1(4)$.*

This is an elementary example of the Frobenius automorphisms Fr_p in the Galois group $G=\mathrm{Gal}(E/\mathbb{Q}\,)$. Remark that $\mathrm{Fr}_p=id$ when p splits completely into E , i.e. when the ideal it generates in $\mathcal{O}_E$ factors into distinct prime ideals of $\mathcal{O}_E$.
A major objective consists in proving that the splitting properties of p in E depend only on its residue modulo some fixed modulus N . This can be achieved if $G=\mathrm{Gal}(E/\mathbb{Q}\,)$ is abelian and if $\sigma:G\to\mathbb{C}^*$ is a homomorphism. Then, there exists an integer $N>0$ and a Dirichlet character $\chi_\sigma:(\mathbb{Z}\,/N\,\mathbb{Z}\,)^*\to\mathbb{C}^*$ such that $\sigma(\mathrm{Fr}_p)=\chi_\sigma(p)$ for all primes p unramified in E : this is E. Artin's fundamental reciprocity law of abelian class field theory. All that was developed very clearly by S. Gelbart in [Gel].

1.1.2 How do global algebraic number fields proceed from this classical background?

An interesting step of this paper in relation with section 1.1.1, consists in replacing the finite adele ring by the infinite places of an algebraic extension (i.e. a splitting (semi)field) $\widetilde{F}^+$ of a global number (semi)field k of characteristic zero such that:

a) the degrees or ranks of the generated "pseudo-ramified" completions of $\widetilde{F}^+$ are integers modulo N where N is the order of the associated global inertia subgroups.

   The completions of $\widetilde{F}_+$ are constructed from irreducible closed algebraic subsets in such a way that they are isomorphically covered by the corresponding algebraic extensions.

These completions result from an isomorphism of compactifications of the corresponding algebraic extensions.

b) a character $\chi_{v_{j_\delta}} : (\mathbb{Z}/N\,\mathbb{Z})^* \to \mathbb{C}^*$ is associated to every pseudo-ramified completion $F^+_{v_{j_\delta}}$ at the v_{j_δ}-th place of F^+ such that its decomposition group $D_{v_{j_\delta}}$ generates the Dirichlet character $\chi_{v_{j_{\delta,m_{j_\delta}}}}$ of the equivalent pseudo-ramified completion $F^+_{v_{j_{\delta,m_{j_\delta}}}}$ following:

$$D_{v_{j_\delta}}(\chi_{v_{j_\delta}}) = \chi_{v_{j_{\delta,m_{j_\delta}}}}\;, \qquad m_{j_\delta} \in \mathbb{N}\ .$$

The point a) deals with the definition of a set of increasing "pseudo-ramified" completions $F^+_{v_{j_\delta}}$ which have increasing ranks equal to the associated extension degrees $[\widetilde{F}^+_{v_{j_\delta}} : k] \simeq j_\delta \cdot N$ such that every (pseudo-)ramified completion $F^+_{v_{j_\delta}}$ is generated from an irreducible "central" completion $F^+_{v^1_{j_\delta}}$ having a rank equal to N .
And the point b) refers to the construction of equivalent pseudo-ramified completions $F^+_{v_{j_{\delta,m_{j_\delta}}}}$.
As a consequence, the residue subfield of $F^+_{v_{j_\delta}}$ is defined by $F^{+,nr}_{v_{j_\delta}} = F^+_{v_{j_\delta}} \big/ F^+_{v^1_{j_\delta}}$: this implies that this residue subfield $F^{+,nr}_{v_{j_\delta}}$ is a pseudo-unramified completion of F^+ whose rank is given by the extension degree $[\widetilde{F}^{+,nr}_{v_{j_\delta}} : k] \simeq j_\delta$ of the corresponding subfield $\widetilde{F}^{+,nr}_{j_\delta}$ of $\widetilde{F}^+$, also called a global residue degree.

The generation of global algebraic extension (semi)fields thus originates from classical algebraic number theory as it will be developed in the following sections and presents some analogy with the construction of local p-adic number fields as envisaged in the following section.

1.1.3 Classical notions of local number fields

The classical notions concerning local number fields can be summarized as follows.

Let $\mathcal{O}_K$ denote the ring of integers of a finite extension K of $\mathbb{Q}_p$. Its residue field is $k(v_K) = k(\wp_K) = \mathcal{O}_K/\wp_K$ where $\wp_K$ is the unique maximal ideal of $\mathcal{O}_K$. Let $v_K : K^* \to \mathbb{Z}$ be the unique valuation so that the absolute value on K is given by

$$|\cdot|_K = |\cdot|_{v_K} \qquad \text{with} \quad |x|_K = (\#k(v_K))^{-v_K(x)} \quad \text{for} \quad x \in K^*\ .$$

The number of elements in $k(\wp_K)$ is $q = p^f$ where $f_{v_K} = [k(v_K) : \mathbb{F}_p]$ is the residue degree over $\mathbb{Q}_p$.
The ideal $\wp_K\ \mathcal{O}_K$ of $\mathcal{O}_K$ has the form $\wp_K^{e_{v_K}} = \widetilde{\omega}_K^{e_{v_K}}$ where

- $\widetilde{\omega}_K$ is a uniformiser in $\mathcal{O}_K$;
- e_{v_K} is the ramification degree of K over $\mathbb{Q}_p$.

Then, we have $[K:\mathbb{Q}_p]=e_{v_K}\cdot f_{v_K}$ such that $e_{v_K}=[K:\mathbb{Q}_p]\big/f_{v_K}$.

If K^{nr} denotes the maximal unramified extension of K , the inertia subgroup I_K can be defined by:

$$\mathrm{Gal}(K^{ac}/K)\Big/I_K \xrightarrow{\ \sim\ } \mathrm{Gal}(K^{nr}/K) \xrightarrow{\ \sim\ } \mathrm{Gal}(k(v_K)^{ac}/k(v_K))\ .$$

1.1.4 Infinite places of a global number field of characteristic 0

Let k be a global number field of characteristic zero and let $\widetilde{F}$ denote a finite (algebraically closed) extension of k such that $\widetilde{F}=\widetilde{F}_R\cup\widetilde{F}_L$ is a symmetric splitting field composed of a right and a left algebraic extension semifields $\widetilde{F}_R$ and $\widetilde{F}_L$ in one-to-one correspondence. In the complex case, $\widetilde{F}_L$ (resp. $\widetilde{F}_R$) is the set of complex (resp. conjugate complex) simple roots of a polynomial ring $k[x]$ over k . In the real case, the symmetric splitting field is noted $\widetilde{F}^+=\widetilde{F}^+_R\cup\widetilde{F}^+_L$ where $\widetilde{F}^+_L$ (resp. $\widetilde{F}^+_R$) is the algebraic extension semifield composed of the set of positive (resp. symmetric negative) simple real roots as developed in the appendix. Remark that k can also be written following $k=k_R\cup k_L$.
The left and right equivalence classes of the local completions of $\widetilde{F}^{(+)}_L$ and $\widetilde{F}^{(+)}_R$ are the left and right real (resp. complex) infinite places of $F^{(+)}_L$ and $F^{(+)}_R$:
they are noted $v=\{v_{1_\delta},\cdots,v_{j_\delta},\cdots,v_{r_\delta}\}$ and $\overline{v}=\{\overline{v}_{1_\delta},\cdots,\overline{v}_{j_\delta},\cdots,\overline{v}_{r_\delta}\}$ in the real case and $\omega=\{\omega_1,\cdots,\omega_j,\cdots,\omega_r\}$ and $\overline{\omega}=\{\overline{\omega}_1,\cdots,\overline{\omega}_j,\cdots,\overline{\omega}_r\}$ in the complex case and are constructed in such a way that each complex place is covered by its real equivalent.

Recall that the completions of $\widetilde{F}^{(+)}_L$ and $\widetilde{F}^{(+)}_R$ at infinite places are defined for the topology by their archimedean absolute values such that the Cauchy sequences in these completions converge (i.e. have a limit).

1.1.5 Completions of a global number field

- Let F_{ω_j} (resp. $F_{\overline{\omega}_j}$) denote a left (resp. right) complex pseudo-ramified completion of $\widetilde{F}_L$ (resp. $\widetilde{F}_R$) in ω_j (resp. $\overline{\omega}_j$) and let $F^+_{v_{j_\delta}}$ (resp. $F^+_{\overline{v}_{j_\delta}}$) denote a left (resp. right) real pseudo-ramified completion of $\widetilde{F}^+_L$ (resp. $\widetilde{F}^+_R$) in v_{j_δ} (resp. $\overline{v}_{j_\delta}$): they will be assumed to be generated from an "irreducible" central completion $F_{\omega^1_j}$ (resp. $F_{\overline{\omega}^1_j}$) of rank $N\cdot m^{(j)}$ in the complex case and from a irreducible completion $F_{v^1_{j_\delta}}$ (resp.

$F_{\overline{v}^1_{j_\delta}}$) of rank N in the real case where $m^{(j)}=\sup(m_j)+1$ is the multiplicity of the j-th real completion covering its complex equivalent.

So, the ranks (i.e. the Galois extension degrees of the associated extensions) of the complex and real completions will be given by the integers modulo N , $N\in\mathbb{N}$, where the integer N corresponds to the Artin conductor.

- In the complex case, the ranks of the complex pseudo-ramified completions will be given by:

  $$[F_{\omega_j}:k]=(*+j\cdot N)\ m^{(j)} \qquad\qquad [F_{\overline{\omega}_j}:k]=(*+j\cdot N)\ m^{(j)}\ ,$$

  where $*$ is an integer inferior to N , in such a way that $[F_{\omega_j}:k]=[F_{\overline{\omega}_j}:k]=(j\cdot N)\ m^{(j)}$ if $[F_{\omega_j}:k]=[F_{\overline{\omega}_j}:k]=0\bmod N$, $1\le j\le r\le\infty$.

  The "irreducible" complex completions have ranks:

  $$[F_{\omega^1_j}:k]=N\cdot m^{(j)}\ , \qquad\qquad [F_{\overline{\omega}^1_j}:k]=N\cdot m^{(j)}\ .$$

- In the real case, the ranks of the pseudo-ramified completions will be:

  $$[F^+_{v_{j_\delta}}:k]=*+j\cdot N \qquad\qquad [F^+_{\overline{v}_{j_\delta}}:k]=*+j\cdot N\ ,$$

  in such a way that $[F^+_{v_{j_\delta}}:k]=[F^+_{\overline{v}_{j_\delta}}:k]=j\cdot N$ if $[F^+_{v_{j_\delta}}:k]=[F^+_{\overline{v}_{j_\delta}}:k]=0\bmod N$, $1\le j\le r\le\infty$.

  The irreducible complex completions have ranks:

  $$[F^+_{v^1_{j_\delta}}:k]=N\ , \qquad\qquad [F^+_{\overline{v}^1_{j_\delta}}:k]=N\ .$$

- The introduction of "irreducible" central completions $F_{\omega^1_j}$ (resp. $F_{\overline{\omega}^1_j}$) in the complex case and of irreducible central completions $F^+_{v^1_{j_\delta}}$ (resp. $F^+_{\overline{v}^1_{j_\delta}}$) in the real case leads to consider that:

  - the complex pseudo-ramified completion F_{ω_j} (resp. $F_{\overline{\omega}_j}$), $1\le j\le r$, can be cut into a set of j irreducible equivalent complex completions $F_{\omega^{j'}_j}$, $1\le j'\le j$ (resp. $F_{\overline{\omega}^{j'}_j}$) of rank $N\cdot m^{(j)}$;
  - the real pseudo-ramified completions $F^+_{v_{j_\delta}}$ (resp. $F^+_{\overline{v}_{j_\delta}}$), $1\le j_\delta\le r_\delta$, can be cut into a set of j_δ irreducible equivalent real completions $F^+_{v^{j'_\delta}_{j_\delta}}$, $1\le j'_\delta\le j_\delta$ (resp. $F^+_{\overline{v}^{j'_\delta}_{j_\delta}}$) of rank N .

However, remark that the extension degrees characterize more exactly the splitting subfields in bijection with the corresponding completions of $\widetilde{F}_L^{(+)}$ and $\widetilde{F}_R^{(+)}$.

- The introduction of "irreducible" central complex left completions $F_{\omega^1_j}$ (resp. right $F_{\overline{\omega}^1_j}$) of rank $N\cdot m^{(j)}$ at the complex places ω_j (resp. $\overline{\omega}_j$), $1\le j\le r$, and of irreducible real left completions $F^+_{v^1_{j_\delta}}$ (resp. right $F^+_{\overline{v}^1_{j_\delta}}$) of rank N at the real places v_{j_δ} (resp. $\overline{v}_{j_\delta}$), $1\le j_\delta\le r$, allow to define the global residue subfields of these completions as quotient subfields given by:

  $$F^{nr}_{\omega_j}=F_{\omega_j}\Big/F_{\omega^1_j}\qquad(\text{resp.}\quad F^{nr}_{\overline{\omega}_j}=F_{\overline{\omega}_j}\Big/F_{\overline{\omega}^1_j}\ )$$

  in the complex case and by

  $$F^{+,nr}_{v_{j_\delta}}=F^+_{v_{j_\delta}}\Big/F^+_{v^1_{j_\delta}}\qquad(\text{resp.}\quad F^{+,nr}_{\overline{v}_{j_\delta}}=F^+_{\overline{v}_{j_\delta}}\Big/F^+_{\overline{v}^1_{j_\delta}}\ )$$

  in the real case: they are called global residue completions or pseudo-unramified completions of $F_L^{(+)}$ (resp. $F_R^{(+)}$) characterized by their ranks, or global residue degrees, given respectively by

  $$[F^{nr}_{\omega_j}:k]=j\cdot m^{(j)}\qquad(\text{resp.}\quad[F^{nr}_{\overline{\omega}_j}:k]=j\cdot m[(j)\ )$$

  in the complex case and by

  $$[F^{+,nr}_{v_{j_\delta}}:k]=j\qquad(\text{resp.}\quad[F^{+,nr}_{\overline{v}_j}:k]=j\ )$$

  in the real case.

- As a place is an equivalence class of completions, we have to consider a set of complex completions $\{F_{\omega_{j,m_j}}\}_{m_j}$, $m_j\in\mathbb{N}$, equivalent to F_{ω_j} , for all $1\le j\le r$ and characterized by the same rank as F_{ω_j} .

  Similarly, a real place v_{j_δ} will be given by the basic real completion $F_{v_{j_\delta}}$ and by the set of real equivalent completions $\{F^+_{v_{j_\delta,m_{j_\delta}}}\}_{m_{j_\delta}}$, $m_{j_\delta}\in\mathbb{N}$, characterized by the same rank as $F_{v_{j_\delta}}$.

  A complex (resp. real) equivalent completion $F_{\omega_{j,m_j}}$ (resp. $F^+_{v_{j,m_{j_\delta}}}$) is generated from the respective basic completion F_{ω_j} (resp. $F^+_{v_{j_\delta}}$) by the action of the nilpotent group element $u_{\omega_{j,m_j}}$ (resp. $u_{v_{j_\delta,m_{j_\delta}}}$) following:

  $$u_{\omega_{j,m_j}}F_{\omega_j}=F_{\omega_{j,m_j}}\quad(\text{resp.}\quad u_{v_{j_\delta,m_{j_\delta}}}F^+_{v_{j_\delta}}=F^+_{v_{j_\delta,m_{j_\delta}}}\ )$$

Since only bilinear cases are relevant, it is more exactly the nilpotent group element $u_{j^2;m_j^2}$ acting on $(F_{\overline{\omega}_j} \times F_{\omega_j})$ which must be taken into account. It generates the equivalent completion $(F_{\overline{\omega}_{j;m_j}} \times F_{\omega_{j;m_j}})$ in the complex case according to:

$$u_{j^2;m_j^2}(F_{\overline{\omega}_j} \times F_{\omega_j}) = F_{\overline{\omega}_{j;m_j}} \times F_{\omega_{j;m_j}} \ ;$$

in the real case, we should have:

$$u_{j_\delta^2;m_{j_\delta^2}}(F^+_{\overline{v}_{j_\delta}} \times F^+_{v_{j_\delta}}) = F^+_{\overline{v}_{j_\delta};m_{j_\delta}} \times F^+_{v_{j_\delta};m_{j_\delta}}$$

as it will be developed in the next chapter.

1.1.6 Infinite adele semirings and semigroups $F_{\omega_\oplus}$, $F^{nr}_{\omega_\oplus}$, $F^+_{v_\oplus}$ and $F^{+,nr}_{v_\oplus}$

- So, a left (resp. right) infinite "pseudo-ramified" adele semiring $\mathbb{A}^{\infty}_{F_\omega}$ (resp. $\mathbb{A}^{\infty}_{F_{\overline{\omega}}}$) can then be introduced by considering the product over Archimedean prime places j_p of $\widetilde{F}_L$ (resp. $\widetilde{F}_R$) of the basic completions $F_{\omega_{j_p}}$ (resp. $F_{\overline{\omega}_{j_p}}$) according to

$$\mathbb{A}^{\infty}_{F_\omega} = \prod_{j_p} F_{\omega_{j_p}} \ , \quad 1 \leq j_p \leq r \leq \infty \ , \qquad\qquad (\text{resp.} \quad \mathbb{A}^{\infty}_{F_{\overline{\omega}}} = \prod_{j_p} F_{\overline{\omega}_{j_p}} \ ).$$

In the same way, a left (resp. right) "pseudo-unramified" adele semiring $\mathbb{A}^{nr,\infty}_{F_\omega}$ (resp. $\mathbb{A}^{nr,\infty}_{F_{\overline{\omega}}}$) can then be defined from the product of pseudo-unramified "prime" completions following:

$$\mathbb{A}^{nr,\infty}_{F_\omega} = \prod_{j_p} F^{nr}_{\omega_{j_p}} \qquad\qquad (\text{resp.} \quad \mathbb{A}^{nr,\infty}_{F_{\overline{\omega}}} = \prod_{j_p} F^{nr}_{\overline{\omega}_{j_p}} \ ).$$

And, in the real case, left and right pseudo-ramified and pseudo-unramified adele semirings are introduced similarly by:

$$\mathbb{A}^{\infty}_{F^+_v} = \prod_{j_{\delta_p}} F^+_{v_{j_p}} \ , \quad 1 \leq j_{\delta_p} \leq r_p \leq \infty \ , \qquad\qquad (\text{resp.} \quad \mathbb{A}^{\infty}_{F^+_{\overline{v}}} = \prod_{j_{\delta_p}} F^+_{\overline{v}_{j_{\delta_p}}} \ )$$

and by

$$\mathbb{A}^{nr,\infty}_{F^+_v} = \prod_{j_{\delta_p}} F^{+,nr}_{v_{j_{\delta_p}}} \qquad\qquad (\text{resp.} \quad \mathbb{A}^{nr,\infty}_{F^+_{\overline{v}}} = \prod_{j_{\delta_p}} F^{+,nr}_{\overline{v}_{j_{\delta_p}}} \ ).$$

- Let

$$\begin{aligned} F_\omega &= \{F_{\omega_1},\cdots,F_{\omega_j},F_{\omega_{j,m_j}},\cdots,F_{\omega_r}\} \\ (\text{resp.}\quad F_{\overline{\omega}} &= \{F_{\overline{\omega}_1},\cdots,F_{\overline{\omega}_j},F_{\overline{\omega}_{j,m_j}},\cdots,F_{\overline{\omega}_r}\}\ ) \end{aligned}$$

denote the set of pseudo-ramified completions at all Archimedean complex places ω_j (resp. $\overline{\omega}_j$). Then, the direct sum of these pseudo-ramified completions is given by:

$$\begin{aligned} F_{\omega_\oplus} &= \bigoplus_j \bigoplus_{m_j} F_{\omega_{j,m_j}}\ , \qquad 1\le j\le r\le\infty\ , \\ (\text{resp.}\quad F_{\overline{\omega}_\oplus} &= \bigoplus_j \bigoplus_{m_j} F_{\overline{\omega}_{j,m_j}}\ ). \end{aligned}$$

Similarly, if

$$\begin{aligned} F^{nr}_\omega &= \{F^{nr}_{\omega_1},\cdots,F^{nr}_{\omega_j},F^{nr}_{\omega_{j,m_j}},\cdots,F^{nr}_{\omega_r}\} \\ (\text{resp.}\quad F^{nr}_{\overline{\omega}} &= \{F^{nr}_{\overline{\omega}_1},\cdots,F^{nr}_{\overline{\omega}_j},F^{nr}_{\overline{\omega}_{j,m_j}},\cdots,F^{nr}_{\overline{\omega}_r}\}\ ) \end{aligned}$$

denote the set of corresponding pseudo-unramified completions, their direct sum is given by:

$$\begin{aligned} F^{nr}_{\omega_\oplus} &= \bigoplus_j \bigoplus_{m_j} F^{nr}_{\omega_{j,m_j}}\ , \qquad 1\le j\le r\le\infty\ , \\ (\text{resp.}\quad F^{nr}_{\overline{\omega}_\oplus} &= \bigoplus_j \bigoplus_{m_j} F^{nr}_{\overline{\omega}_{j,m_j}}\ ). \end{aligned}$$

In the real case, if

$$\begin{aligned} F^+_v &= \{F^+_{v_{1_\delta}},\cdots,F^+_{v_{j_\delta,m_{j_\delta}}},\cdots,F^+_{v_{r_\delta}}\} \\ (\text{resp.}\quad F^+_{\overline{v}} &= \{F^+_{\overline{v}_{1_\delta}},\cdots,F^+_{\overline{v}_{j_\delta,m_{j_\delta}}},\cdots,F^+_{\overline{v}_{r_\delta}}\}\ ) \end{aligned}$$

denotes the set of pseudo-ramified completions at all Armimedean real places v_{j_δ} (resp. $\overline{v}_{j_\delta}$), the direct sum of these gives:

$$\begin{aligned} F^+_{v_\oplus} &= \bigoplus_{j_\delta} \bigoplus_{m_{j_\delta}} F^+_{v_{j_\delta,m_{j_\delta}}}\ , \qquad 1\le j_\delta\le r_\delta\le\infty\ , \\ (\text{resp.}\quad F^+_{\overline{v}_\oplus} &= \bigoplus_{j_\delta} \bigoplus_{m_{j_\delta}} F^+_{\overline{v}_{j_\delta,m_{j_\delta}}}\ ). \end{aligned}$$

Similarly, the direct sum of all pseudo-unramified real completions gives:

$$\begin{aligned} F^{+,nr}_{v_\oplus} &= \bigoplus_{j_\delta} \bigoplus_{m_{j_\delta}} F^{+,nr}_{v_{j_\delta,m_{j_\delta}}}\ , \qquad 1\le j_\delta\le r_\delta\le\infty\ , \\ (\text{resp.}\quad F^{+,nr}_{\overline{v}_\oplus} &= \bigoplus_{j_\delta} \bigoplus_{m_{j_\delta}} F^{+,nr}_{\overline{v}_{j_\delta,m_{j_\delta}}}\ ). \end{aligned}$$

Galois groups and Weil groups will now be introduced in the complex case by taking into account that the real case can be handled similarly. Furthermore, the considered Galois subgroups refer to Galois subgroups of splitting subfields $\widetilde{F}_{\omega_j}$ in bijection with the corresponding completions F_{ω_j} .

1.1.7 Galois and inertia subgroups

Let $\mathrm{Gal}^D(\widetilde{F}_{\omega_j}/k)$ (resp. $\mathrm{Gal}^D(\widetilde{F}_{\overline{\omega}_j}/k)$), $\mathrm{Gal}(\widetilde{F}_{\omega_{j,m_j}}/k)$ (resp. $\mathrm{Gal}(\widetilde{F}_{\overline{\omega}_{j,m_j}}/k)$) and $\mathrm{Gal}(\widetilde{F}_{\omega_j}/k)$ (resp. $\mathrm{Gal}(\widetilde{F}_{\overline{\omega}_j}/k)$) denote respectively the Galois subgroups of the basic (non-compact) extension $\widetilde{F}_{\omega_j}$ (resp. $\widetilde{F}_{\overline{\omega}_j}$), of the m_j-th equivalent extension $\widetilde{F}_{\omega_{j,m_j}}$ (resp. $\widetilde{F}_{\overline{\omega}_{j,m_j}}$) and of the set of extensions $\{\widetilde{F}_{\omega_{j,m_j}}\}$ (resp. $\{\widetilde{F}_{\overline{\omega}_{j,m_j}}\}$) including $\widetilde{F}_{\omega_j}$ (resp. $\widetilde{F}_{\overline{\omega}_j}$). So, we have that:

$$\mathrm{Gal}(\widetilde{F}_{\omega_j}/k) = \mathrm{Gal}^D(\widetilde{F}_{\omega_j}/k) \underset{m_j}{\oplus} \mathrm{Gal}(\widetilde{F}_{\omega_{j,m_j}}/k) \; .$$

For the respective pseudo-unramified extension, we should have:

$$\mathrm{Gal}(\widetilde{F}^{nr}_{\omega_j}/k) = \mathrm{Gal}^D(\widetilde{F}^{nr}_{\omega_j}/k) \underset{m_j}{\oplus} \mathrm{Gal}(\widetilde{F}^{nr}_{\omega_{j,m_j}}/k) \; .$$

The Galois subgroup of the "irreducible" central extension $\widetilde{F}_{\omega^1_j}$ having a rank equal to $N \cdot m^{(j)}$ is obviously the global inertia subgroup $I^D_{F_{\omega_j}}$ of $\mathrm{Gal}^D(\widetilde{F}_{\omega_j}/k)$ because it can be defined by:

$$I^D_{F_{\omega_j}} = \mathrm{Gal}^D(\widetilde{F}_{\omega_j}/k)\Big/ \mathrm{Gal}^D(\widetilde{F}^{nr}_{\omega_j}/k) \; .$$

Similarly, the global inertia subgroup $I_{F_{\omega_{j,m_j}}}$ of $\mathrm{Gal}(\widetilde{F}_{\omega_{j,m_j}}/k)$ is defined by:

$$I_{F_{\omega_{j,m_j}}} = \mathrm{Gal}(\widetilde{F}_{\omega_{j,m_j}}/k)\Big/ \mathrm{Gal}(\widetilde{F}^{nr}_{\omega_{j,m_j}}/k) \; .$$

As the global inertia subgroups are of Galois type, they are all isomorphic:

$$\begin{array}{ccccccccc}
I_{F^D_{\omega_1}} & \simeq & \cdots & \simeq & I_{F^D_{\omega_j}} & \simeq & \cdots & \simeq & I_{F^D_{\omega_r}} \; , \qquad 1 \leq j \leq r \; , \\[2ex]
I_{F^D_{\omega_{1,m_1}}} & \simeq & \cdots & \simeq & I_{F^D_{\omega_{j,m_j}}} & \simeq & \cdots & \simeq & I_{F^D_{\omega_{r,m_r}}} \; ,
\end{array}$$

which has for consequence that the kernel of the map:

$$\mathrm{Gal}_j : \quad \mathrm{Gal}(\widetilde{F}_{\omega_j}/k) \longrightarrow \mathrm{Gal}(\widetilde{F}^{nr}_{\omega_j}/k)$$

is a general global inertia subgroup $I_{F_{\omega_j}}$ verifying

$$I_{F_{\omega_j}} = \mathrm{Gal}(\widetilde{F}_{\omega_j}/k)\Big/ \mathrm{Gal}(\widetilde{F}^{nr}_{\omega_j}/k) \; .$$

1.1.8 Proposition

There exists an injective nilpotent morphism

$$\mathcal{N}_{\mathrm{Gal}_{\omega_j}} : \quad \mathrm{Gal}^D(\widetilde{F}_{\omega_j}/k) \longrightarrow \mathrm{Gal}(\widetilde{F}_{\omega_j}/k)$$

from the Galois subgroup $\mathrm{Gal}^D(\widetilde{F}_{\omega_j}/k)$ *of the basic extension* $\widetilde{F}_{\omega_j}$ *to the Galois subgroup* $\mathrm{Gal}(\widetilde{F}_{\omega_j}/k)$ *of the set of extensions* $\{\widetilde{F}_{\omega_{j,m_j}}\}$ *.*

Proof: According to section 1.1.5, the extension $\widetilde{F}_{\omega_{j,m_j}}$ is generated from the basic extension $\widetilde{F}_{\omega_j}$ by the action of the nilpotent group element $u_{\omega_{j,m_j}}$ following:

$$u_{\omega_{j,m_j}} \widetilde{F}_{\omega_j} = \widetilde{F}_{\omega_{j,m_j}} \; . \qquad \blacksquare$$

1.1.9 Definition: global Weil groups

Referring to section 1.1.7, we have that:

- $\mathrm{Gal}(\widetilde{F}^{ac}_L/k) = \bigoplus\limits_{j=1}^{r} \mathrm{Gal}(\widetilde{F}_{\omega_j}/k)$;
- $\mathrm{Gal}(\widetilde{F}^{ac}_R/k) = \bigoplus\limits_{j=1}^{r} \mathrm{Gal}(\widetilde{F}_{\overline{\omega}_j}/k)$;
- $\mathrm{Gal}^D(\widetilde{F}^{ac}_L/k) = \bigoplus\limits_{j=1}^{r} \mathrm{Gal}^D(\widetilde{F}_{\omega_j}/k)$;
- $\mathrm{Gal}^D(\widetilde{F}^{ac}_R/k) = \bigoplus\limits_{j=1}^{r} \mathrm{Gal}^D(\widetilde{F}_{\overline{\omega}_j}/k)$;

where $\widetilde{F}^{ac}_L$ is the union of all finite abelian extensions $\widetilde{F}_{\omega_j}$ of F_L in $\overline{F}_L$ leading to:

- $\mathrm{Gal}(\widetilde{F}^{ac}_R/k) \times \mathrm{Gal}(\widetilde{F}^{ac}_L/k) = \bigoplus\limits_{j=1}^{r} (\mathrm{Gal}(\widetilde{F}_{\overline{\omega}_j}/k) \times \mathrm{Gal}(\widetilde{F}_{\omega_j}/k))$;
- $\mathrm{Gal}^D(\widetilde{F}^{ac}_R/k) \times \mathrm{Gal}^D(\widetilde{F}^{ac}_L/k) = \bigoplus\limits_{j=1}^{r} (\mathrm{Gal}^D(\widetilde{F}_{\overline{\omega}_j}/k) \times \mathrm{Gal}^D(\widetilde{F}_{\omega_j}/k))$.

Similarly, in the pseudo-unramified case, we get:

- $\mathrm{Gal}(\widetilde{F}^{nr}_R/k) \times \mathrm{Gal}(\widetilde{F}^{nr}_L/k) = \bigoplus\limits_{j=1}^{r} (\mathrm{Gal}(\widetilde{F}^{nr}_{\overline{\omega}_j}/k) \times \mathrm{Gal}(\widetilde{F}^{nr}_{\omega_j}/k))$;
- $\mathrm{Gal}^D(\widetilde{F}^{nr}_R/k) \times \mathrm{Gal}^D(\widetilde{F}^{nr}_L/k) = \bigoplus\limits_{j=1}^{r} (\mathrm{Gal}^D(\widetilde{F}^{nr}_{\overline{\omega}_j}/k) \times \mathrm{Gal}^D(\widetilde{F}^{nr}_{\omega_j}/k))$.

As in the p-adic case, the Weil group [Tat], [H-T] is the Galois subgroup of the elements inducing on the residue field an integer power of a Frobenius element, we shall assume that, in the characteristic zero case, the Weil group will be the Galois subgroup of the pseudo-ramified extensions characterized by extension degrees $d = 0 \bmod N \cdot m^{(j)}$.
In this respect, let $\dot{\widetilde F}_{\omega_j}$ (resp. $\dot{\widetilde F}_{\overline\varpi_j}$) denote a Galois extension characterized by a degree

$$[\dot{\widetilde F}_{\omega_j} : k] = (j\cdot N)\ m^{(j)} \qquad (\text{resp.} \quad [\dot{\widetilde F}_{\overline\varpi_j} : k] = (j\cdot N)\ m^{(j)}\ ),$$

and let $\dot{\widetilde F}^{nr}_{\omega_j}$ (resp. $\dot{\widetilde F}^{nr}_{\overline\varpi_j}$) denote the respective pseudo-unramified extension characterized by the global residue degree:

$$[\dot{\widetilde F}^{nr}_{\omega_j} : k] = j\cdot m^{(j)} \qquad (\text{resp.} \quad [\dot{\widetilde F}^{nr}_{\overline\varpi_j} : k] = j\cdot m^{(j)}\ ).$$

The sum of the Galois subgroups of such extensions is then given by:

$$\mathrm{Gal}(\dot{\widetilde F}^{ac}_L/k) = \bigoplus_{j=1}^{r} \mathrm{Gal}(\dot{\widetilde F}_{\omega_j}/k) = W^{ab}_{F_L}$$

$$(\text{resp.} \quad \mathrm{Gal}(\dot{\widetilde F}^{ac}_R/k) = \bigoplus_{j=1}^{r} \mathrm{Gal}(\dot{\widetilde F}_{\overline\varpi_j}/k) = W^{ab}_{F_R}\ )$$

and corresponds to the Weil global group $W^{ab}_{F_L}$ (resp. $W^{ab}_{F_R}$).
The product, right by left, of these Weil groups thus is:

$$W^{ab}_{F_R} \times W^{ab}_{F_L} = \mathrm{Gal}(\dot{\widetilde F}^{ac}_R/k) \times \mathrm{Gal}(\dot{\widetilde F}^{ac}_L/k) \subset \mathrm{Gal}(\widetilde F^{ac}_R/k) \times \mathrm{Gal}(\widetilde F^{ac}_L/k)\ .$$

1.1.10 Remark concerning the pseudo-ramified extensions taken into account the Weil global groups

As it was mentioned in section 1.1.9, the Weil global groups are the Galois subgroups of the pseudo-ramified extensions characterized by extension degrees $d = 0 \bmod N$: they were noted $\dot{\widetilde F}_{\omega_j}$ and $\dot{\widetilde F}_{\overline\varpi_j}$, $1 \le j \le r \le \infty$.
Since the Langlands program concerns the n-dimensional representations of Weil global groups in bijection with the corresponding automorphic representations, we shall be essentially interested by the completions corresponding to these pseudo-ramified extensions $\dot{\widetilde F}_{\omega_{j,m_j}}$ and $\dot{\widetilde F}_{\overline\varpi_{j,m_j}}$. In order to simplify the notations, we shall consider, until the end of chapter 4, these restricted completions with the notations introduced in section 1.1.6 for the general case.

Let us now introduce the Suslin-Voevodsky motivic bicomplexes and the Chow bimotives which will be especially used in chapter 4.
The proposed treatment shows how it is possible to construct enough algebraic cycles.

1.2 Suslin-Voevodsky motivic bicomplexes

Let F_ω (resp. $F_{\overline{\omega}}$) denote the set of the finite completions of F_L (resp. F_R) at the set of infinite places ω (resp. $\overline{\omega}$).

1.2.1 Definition: CW semicomplexes

Let T_L (resp. T_R) be a left (resp. right) topological semispace over F_ω (resp. $F_{\overline{\omega}}$) restricted respectively to the upper (resp. lower) half space. T_L (resp. T_R) is identified with a left (resp. right) CW complex having a partition in left (resp. right) closed cells noted $\mathrm{CEL}_L(F_{\omega_j})$ (resp. $\mathrm{CEL}_R(F_{\overline{\omega}_j})$) such that every left (resp. right) closed cell $\mathrm{CEL}_L(F_{\omega_j})$ (resp. $\mathrm{CEL}_R(F_{\overline{\omega}_j})$) be defined over a left (resp. right) place ω_j (resp. $\overline{\omega}_j$). Furthermore, it will be assumed that every left (resp. right) reducible closed cell $\mathrm{CEL}_L(F_{\omega_j})$ (resp. $\mathrm{CEL}_R(F_{\overline{\omega}_j})$) decomposes into a set of one-dimensional irreducible subcells $C_L(F_{\omega_j})$ (resp. $C_R(F_{\overline{\omega}_j})$). Each left (resp. right) CW complex T_L (resp. T_R) is isomorphic to an object of the category $Sm_L(k)$ (resp. $Sm_R(k)$) of the left (resp. right) smooth quasi-projective semivarieties over k . The Eilenberg-MacLane topological semispace generated by the left (resp. right) CW complex T_L (resp. T_R) is the free topological abelian semigroup generated by T_L (resp. T_R), i.e. the semigroup completion $L^{\mathrm{top}}[T_L]$ (resp. $L^{\mathrm{top}}[T_R]$) of the topological commutative monoid $\coprod\limits_{i_\ell\geq 0} SP^{i_\ell}(T_L)$ (resp. $\coprod\limits_{i_\ell\geq 0} SP^{i_\ell}(T_R)$), where $SP^{i_\ell}(T_L)$ (resp. $SP^{i_\ell}(T_R)$) denotes the i_ℓ-th symmetric product of T_L (resp. T_R).

1.2.2 Definition: Suslin-Voevodsky motivic presheaf

Let $\Delta_L^{2n_\ell}$ (resp. $\Delta_R^{2n_\ell}$) denote a left (resp. right) real topological $2n_\ell$-simplex, $2n_\ell\in\mathbb{N}$, i.e. a closed subscheme in the affine $(2n_\ell+2)$-semispace $\mathbb{A}_k^{2n_\ell+2}$ (resp. $\mathbb{A}_k^{2n_\ell+2}$) and let $\Delta_L^\bullet$ (resp. $\Delta_R^\bullet$) be a cosimplicial object in $Sm_R(k)$ (resp. $Sm_L(k)$) from the collection of the $\Delta_R^{2n_\ell}$ (resp. $\Delta_L^{2n_\ell}$). A presheaf of complexes on $Sm_L(k)$ (resp. $Sm_R(k)$) is a functor from $Sm_L(k)$ (resp. Sm_Rk) to the left (resp. right) chain complexes of abelian semigroups. On the other hand, let X_L^{sv} (resp. X_R^{sv}) denote a left (resp. right) Suslin-Voevodsky smooth semischeme of real dimension 2ℓ isomorphic to the left (resp. right) topological semispace T_L (resp. T_R) over F_ω (resp. $F_{\overline{\omega}}$) decomposing into closed cells $\mathrm{CEL}_L(F_{\omega_j})$ (resp. $\mathrm{CEL}_R(F_{\overline{\omega}_j})$) which are also isomorphic to their algebraic analogues $\mathrm{CEL}_L(\widetilde{F}_{\omega_j})$ (resp. $\mathrm{CEL}_R(\widetilde{F}_{\overline{\omega}_j})$). Fix $2n_\ell=i_\ell\times 2\ell$. Then, a Suslin-Voevodsky motivic left (resp. right) presheaf of X_L^{sv} (resp. X_R^{sv}) on $Sm_L(k)$ (resp. $Sm_R(k)$), denoted $\underline{C}_*(X_L^{sv})$ (resp. $\underline{C}_*(X_R^{sv})$) and called in short a Suslin-Voevodsky left (resp. right) motive, is any functor from (X_L^{sv}) (resp. (X_R^{sv})) to the left (resp. right) chain complex associated to the left (resp. right) abelian semigroup $\coprod\limits_{i_\ell} \text{Hom}_{Sm_{L/k}}(\Delta^\bullet_L, SP^{i_\ell}(X^{sv}_L))$ (resp. $\coprod\limits_{i_\ell} \text{Hom}_{Sm_{R/k}}(\Delta^\bullet_R, SP^{i_\ell}(X^{sv}_R))$).

1.2.3 Definition: reducibility of Suslin-Voevodsky presheaf

If we take into account the decomposition of the left (resp. right) smooth semischeme X^{sv}_L (resp. X^{sv}_R) in closed subsemischemes isomorphic to algebraic cells $\text{CEL}_L(F_{\omega_j})$ (resp. $\text{CEL}_R(F_{\overline{\omega}_j})$, then the Suslin-Voevodsky left (resp. right) motive $\underline{C}_*(X^{sv}_L)$ (resp. $\underline{C}_*(X^{sv}_R)$) is the functor from X^{sv}_L (resp. X^{sv}_R) to

$$\coprod_{i_\ell}\coprod_j \text{Hom}_{Sm_{L/k}}(\Delta^\bullet_L, SP^{i_\ell}_L(X^{sv}_L[j])) \qquad (\text{resp.} \quad \coprod_{i_\ell}\coprod_j \text{Hom}_{Sm_{R/k}}(\Delta^\bullet_L, SP^{i_\ell}_L(X^{sv}_R[j])) \ )$$

where $SP^{i_\ell}(X^{sv}_L[j])$ (resp. $SP^{i_\ell}(X^{sv}_R[j])$) denotes the j-th equivalence class of the i_ℓ-th symmetric product of X^{sv}_L (resp. X^{sv}_R) associated with the left (resp. right) place ω_j (resp. $\overline{\omega}_j$) of the completion of the semifield $\widetilde{F}_L$ (resp. $\widetilde{F}_R$). On the other hand, if we consider the decomposition of the closed cells into one-dimensional irreducible affine curves $C_L(j)$ (resp. $C_R(j)$), then the Suslin-Voevodsky left (resp. right) motive $\underline{C}_*(X^{sv}_L)$ (resp. $\underline{C}_*(X^{sv}_R)$) will be represented by

$$\begin{aligned} &\coprod_{i_\ell}\coprod_j \coprod_{m=1} \text{Hom}_{Sm_{L/k}}(\Delta^\bullet_L, C_m(SP^{i_\ell}(X^{sv}_L[j]))) \\ (\text{resp.} \qquad &\coprod_{i_\ell}\coprod_j \coprod_{m=1} \text{Hom}_{Sm_{R/k}}(\Delta^\bullet_R, C_m(SP^{i_\ell}(X^{sv}_R[j]))) \ ) \end{aligned}$$

where $\{C_m(SP^{i_\ell}(X^{sv}_L[j]))\}^{2n_\ell}_{m=1}$ (resp. $\{C_m(SP^{i_\ell}(X^{sv}_R[j]))\}^{2n_\ell}_{m=1}$) is a set of affine curves, having multiplicities m_{j_ℓ} . Finally, if $2n_\ell = i_\ell \times 2\ell$, let $L_{2n_\ell;L}$ (resp. $L_{2n_\ell;R}$) denote the restriction of $\underline{C}_*(X^{sv}_L)$ (resp. $\underline{C}_*(X^{sv}_R)$) to its $2n_\ell$-th element and let $Z_L(2n_\ell)$ (resp. $Z_R(2n_\ell)$) denote the $2n_\ell$-th left (resp. right) Suslin-Voevodsky submotive given by $\underline{C}_*(L_{2n_\ell;L})[-2n_\ell]$ (resp. $\underline{C}_*(L_{2n_\ell;R})[-2n_\ell]$) which corresponds to the $2n_\ell$-th desuspension of $\underline{C}_*(L_{2n_\ell;L})$ (resp. $\underline{C}_*(L_{2n_\ell;R})$) (the desuspension sends objects of homological degree $(2n_\ell+2)$ to degree $2n_\ell$) [Mor].

1.2.4 Presheaf with transfers

$\underline{C}_*(X^{sv}_L)$ (resp. $\underline{C}_*(X^{sv}_R)$) has the property to be a presheaf with transfer, i.e. has the correspondence property of motives. A left (resp. right) correspondence between the smooth left (resp. right) semischemes X^{sv}_L (resp. X^{sv}_R) and Y^{sv}_L (resp. Y^{sv}_R), noted $\text{Corr}(X^{sv}_L, Y^{sv}_L)$ (resp. $\text{Corr}(X^{sv}_R, Y^{sv}_R)$), is the free abelian semigroup on the set of closed irreducible subsemischemes XY^{sv}_L (resp. XY^{sv}_R) of $X^{sv}_L \times_k Y^{sv}_L$ (resp. $X^{sv}_R \times_k Y^{sv}_R$) [Voe] endowed with the projection from XY^{sv}_L (resp. XY^{sv}_R) on one of the irreducible components of X^{sv}_L or of Y^{sv}_L (resp. X^{sv}_R or of Y^{sv}_R).

1.2.5 Bilinear correspondence on Suslin-Voevodsky semischemes

As left and right semischemes X_L^{sv} and X_R^{sv} are taken into account, we have to introduce a new kind of correspondence called bilinear left-right (resp. right-left) correspondence between the left (resp. right) smooth semischeme X_L^{sv} (resp. X_R^{sv}) and the right (resp. left) smooth semischeme Y_R^{sv} (resp. Y_L^{sv}). Then, a bilinear left-right (resp. right-left) correspondence will be given by $\mathrm{Corr}(Y_R^{sv}, X_L^{sv})$ (resp. $\mathrm{Corr}(Y_L^{sv}, X_R^{sv})$) and defined as the free abelian bilinear semigroup on the set of closed irreducible subsemischemes $XY_{R\times L}^{sv}$ (resp. $XY_{L\times R}^{sv}$) of $Y_R^{sv}\times X_L^{sv}$ (resp. $Y_L^{sv}\times X_R^{sv}$). This leads to introduce a left (resp. right) Suslin-Voevodsky motivic bilinear presheaf $\underline{C}_*(Y_R^{sv}\times X_L^{sv})$ (resp. $\underline{C}_*(Y_L^{sv}\times X_R^{sv})$) from $Y_R^{sv}\times X_L^{sv}$ (resp. $Y_L^{sv}\times X_R^{sv}$) to the right-left (resp. left-right) chain complex associated to the bilinear semigroup

$$\begin{aligned} &\coprod_{i_\ell} \mathrm{Hom}_{Sm_R\otimes Sm_L}(\Delta^\bullet_{R\times L}, SP^{i_\ell}(\mathrm{Corr}(Y_R^{sv}, X_L^{sv})) \\ (\text{resp.}\quad &\coprod_{i_\ell} \mathrm{Hom}_{Sm_R\otimes Sm_L}(\Delta^\bullet_{L\times R}, SP^{i_\ell}(\mathrm{Corr}(Y_L^{sv}, X_R^{sv}))\ ). \end{aligned}$$

1.3 Chow motives

1.3.1 Equivalence classes of algebraic cycles

Let X_L (resp. X_R) denote a left (resp. right) smooth semischeme of complex dimension n isomorphic to an algebraic linear semigroup of the same dimension over the extensions of a field k of characteristic 0 such that

- $2n\geq 2\ell$ (2ℓ is the real dimension of the Suslin-Voevodsky semischeme X_L^{sv} (resp. X_R^{sv});
- $2n=\sum\limits_\ell i_\ell\times 2\ell=\sum\limits_\ell 2n_\ell$;
- i is an integer corresponding to the dimension $2n_\ell$ used in section 1.2;

  so, we have that: $i\equiv 2n_\ell$.

Let $Z^i(X_L)$ (resp. $Z^i(X_R)$) be the semigroup of algebraic semicycles $\mathrm{CY}^i(X_L)$ (resp. $\mathrm{CY}^i(X_R)$) of codimension i on X_L (resp. X_R). As the left (resp. right) semischeme X_L (resp. X_R) is isomorphic to an algebraic linear semigroup of the same dimension decomposing into conjugacy classes labeled by the integer j (see the introduction and section 2.4), it can be assumed that the semigroup of algebraic semicycles $Z^i(X_L)$ (resp. $Z^i(X_R)$) is partitioned into equivalence classes corresponding to the set of left (resp. right) places $\omega_1, \cdots, \omega_r$ (resp. $\overline{\omega}_1, \cdots, \overline{\omega}_r$). So, let $\mathcal{Z}^i(X_L[j])$ (resp. $\mathcal{Z}^i(X_R[j])$) denote the j-th equivalence class of the semigroup of algebraic semicycles $\mathrm{CY}^i(X_L[j])$ (resp. $\mathrm{CY}^i(X_R[j])$) associated to the place ω_j (resp. $\overline{\omega}_j$) of F_L (resp. F_R).

1.3.2 Definition: rational equivalence

On the other hand, the left (resp. right) semicycle $\mathrm{CY}^i(X_L[j])$ (resp. $\mathrm{CY}^i(X_R[j])$) must be rationally equivalent to zero, i.e. there must exist rational functions $f_{\mathrm{CY}^i_L}$ (resp. $f_{\mathrm{CY}^i_R}$) on irreducible subschemes of X_L (resp. X_R) such that $\mathrm{CY}^i(X_L[j])$ (resp. $\mathrm{CY}^i(X_R[j])$) can be decomposed into a formal sum of (Weil) divisors on $f_{\mathrm{CY}^i_L}$ (resp. $f_{\mathrm{CY}^i_R}$) [Mur]. Let $\mathcal{Z}^i_{\mathrm{rat}}(X_L[j])$ (resp. $\mathcal{Z}^i_{\mathrm{rat}}(X_R[j])$) denote the j-th equivalence class of the semigroup of algebraic semicycles of codimension i rationally equivalent to zero, i.e. $\mathcal{Z}^i_{\mathrm{rat}}(X_L[j]) = \{\mathrm{CY}^i(X_L[j]) \in \mathcal{Z}^i(X_L[j]) \mid \mathrm{CY}^i(X_L[j])$ be rationally equivalent to zero$\}$ (idem for $\mathcal{Z}^i_{\mathrm{rat}}(X_R[j])$). Thus, we have that

$$\mathrm{CH}^i(X_L) = \frac{\mathcal{Z}^i(X_L)}{\mathcal{Z}^i_{\mathrm{rat}}(X_L)} \qquad (\text{resp.} \quad \mathrm{CH}^i(X_R) = \frac{\mathcal{Z}^i(X_R)}{\mathcal{Z}^i_{\mathrm{rat}}(X_R)}\ )$$

is the i-th left (resp. right) Chow semigroup of X_L (resp. X_R) and that

$$\mathrm{CH}^i(X_L[j]) = \frac{\mathcal{Z}^i(X_L[j])}{\mathcal{Z}^i_{\mathrm{rat}}(X_L[j])} \qquad (\text{resp.} \quad \mathrm{CH}^i(X_R[j]) = \frac{\mathcal{Z}^i(X_R[j])}{\mathcal{Z}^i_{\mathrm{rat}}(X_R[j])}\ )$$

is the j-th equivalence class of the i-th left (resp. right) Chow semigroup.

1.3.3 Lemma

If a numerical equivalence is considered for the i-th left (resp. right) Chow semigroup $\mathrm{CH}^i(X_L)$ (resp. $\mathrm{CH}^i(X_R)$), then it can be partitioned according to:

$$\mathrm{CH}^i(X_L) = \bigoplus_{m=1}^{i} \bigoplus_j \mathrm{CH}^1_m(X_L[j])$$

$$\textit{(resp.} \quad \mathrm{CH}^i(X_R) = \bigoplus_{m=1}^{i} \bigoplus_j \mathrm{CH}^1_m(X_R[j])\ )$$

where $\mathrm{CH}^1_m(X_L[j])$ (resp. $\mathrm{CH}^1_m(X_R[j])$) is the j-th equivalence class of the one-dimensional simple left (resp. right) Chow semigroup $\mathrm{CH}^1_m(X_L)$ (resp. $\mathrm{CH}^1_m(X_R)$).

Proof: an algebraic semicycle $\mathrm{CY}^i(X_L) \in \mathcal{Z}^i(X_L)$ (resp. $\mathrm{CY}^i(X_R) \in \mathcal{Z}^i(X_L)$) is said to be numerically equivalent to zero if, for all algebraic semicycles $\mathrm{CY}^{2n-i}(X_L) \in \mathcal{Z}^{2n-i}(X_L)$ (resp. $\mathrm{CY}^{2n-i}(X_R) \in \mathcal{Z}^{2n-i}(X_R)$), the intersection number $\#(\mathrm{CY}^i(X_L) \cdot \mathrm{CY}^{2n-i}(X_L)) = 0$ (idem for right part).

According to U. Jannsen [Jan2], the category of motives is a semisimple abelian category if there is a numerical equivalence on the cycles of the Chow semigroup $\mathrm{CH}^i(X_L)$ (resp. $\mathrm{CH}^i(X_R)$) of the motives. So, if there is a numerical equivalence on the algebraic semicycles $\mathrm{CY}^i(X_L)$ (resp. $\mathrm{CH}^i(X_R)$), the i-th Chow semigroup can decompose as a direct sum of simple one-dimensional semigroups $\mathrm{CH}^1_m(X_L)$ (resp. $\mathrm{CH}^1_m(X_R)$). So, the numerical equivalence on algebraic cycles is stronger than rational equivalence and implies it. ■

1.3.4 Definition: reducibility of algebraic semicycles

A left (resp. right) algebraic semicycle thus decomposes following:

$$\mathrm{CY}^i(X_L) = \bigoplus_j \bigoplus_{m=1}^{i} \bigoplus_{m_{j_m}} \mathrm{CY}^1_m(X_L[j])$$

$$\text{(resp.} \qquad \mathrm{CY}^i(X_R) = \bigoplus_j \bigoplus_{m=1}^{i} \bigoplus_{m_{j_m}} \mathrm{CY}^1_m(X_R[j]) \ )$$

where m_{j_m} denotes the equivalent representatives of the m-th one-dimensional semicycle $\mathrm{CY}^1_m(X_L[j])$ (resp. $\mathrm{CY}^1_m(X_R[j])$).

1.3.5 Proposition

Let $\mathrm{CH}(X_L) = \bigoplus_{i=1}^{2n} \mathrm{CH}^i(X_L)$ *(resp.* $\mathrm{CH}(X_R) = \bigoplus_{i=1}^{2n} \mathrm{CH}^i(X_R)$ *) be the left (resp. right) Chow semiring.*
If numerical equivalence is considered for the algebraic semicycles of all codimensions, the Chow semiring $\mathrm{CH}(X_L)$ *(resp.* $\mathrm{CH}(X_R)$ *) develops according to:*

$$\mathrm{CH}(X_L) = \bigoplus_{i=1}^{2n} \bigoplus_j \bigoplus_{m=1}^{i} \mathrm{CH}^1_{i;m}(X_L[j])$$

$$\textit{(resp.} \qquad \mathrm{CH}(X_R) = \bigoplus_{i=1}^{2n} \bigoplus_j \bigoplus_{m=1}^{i} \mathrm{CH}^1_{i;m}(X_R[j]) \ )$$

and has for elements:

$$\mathrm{CY}(X_L) = \bigoplus_{i=1}^{2n} \bigoplus_j \bigoplus_{m=1}^{i} \bigoplus_{m_{j_m}} \mathrm{CY}^1_{i;m}(X_L[j])$$

$$\textit{(resp.} \qquad \mathrm{CY}(X_R) = \bigoplus_{i=1}^{2n} \bigoplus_j \bigoplus_{m=1}^{i} \bigoplus_{m_{j_m}} \mathrm{CY}^1_{i;m}(X_R[j]) \ ).$$

$\mathrm{CH}(X_L)$ *(resp.* $\mathrm{CH}(X_R)$ *) is thus a semisimple semiring (see* [D-M], [Ram] *for a recent literature on this subject).*

1.3.6 Definition

A *pure Chow motive* restricted to the left or right case considered here is commonly defined as being a pair $(X_L, \mathrm{Corr}^0(X_L,X_L))$ (resp. $(X_R, \mathrm{Corr}^0(X_R,X_R))$) consisting in a left (resp. right) smooth semischeme X_L (resp. X_R) having a decomposition in the left (resp. right) Chow semiring $\mathrm{CH}(X_L)$ (resp. $\mathrm{CH}(X_R)$) as developed in Proposition 1.3.5 and a left (resp. right) correspondence [Mur] $\mathrm{Corr}^0(X_L,X_L) = \mathrm{CH}^0(X_L,X_L)$ (resp. $\mathrm{Corr}^0(X_R,X_R) = \mathrm{CH}^0(X_R,X_R)$) which is the set of left (resp. right) projectors of X_L (resp. X_R).

1.3.7 Proposition

The category of left (resp. right) Chow semimotives is additive, semisimple and graded by weights.

Proof: results directly from Proposition 1.3.5.

1.3.8 Definition: product of Chow semirings

As in 1.2.5, a bilinear correspondence $\mathrm{Corr}(X_R,X_L)$ can be introduced and defined by $\mathrm{Corr}(X_R,X_L) = \mathrm{CH}(X_R \times X_L) \simeq \mathrm{CH}(X_R) \times \mathrm{CH}(X_L)$. This leads to introduce the product between a right and a left Chow semiring which can be developed according to:

$$\begin{aligned}
\mathrm{CH}(X_R)\times\mathrm{CH}(X_L) &= \left(\bigoplus_i \bigoplus_j \bigoplus_{m_R=1}^{i} \mathrm{CH}^1_{i;m_R}(X_R[j])\right) \times \left(\bigoplus_i \bigoplus_j \bigoplus_{i;m_L}^{i} \mathrm{CH}^1_{i;m_L}(X_L[j])\right)\\
&= \bigoplus_i \bigoplus_j \bigoplus_{m_R=m_L=m} \mathrm{CH}^1_{i;m}(X_R[j]) \times \mathrm{CH}^1_{i;m}(X_L[j])\\
&\quad \bigoplus_i \bigoplus_j \bigoplus_{m_R\neq m_L} \mathrm{CH}^1_{i;m_R}(X_R[j]) \times \mathrm{CH}^1_{i;m_L}(X_L[j])
\end{aligned}$$

where the first direct sum gives rise to a bilinear intersection pairing:

$$\mathrm{CH}^1_{i;m}(X_R[j]) \times \mathrm{CH}^1_{i;m}(X_L[j]) \to \mathrm{CH}^1_{i;m}(X_{R\times L}[j])$$

while the second direct sum yields a bilinear mixed intersection pairing:

$$\mathrm{CH}^1_{i;m_R}(X_R[j]) \times \mathrm{CH}^1_{i;m_L}(X_L[j]) \to \mathrm{CH}^0_{i;m_R;m_L}(X_{R\times L}[j])\ .$$

1.3.9 Proposition

Let $\underline{C}_*(X_L^{sv})$ *(resp.* $\underline{C}_*(X_R^{sv})$ *) be the left (resp. right) Suslin-Voevodsky reducible motivic presheaf represented by*

$$\begin{aligned} & \coprod_{i_\ell}\coprod_{j}\coprod_{m} \mathrm{Hom}_{Sm_L/k}(\Delta_L^\bullet, C_m(SP^{i_\ell}(X_L^{sv}[j]))) \\ \textit{(resp.} \quad & \coprod_{i_\ell}\coprod_{j}\coprod_{m} \mathrm{Hom}_{Sm_R/k}(\Delta_R^\bullet, C_m(SP^{i_\ell}(X_R^{sv}[j]))) \ ) \end{aligned}$$

and let $\mathrm{CH}(X_L)=\bigoplus_i\bigoplus_j\bigoplus_m \mathrm{CH}^1_{i;m}(X_L[j])$ *(resp.* $\mathrm{CH}(X_R)=\bigoplus_i\bigoplus_j\bigoplus_m \mathrm{CH}^1_{i;m}(X_R[j])$ *) be the left (resp. right) reducible Chow semiring of* X_L *(resp.* X_R *).*
Then, we can envisage the maps:

$$\begin{aligned} \mathrm{Mot}_L : \quad & \underline{C}_*(X_L^{sv}) \xrightarrow{\sim} (X_L, \mathrm{Corr}^0(X_L,X_L)) \\ \textit{(resp.} \quad Mot_R : \quad & \underline{C}_*(X_R^{sv}) \xrightarrow{\sim} (X_R, \mathrm{Corr}^0(X_R,X_R)) \ ) \end{aligned}$$

which are bijective between the pure left (resp. right) Suslin-Voevodsky semimotive and the corresponding pure left (resp. right) Chow semimotive $(X_L, \mathrm{Corr}^0(X_L,X_L))$ *(resp.* $(X_R, \mathrm{Corr}^0(X_R,X_R))$ *).*

Proof: this is a direct consequence of the developments of this chapter connecting:

- the dimension $2n$ of X_L (resp. X_R) with the dimension 2ℓ of X_L^{sv} (resp. X_R^{sv}) by the relation $2n=\sum\limits_{\ell} i_\ell \times 2\ell$;
- the reducibility of $\underline{C}_*(X_L^{sv})$ (resp. $\underline{C}_*(X_R^{sv})$) to the reducibility of $\mathrm{CH}(X_L)$ (resp. $\mathrm{CH}(X_R)$). ■

2 Nonabelian global class field concepts based on the representation of an algebraic bilinear semigroup

2.1 The general bilinear algebraic semigroup

2.1.1 From the abelian global class field theory towards a nonabelian global class field theory: a summary

In section 1.1, packets of left and right equivalent complex pseudo-ramified completions associated respectively with the left and right algebraic extension semifields $\widetilde{F}_L$ and $\widetilde{F}_R$ were introduced so that:

a) the j-th packet of the left (resp. right) complex pseudo-ramified completions is composed of the basic completion F_{ω_j} (resp. $F_{\overline{\omega}_j}$) and of the equivalent completions $F_{\omega_{j,m_j}}$ (resp. $F_{\overline{\omega}_{j,m_j}}$): these completions are characterized by a rank given by the extension degree of the associated extension:

$$[\widetilde{F}_{\omega_j} : k] = * + (j \cdot N)\ m^{(j)}\ ,$$
$$\text{(resp.}\quad [\widetilde{F}_{\overline{\omega}_j} : k] = * + (j \cdot N)\ m^{(j)}\ );$$

but, as mentioned in sections 1.1.9 and 1.1.10, the completions with ranks$[\widetilde{F}_{\omega_j} : k] = (j \cdot N)\ m^{(j)}$ (resp. $[\widetilde{F}_{\overline{\omega}_j} : k] = (j \cdot N)\ m^{(j)}$) and $[\widetilde{F}_{\omega_{j,m_j}} : k] = (j \cdot N)\ m^{(j)}$ (resp. $[\widetilde{F}_{\overline{\omega}_{j,m_j}} : k] = (j \cdot N)\ m^{(j)}$) will only be taken into account.

b) as the basic completion F_{ω_j} (resp. $F_{\overline{\omega}_j}$) is generated from an irreducible central completion $F_{\omega^1_j}$ (resp. $F_{\overline{\omega}^1_j}$), the basic completions F_{ω_j} (resp. $F_{\overline{\omega}_j}$) and the equivalent completions $F_{\omega_{j,m_j}}$ (resp. $F_{\overline{\omega}_{j,m_j}}$) will be respectively decomposed into j irreducible equivalent subcompletions $F_{\omega^{j'}_j}$, $1 \le j' \le j$, (resp. $F_{\overline{\omega}^{j'}_j}$) and $F_{\omega^{j'}_{j,m_j}}$ (resp. $F_{\overline{\omega}^{j'}_{j,m_j}}$) or rank $N \cdot m^{(j)}$.

c) a character χ_{ω_j} (resp. $\chi_{\overline{\omega}_j}$) is associated to each basic completion F_{ω_j} (resp. $F_{\overline{\omega}_j}$); and an equivalent Dirichlet character $\chi_{\omega_{j,m_j}}$ (resp. $\chi_{\overline{\omega}_{j,m_j}}$) corresponds the equivalent completion $F_{\omega_{j,m_j}}$ (resp. $F_{\overline{\omega}_{j,m_j}}$).

d) the left (resp. right) j-th packet of complex pseudo-ramified extensions corresponds to the j-th left (resp. right) place ω_j (resp. $\overline{\omega}_j$) of F_L (resp. F_R).

e) a left (resp. right) infinite adele semiring was constructed from the tower F_ω (resp. $F_{\overline{\omega}}$) of these packets of equivalent completions characterized by increasing ranks according to:

$$\mathbb{A}^{\infty}_{F_\omega} = \prod_{j_p=1} F_{\omega_{j_p}} \qquad (\text{resp.} \quad \mathbb{A}^{\infty}_{F_{\overline{\omega}}} = \prod_{j_p=1} F_{\overline{\omega}_{j_p}} \ ),$$

where:

- $F_\omega = \{F_{\omega_1}, \cdots, F_{\omega_{j,m_j}}, \cdots, F_{\omega_r}\}$ (resp. $F_{\overline{\omega}} = \{F_{\overline{\omega}_1}, \cdots, F_{\overline{\omega}_{j,m_j}}, \cdots, F_{\overline{\omega}_r}\}$);
- $F_{\omega_\oplus} = \bigoplus\limits_j \bigoplus\limits_{m_j} F_{\omega_{j,m_j}}$ (resp. $F_{\overline{\omega}_\oplus} = \bigoplus\limits_j \bigoplus\limits_{m_j} F_{\overline{\omega}_{j,m_j}}$) (see section. 1.1.6).

- So, the set of packets of left (resp. right) complex pseudo-ramified extensions covering the associated completions is a left (resp. right) affine semigroup $\mathbb{S}^1_L$ (resp. $\mathbb{S}^1_R$) whose complex fibers are one-dimensional and to which a “complex” Picard semigroup $\text{Pic}(\widetilde{F}_\omega)$ (resp. $\text{Pic}(\widetilde{F}_{\overline{\omega}})$) corresponds: $\text{Pic}(\widetilde{F}_\omega)$ (resp. $\text{Pic}(\widetilde{F}_{\overline{\omega}})$) thus is the set of r isomorphism classes of finitely generated semimodules of complex dimension 1 over $\widetilde{F}_\omega$ (resp. $\widetilde{F}_{\overline{\omega}}$).

- These few considerations relative to complex pseudo-ramified extensions (real pseudo-ramified extensions can be handled similarly) deal with the abelian class field theory. Now, the n-dimensional global Langlands program is based on nonabelian global class field theory [Lan2]. The challenge then consists in constructing the n-dimensional analog of the affine semigroup $\mathbb{S}^1_L$ (resp. $\mathbb{S}^1_R$) such that we get an injective homomorphism:

$$\begin{aligned} \sigma_L : \quad & W^{ab}_{F_L} \longrightarrow \text{GL}_n(\widetilde{F}_{\omega_\oplus}) \\ (\text{resp.} \quad \sigma_R : \quad & W^{ab}_{F_R} \longrightarrow \text{GL}_n(\widetilde{F}_{\overline{\omega}_\oplus}) \ ) \end{aligned}$$

from the Weil global group $W^{ab}_{F_L}$ (resp. $W^{ab}_{F_R}$) to $\text{GL}_n(\widetilde{F}_{\omega_\oplus})$ (resp. $\text{GL}_n(\widetilde{F}_{\overline{\omega}_\oplus})$).

More concretely, as it was justified in the introduction and in [Pie3], bilinearity, instead of linearity, will be envisaged: so, enveloping (semi)algebras $A^e = A \otimes_R A^{\text{op}}$ of a given R-(semi)algebra A , where A^{op} denotes the opposite (semi)algebra of A , as well as bisemialgebras will be considered. Thus, we are interested in the products, right by left:

  - of infinite adele semirings $\mathbb{A}^{\infty}_{F_{\overline{\omega}}} \times \mathbb{A}^{\infty}_{F_\omega}$ and of semigroups $F_{\overline{\omega}} \times F_\omega$ and $F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus}$.
  - of Picard semigroups $\text{Pic}(\widetilde{F}_{\overline{\omega}}) \times \text{Pic}(\widetilde{F}_\omega)$.
  - of affine semigroups $\mathbb{S}^1_R \times \mathbb{S}^1_L$.
  - of Weil global groups $W^{ab}_{F_R} \times W^{ab}_{F_L}$.

- The n-dimensional analog of the bilinear affine semigroup $\mathbb{S}^1_R \times \mathbb{S}^1_L$ will be a $2n$-dimensional bilinear affine semigroup which is a reductive (and semisimple) bilinear algebraic semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$ considered as generated from the product $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega) \equiv T^t_n(\widetilde{F}_{\overline{\omega}}) \times T_n(\widetilde{F}_\omega)$ of the group $T^t_n(\widetilde{F}_{\overline{\omega}})$ of lower triangular matrices with entries in $\widetilde{F}_{\overline{\omega}}$ by the group $T_n(\widetilde{F}_\omega)$ of upper triangular matrices with entries in $\widetilde{F}_\omega$. So, the algebraic bilinear affine semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$ over $\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega$, resulting from the general bilinear algebraic semigroup of matrices $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$, is a $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$-bisemimodule $\widetilde{M}_R \otimes \widetilde{M}_L$ (see the appendix).

  Recall that a reductive (resp. semisimple) group is a group having no unipotent (resp. solvable) infinite normal subgroup.

- The consideration of a bilinear algebraic semigroup is justified by the fact that a bilinear algebraic (semi)group covers its linear equivalent as it is proved in proposition 2.1.7: this leads us to take into account bialgebras and enveloping algebras.

  As the bilinear algebraic semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$ is built over $(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$, it is composed of r conjugacy classes, $1 \leq j \leq r$, having multiplicities $m^{(r)} = \sup(m_r+1)$ and corresponding to the r biplaces of $(F_{\overline{\omega}} \times F_\omega)$. Note that $m^{(r)}$ denotes the number of equivalent representatives in the r-th conjugacy class.

  The algebraic representation of the bilinear algebraic semigroup of matrices $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$ into the $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$ -bisemimodule $(\widetilde{M_R} \otimes \widetilde{M_L})$ corresponds to an algebraic morphism from $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$ into $\mathrm{GL}(\widetilde{M_R} \otimes \widetilde{M_L})$ where $\mathrm{GL}(\widetilde{M_R} \otimes \widetilde{M_L})$ denotes the group of automorphisms of $\widetilde{M_R} \otimes \widetilde{M_L}$.

  Let $\widetilde{M_{R_\oplus}} \otimes \widetilde{M_{L_\oplus}}$ be the representation space of $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}_\oplus} \times \widetilde{F}_{\omega_\oplus})$ decomposing into the direct sum of subbisemimodules representing its conjugacy classes as it will be seen in section 2.1.3.

  Then, $\mathrm{GL}(\widetilde{M_{R_\oplus}} \otimes \widetilde{M_{L_\oplus}})$ constitutes the $2n$-dimensional equivalent of the product $W^{ab}_{F_R} \times W^{ab}_{F_L}$ of the global Weil groups and the bilinear algebraic semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}_\oplus} \times \widetilde{F}_{\omega_\oplus})$ becomes naturally the $2n$-dimensional (irreducible) representation space $(\mathrm{Irr}) \operatorname{Rep}^{(2n)}_{W_{F_{R\times L}}}(W^{ab}_{F_R} \times W^{ab}_{R_L})$ of $(W^{ab}_{F_R} \times W^{ab}_{R_L})$. The isomorphisms [Bor2]:

$$
\begin{aligned}
i_{\mathrm{alg}} \quad &: \quad \mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega) \quad \longrightarrow \quad G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega) \; , \\
i_{\mathrm{aut}} \quad &: \quad \mathrm{GL}(\widetilde{M_R} \otimes \widetilde{M_L}) \quad \longrightarrow \quad \mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega) \; ,
\end{aligned}
$$

leading to the (irreducible) representation

$$(\text{Irr})\,\text{Rep}^{(2n)}_{W_{\widetilde F_{R\times L}}}(W^{ab}_{F_R}\times W^{ab}_{F_L}) : \quad \text{GL}(\widetilde M_{R_\oplus}\otimes \widetilde M_{L_\oplus}) \longrightarrow G^{(2n)}(\widetilde F_{\overline\omega_\oplus}\times \widetilde F_{\omega_\oplus}) \ ,$$

implying the injective homomorphism:

$$\sigma_R\times\sigma_L : \quad W^{ab}_{F_R}\times W^{ab}_{F_L} \longrightarrow \text{GL}_n(\widetilde F_{\overline\omega_\oplus}\times \widetilde F_{\omega_\oplus})$$

as announced precedingly.

- So, the main purpose of chapter 2 will consist in introducing the bilinear algebraic semigroup $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_\omega)$ and in showing how it can be cut into conjugacy (bi)classes under the actions of Hecke operators. On these basis, nonabelian global class field concepts will be taken up.

2.1.2 Notations

The developments of this chapter will concern complex (algebraically closed) symmetric extension semifields $\widetilde F_L$ and $\widetilde F_R$ of a global number field k , taking into account that the "real" case can be handled similarly.

Let then

$$\widetilde F_{\omega_\oplus} = \overset{r}{\underset{j=1}{\oplus}} \widetilde F_{\omega_j} \underset{j,m_j}{\oplus} \widetilde F_{\omega_{j,m_j}}$$

$$(\text{resp.}\quad \widetilde F_{\overline\omega_\oplus} = \overset{r}{\underset{j=1}{\oplus}} \widetilde F_{\overline\omega_j} \underset{j,m_j}{\oplus} \widetilde F_{\overline\omega_{j,m_j}}\ )$$

denote the sum of the left (resp. right) complex pseudo-ramified basic and equivalent decompositions of the semifield $\widetilde F_L$ (resp. $\widetilde F_R$) at the set of complex places ω_j (resp. $\overline\omega_j$) as introduced in section 1.1.6 and let

$$\widetilde F^{nr}_{\omega_\oplus} = \overset{r}{\underset{j=1}{\oplus}} \widetilde F^{nr}_{\omega_j} \underset{j,m_j}{\oplus} \widetilde F^{nr}_{\omega_{j,m_j}}$$

$$(\text{resp.}\quad \widetilde F^{nr}_{\overline\omega_\oplus} = \overset{r}{\underset{j=1}{\oplus}} \widetilde F_{\overline\omega_j} \underset{j,m_j}{\oplus} \widetilde F^{nr}_{\overline\omega_{j,m_j}}\ )$$

denote the sum of the respective pseudo-unramified extensions.

2.1.3 The general reductive bilinear algebraic semigroups

a) • If we refer to the introduction, the Gauss bilinear decomposition of the algebraic bilinear semigroup of matrices $\text{GL}_n(\widetilde F_R\times\widetilde F_L)$ over the product of the extension semifields $\widetilde{F}_R$ and $\widetilde{F}_L$ can be developed according to:

$$\begin{aligned}\mathrm{GL}_n(\widetilde{F}_R \times \widetilde{F}_L) &\equiv T_n^t(\widetilde{F}_R) \times T_n(\widetilde{F}_L)\\ &= [D_n(\widetilde{F}_R) \times UT_n^t(\widetilde{F}_R)] \times [UT_n(\widetilde{F}_L) \times D_n(\widetilde{F}_L)]\end{aligned}$$

in such a way that

$$\begin{array}{rrcl} & T_n(\widetilde{F}_L): & \widetilde{F}_L \longrightarrow & T^{(2n)}(\widetilde{F}_L)\\ (\text{resp.} & T_n^t(\widetilde{F}_R): & \widetilde{F}_R \longrightarrow & T^{(2n)}(\widetilde{F}_R)\ )\end{array}$$

can be viewed as an operator sending $\widetilde{F}_L$ (resp. $\widetilde{F}_R$) into the affine semispace $T^{(2n)}(\widetilde{F}_L)$ (resp. $T^{(2n)}(\widetilde{F}_R)$) of dimension $2n$.

If F_L (resp F_R) denotes the set of equivalence classes of the completions associated with $\widetilde{F}_L$ (resp. $\widetilde{F}_R$), then we have the commutative diagram:

$$\begin{tikzcd} \widetilde{F}_L \arrow[r, "T_n"] \arrow[d] & T^{(2n)}(\widetilde{F}_L) \arrow[d] \\ F_L \arrow[r, "T_n"] & T^{(2n)}(F_L) \end{tikzcd} \quad \left(\text{resp.} \quad \begin{tikzcd} \widetilde{F}_R \arrow[r, "T_n^t"] \arrow[d] & T^{(2n)}(\widetilde{F}_R) \arrow[d] \\ F_R \arrow[r, "T_n^t"] & T^{(2n)}(F_R) \end{tikzcd} \right)$$

where $T^{(2n)}(F_L)$ (resp. $T^{(2n)}(F_R)$) is the locally compact linear complete semigroup isomorphic to $T^{(2n)}(\widetilde{F}_L)$ (resp. $T^{(2n)}(\widetilde{F}_R)$).

- The question which arises now consists in knowing in what extent the generation of the affine semispace $T^{(2n)}(\widetilde{F}_L)$ (resp. $T^{(2n)}(\widetilde{F}_R)$) from the "action" of the group of upper (resp. lower) triangular matrices $T_n(\centerdot)$ (resp. $T_n^t(\centerdot)$) on the semifield $\widetilde{F}_L$ (resp. $\widetilde{F}_R$) corresponds to the "cross action" of the group of upper (resp. lower) unitriangular matrices $UT_n(\centerdot)$ (resp. $UT_n^t(\centerdot)$) by the group of diagonal matrices $D_n(\centerdot)$.

  The response lies on the **existence of a lattice** decomposing the semifield $\widetilde{F}_L$ (resp. $\widetilde{F}_R$) into conjugacy classes which are in one-to-one correspondence with the places of F_L (resp. F_R) in such a way that, at each place ω_j (resp. $\overline{\omega}_j$), we have a basic completion F_{ω_j} (resp. $F_{\overline{\omega}_j}$) and a set of equivalent completions $\{F_{\omega_{j,m_j}}\}_{m_j\neq0}$ (resp. $\{F_{\overline{\omega}_{j,m_j}}\}_{m_j\neq0}$) as developed in section 1.1.5.

- Let then $\widetilde{F}_{b_L}$ (resp. $\widetilde{F}_{b_R}$) denote the subfield of $\widetilde{F}_L$ (resp. $\widetilde{F}_R$) composed of the extensions which are in one-to-one correspondence with the set $\{F_{\omega_j}\}_j$ (resp. $\{F_{\overline{\omega}_j}\}_j$) of basic completions of F_L (resp. F_R) and let $\widetilde{F}_{eq_L} = \widetilde{F}_L \setminus \widetilde{F}_{b_L}$

(resp. $\widetilde{F}_{eq_R}=\widetilde{F}_R\setminus\widetilde{F}_{b_R}$) denote the subfield of $\widetilde{F}_L$ (resp. $\widetilde{F}_R$) being in one-to-one correspondence with the set of equivalent completions $\{F_{\omega_{j,m_j}}\}_{m_j\neq0}$ (resp. $\{F_{\overline{\omega}_{j,m_j}}\}_{m_j\neq0}$).

- Then, the "action" of $T_n(\centerdot)$ (resp. $T_n^t(\centerdot)$) will clearly correspond to the " cross action" of $UT_n(\centerdot)$ (resp. $UT_n^t(\centerdot)$) by $D_n(\centerdot)$ if:
  - $D_n(\centerdot)$ acts on the basic subfield $\widetilde{F}_{b_L}$ (resp. $\widetilde{F}_{b_R}$);
  - $UT_n(\centerdot)$ (resp. $UT_n^t(\centerdot)$) acts on the complementary subfield $\widetilde{F}_{eq_L}$ (resp. $\widetilde{F}_{eq_R}$).

  By this way, the diagonal group $D_n(\centerdot)$ generates an affine subsemispace $T^{(2n)}(\widetilde{F}_{b_L})\subset T^{(2n)}(\widetilde{F}_L)$ (resp. $T^{(2n)}(\widetilde{F}_{b_R})\subset T^{(2n)}(\widetilde{F}_R)$) in one-to-one correspondence with the set of basic conjugacy class representatives of $T^{(2n)}(F_L)$ (resp. $T^{(2n)}(F_R)$), generated from the set of basic completions $\{F_{\omega_j}\}_j$ (resp. $\{F_{\overline{\omega}_j}\}_j$), while the unitriagonal group $UT_n(\centerdot)$ (resp. $UT_n^t(\centerdot)$) generates an affine subspace $T^{(2n)}(\widetilde{F}_{eq_L})\subset T^{(2n)}(\widetilde{F}_L)$ (resp. $T^{(2n)}(\widetilde{F}_{eq_R})\subset T^{(2n)}(\widetilde{F}_R)$), complementary of $T^{(2n)}(\widetilde{F}_{b_L})\subset T^{(2n)}(\widetilde{F}_L)$ (resp. $T^{(2n)}(\widetilde{F}_{b_R})\subset T^{(2n)}(\widetilde{F}_R)$), in one-to-one correspondence with the set of equivalent conjugacy class representatives of $T^{(2n)}(F_L)$ (resp. $T^{(2n)}(F_R)$), generated from the set of equivalent completions $\{F_{\omega_{j,m_j}}\}_{m_j\neq0}$ (resp. $\{F_{\overline{\omega}_{j,m_j}}\}_{m_j\neq0}$).

- So, we can state more precisely that:

$$\begin{aligned} T_n(\widetilde{F}_L)&=UT_n(\widetilde{F}_{b_L})\times D_n(\widetilde{F}_{b_L})\\ \text{(resp.}\quad T_n^t(\widetilde{F}_R)&=D_n(\widetilde{F}_{b_R})\times UT_n^t(\widetilde{F}_{b_R})\ \text{)}\end{aligned}$$

  where:

$$\widetilde{F}_L=\widetilde{F}_{b_L}\cup\widetilde{F}_{eq_L}\qquad\text{(resp.}\quad \widetilde{F}_R=\widetilde{F}_{b_R}\cup\widetilde{F}_{eq_R}\ \text{)},$$

  but, the standard easier notation

$$\begin{aligned} T_n(\widetilde{F}_L)&=UT_n(\widetilde{F}_L)\times D_n(\widetilde{F}_L)\\ \text{(resp.}\quad T_n(\widetilde{F}_R)&=D_n(\widetilde{F}_R)\times UT_n^t(\widetilde{F}_R)\ \text{)}\end{aligned}$$

  will be commonly used in the following.

We can then summarize all that in the diagram:

$$\begin{tikzcd}
\widetilde{F}_{b_L} \arrow[rr] \arrow[dr] \arrow[dd, "D_n(\cdot)"] & & \widetilde{F}_{eq_L} \arrow[dl] \arrow[dd, "UT_n(\cdot)"] \\
& \widetilde{F}_L \arrow[dd, "T_n(\cdot)"] & \\
T^{(n)}(\widetilde{F}_{b_L}) \arrow[dr] & & T^{(n)}(\widetilde{F}_{eq_L}) \arrow[dl] \\
& T^{(n)}(\widetilde{F}_L) &
\end{tikzcd}$$

(the right case being handled similarly).

b)

- Let B_L and B_R be two division semialgebras of dimension $2n$ respectively over the centers $\widetilde{F}_L$ and $\widetilde{F}_R$ such that B_R be the opposite division semialgebra of B_L . Similarly, let B_ω and $B_{\overline{\omega}}$ denote two division semialgebras of dimension n respectively over F_ω and $F_{\overline{\omega}}$. If we fix an isomorphism: $B_\omega \simeq T_n(F_\omega)$ (resp. $B_{\overline{\omega}} \simeq T^t_n(F_\omega)$), where $T_n(F_\omega)$ (resp. $T^t_n(F_\omega)$) denotes the matrix algebra of upper (resp. lower) Borel triangular matrices, then we have that:

  $$B_{\overline{\omega}} \otimes B_\omega \simeq T^t_n(F_{\overline{\omega}}) \times T_n(F_\omega) \equiv \mathrm{GL}_n(F_{\overline{\omega}} \times F_\omega)\ .$$

  $\mathrm{GL}_n(F_{\overline{\omega}} \times F_\omega)$ thus is a reductive bilinear general complete semigroup of invertible $n \times n$ matrices over $F_{\overline{\omega}} \times F_\omega$ having its representation space given by the tensor product $M_R \otimes M_L$ of a right $2n$-dimensional $B_{\overline{\omega}}$-semimodule M_R by a left $2n$-dimensional B_ω-semimodule M_L so that M_L is provided with a left action of $T_n(F_\omega)$ while M_R is provided with a right action of $T^t_n(F_{\overline{\omega}})$.

- We could also consider the bilinear complete semigroup $\mathrm{GL}_n(F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus})$ over product, right by left, of sums of complex pseudo-ramified completions $F_{\overline{\omega}_\oplus} = \bigoplus\limits_j \bigoplus\limits_{m_j} F_{\overline{\omega}_{j,m_j}}$ and $F_{\omega_\oplus} = \bigoplus\limits_j \bigoplus\limits_{m_j} F_{\omega_{j,m_j}}$.

  Its representation space is given by the tensor product $M_{R_\oplus} \otimes M_{L_\oplus}$ of a right n-dimensional $T^t_n(F_{\overline{\omega}_\oplus})$-semimodule $M_{R_\oplus}$ by its left equivalent $T_n(F_{\omega_\oplus})$-semimodule $M_{L_\oplus}$ in such a way that $M_{R_\oplus} \otimes M_{L_\oplus}$ decomposes according to:

  $$M_{R_\oplus} \otimes M_{L_\oplus} = \bigoplus_j \bigoplus_{m_j} (M_{\overline{\omega}_{j,m_j}} \otimes M_{\omega_{j,m_j}})$$

  where $M_{\omega_{j,m_j}}$ (resp. $M_{\overline{\omega}_{j,m_j}}$) are $B_{\omega_{j,m_j}}$-subsemimodules (resp. $B_{\overline{\omega}_{j,m_j}}$-subsemimodules).

This decomposition is a consequence of the decomposition of $F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus}$ according to:

$$F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus} = \bigoplus_j \bigoplus_{m_j} (F_{\overline{\omega}_{j,m_j}} \times F_{\omega_{j,m_j}}) \; .$$

- Similarly, the bilinear general complete semigroup of matrices over the product $\mathbb{A}^{\infty}_{F_{\overline{\omega}}} \times \mathbb{A}^{\infty}_{F_\omega}$ of infinite adele semirings is given by:

  $$\mathrm{GL}_n(\mathbb{A}^{\infty}_{F_{\overline{\omega}}} \times \mathbb{A}^{\infty}_{F_\omega}) \equiv T^t_n(\mathbb{A}^{\infty}_{F_{\overline{\omega}}}) \times T_n(\mathbb{A}^{\infty}_{F_\omega})$$

  and verify

  $$B_{\mathbb{A}^{\infty}_{F_{\overline{\omega}}}} \otimes B_{\mathbb{A}^{\infty}_{F_\omega}} \simeq \mathrm{GL}_n(\mathbb{A}^{\infty}_{F_{\overline{\omega}}} \times \mathbb{A}^{\infty}_{F_\omega})$$

  where $B_{\mathbb{A}^{\infty}_{F_\omega}}$ (resp. $B_{\mathbb{A}^{\infty}_{F_{\overline{\omega}}}}$) is a division semialgebra of dimension n over the adele semiring $\mathbb{A}^{\infty}_{F_\omega}$ (resp. $\mathbb{A}^{\infty}_{F_{\overline{\omega}}}$).

  Its representation space is given by the tensor product $M_{R_{\circledast}} \otimes M_{L_{\circledast}}$ of the $T_n(\mathbb{A}^{\infty}_{F_\omega})$-semimodule $M_{L_{\circledast}}$ by the $T^t_n(\mathbb{A}^{\infty}_{F_{\overline{\omega}}})$-semimodule $M_{R_{\circledast}}$ which decompose respectively according to:

  $$M_{L_{\circledast}} = \prod_{j_p} M_{\omega_{j_p}} \qquad \text{and} \qquad M_{R_{\circledast}} = \prod_{j_p} M_{\overline{\omega}_{j_p}}$$

  where $M_{\omega_{j_p}}$ (resp. $M_{\overline{\omega}_{j_p}}$) is a $2n$-dimensional $B_{\omega_{j_p}}$-subsemimodule (resp. $B_{\overline{\omega}_{j_p}}$-subsemimodule) on the j_p-th Archimedean prime complex place ω_{j_p} (resp. $\overline{\omega}_{j_p}$) in the sense of section 1.1.6.

- The pseudo-unramified case can be introduced similarly. Indeed, let B^{nr}_ω and $B^{nr}_{\overline{\omega}}$ denote two division semialgebras of dimension $2n$ respectively over the sums F^{nr}_ω and $F^{nr}_{\overline{\omega}}$ of pseudo-unramified completions of the semifields $\widetilde{F}_L$ and $\widetilde{F}_R$. Fixing the isomorphisms: $B^{nr}_\omega \simeq T_n(F^{nr}_\omega)$ and $B^{nr}_{\overline{\omega}} \simeq T^t_n(F^{nr}_{\overline{\omega}})$, we get:

  $$B^{nr}_{\overline{\omega}} \otimes B^{nr}_\omega \simeq T^t_n(F^{nr}_{\overline{\omega}}) \times T_n(F^{nr}_\omega) \equiv \mathrm{GL}_n(F^{nr}_{\overline{\omega}} \times F^{nr}_\omega)$$

  and $\mathrm{GL}_n(F^{nr}_{\overline{\omega}} \times F^{nr}_\omega)$ has for representation space the tensor product $M^{nr}_R \otimes M^{nr}_L$ of a pseudo-unramified right $B^{nr}_{\overline{\omega}}$-semimodule M^{nr}_R by its left equivalent M^{nr}_L .

  The bilinear general complete semigroup of invertible matrices over the product $\mathbb{A}^{nr,\infty}_{F_{\overline{\omega}}} \times \mathbb{A}^{nr,\infty}_{F_\omega}$ of pseudo-unramified infinite adele semirings is given by $\mathrm{GL}_n(\mathbb{A}^{nr,\infty}_{F_{\overline{\omega}}} \times \mathbb{A}^{nr,\infty}_{F_\omega})$ and verifies:

  $$B_{\mathbb{A}^{nr,\infty}_{F_{\overline{\omega}}}} \otimes B_{\mathbb{A}^{nr,\infty}_{F_\omega}} \simeq \mathrm{GL}_n(\mathbb{A}^{nr,\infty}_{F_{\overline{\omega}}} \times \mathbb{A}^{nr,\infty}_{F_\omega}) \; .$$

2.1.4 Proposition

a) *Any matrix* $g_n(F_{\overline{\omega}_j}\times F_{\omega_j})\in \mathrm{GL}_n(F_{\overline{\omega}}\times F_\omega)$ *, restricted over the product* $F_{\overline{\omega}_j}\times F_{\omega_j}$ *of the completions* $F_{\overline{\omega}_j}$ *and* F_{ω_j} *, has the following* **Gauss bilinear decomposition***:*

$$g_n(F_{\overline{\omega}_j}\times F_{\omega_j}) = [(d_n(F_{\omega_j})\times d_n(F_{\overline{\omega}_j})]\times[u_n^t(F_{\omega_j}]\times u_n(F_{\overline{\omega}_j}]$$

where

- $d_n(F_{\omega_j})\in D_n(F_\omega)$ *is an element of the group* $D_n(F_\omega)$ *of diagonal matrices of order* n *.*
- $u_n(F_{\omega_j})\in UT_n(F_\omega)$ *is an element of the group* $UT_n(F_\omega)$ *of upper unitriangular matrices.*

b) *The representation space* $(M_R\otimes M_L)$ *of* $\mathrm{GL}_n(F_{\overline{\omega}}\times F_\omega)$ *is (bi)-homomorphic to the representation space* $M_{R_\oplus}\otimes M_{L_\oplus}$ *of* $\mathrm{GL}_n(F_{\overline{\omega}_\oplus}\times F_{\omega_\oplus})$ *and to the representation space* $M_{R_{\textcircled{\scriptsize$\pi$}}}\otimes M_{L_{\textcircled{\scriptsize$\pi$}}}$ *of* $\mathrm{GL}_n(\mathbb{A}^{\infty}_{F_{\overline{\omega}}}\times\mathbb{A}^{\infty}_{F_\omega})$ *as it can be figured in the commutative diagram:*

$$\begin{tikzcd}
\mathrm{GL}_n(F_{\overline{\omega}}\times F_\omega) \arrow[r] \arrow[d] & \mathrm{GL}_n(F_{\overline{\omega}_\oplus}\times F_{\omega_\oplus}) \arrow[d] \\
M_R\otimes M_L \arrow[r, "H_\oplus"] \arrow[d, "H_{\textcircled{\scriptsize$\pi$}}"] & M_{R_\oplus}\otimes M_{L_\oplus} \arrow[dl, shift left, "H_{\oplus\to\textcircled{\scriptsize$\pi$}}"] \\
M_{R_{\textcircled{\scriptsize$\pi$}}}\otimes M_{L_{\textcircled{\scriptsize$\pi$}}} \arrow[ur, shift left, "H_{\textcircled{\scriptsize$\pi$}\to\oplus}"] & \\
\mathrm{GL}_n(\mathbb{A}^{\infty}_{F_{\overline{\omega}}}\times\mathbb{A}^{\infty}_{F_\omega}) \arrow[u] &
\end{tikzcd}$$

Proof:

a) The Gauss bilinear decomposition of $g_n(F_{\overline{\omega}_j}\times F_{\omega_j})$ is an evident and natural generalization of the classical Gauss linear decomposition of the elements $g_n(F)$ of the general linear group $\mathrm{GL}_n(F)$ where F denotes a number field.

The diagonal $(B_{\overline{\omega}_j}\otimes B_{\omega_j})$-subbisemimodules $M_{\overline{\omega}_j}\otimes M_{\omega_j}$ correspond to the diagonal representation space of $D_n(F_{\overline{\omega}}\times F_\omega)$, while the off-diagonal $(B_{\overline{\omega}_{j,m_j}}\otimes B_{\omega_{j,m_j}})$-subbisemimodules $M_{\overline{\omega}_{j,m_j}}\otimes M_{\omega_{j,m_j}}$ correspond to the off-diagonal representation space of $\mathrm{GL}_n(F_{\overline{\omega}}\times F_\omega)$.

Thus, the off-diagonal subsemimodules $M_{\overline{\omega}_{j,m_j}} \otimes M_{\omega_{j,m_j}}$ are obtained from the diagonal subbisemimodules under the off-diagonal nilpotent action of $UT_n^t(F_{\overline{\omega}}) \times UT_n(F_\omega)$.

b) As $\mathrm{GL}_n(F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus})$ and $\mathrm{GL}_n(\mathbb{A}^{\infty}_{F_{\overline{\omega}}} \times \mathbb{A}^{\infty}_{F_\omega})$ are bilinear subgroups of $\mathrm{GL}_n(F_{\overline{\omega}} \times F_\omega)$ by construction according to section 2.1.3, the homomorphisms:

$$H_\oplus : \quad M_R \otimes M_L \longrightarrow M_{R_\oplus} \otimes M_{L_\oplus} \qquad \text{and} \qquad H_{\textcircled{\pi}} : \quad M_R \otimes M_L \longrightarrow M_{R_{\textcircled{\pi}}} \otimes M_{L_{\textcircled{\pi}}}$$

on their representation spaces follow directly.

On the other hand, the isomorphism

$$H_{\oplus \to \textcircled{\pi}} : \quad M_{R_\oplus} \otimes M_{L_\oplus} \longrightarrow M_{R_{\textcircled{\pi}}} \otimes M_{L_{\textcircled{\pi}}} \ ,$$

which is postutaled, results from the equality between the developments in series and in Eulerian products of the associated L-functions. ■

2.1.5 Corollary

1. *The* $\mathrm{GL}_n(F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus})$*-bisemimodule* $M_{R_\oplus} \otimes M_{L_\otimes}$ *and the* $\mathrm{GL}_n(\mathbb{A}^{\infty}_{F_{\overline{\omega}}} \times \mathbb{A}^{\infty}_{F_\omega})$*-bisemimodule* $M_{R_{\textcircled{\pi}}} \otimes M_{L_{\textcircled{\pi}}}$ *are representation spaces of the bilinear complete semigroup* $\mathrm{GL}_n(F_{\overline{\omega}} \times F_\omega)$ *.*

2. *The* $\mathrm{GL}_n(F_{\overline{\omega}} \times F_\omega)$*-bisemimodule* $M_R \otimes M_L$ *is a representation space of the bilinear algebraic semigroup* $\mathrm{GL}_n(\mathbb{A}^{\infty}_{F_{\overline{\omega}}} \times \mathbb{A}^{\infty}_{F_\omega})$ *.*

Proof:

1. Part 1. of the corollary results from the homomorphism:

$$\begin{tikzcd}
\mathrm{GL}_n(F_{\overline{\omega}} \times F_\omega) \arrow[r] & M_R \otimes M_L \arrow[r] \arrow[d] & M_{R_{\textcircled{\pi}}} \otimes M_{L_{\textcircled{\pi}}} \\
& M_{R_\oplus} \otimes M_{L_\oplus} &
\end{tikzcd}$$

2. Part 2. of the corollary results from the homomorphism:

$$\mathrm{GL}_n(\mathbb{A}^{\infty}_{F_{\overline{\omega}}} \times \mathbb{A}^{\infty}_{F_\omega}) \longrightarrow M_{R_{\textcircled{\pi}}} \otimes M_{L_{\textcircled{\pi}}} \longrightarrow M_R \otimes M_L$$

3. and so on. ■

2.1.6 Linear and bilinear algebraic (semi)groups

Let $\widetilde{F}_{\overline{\omega}-\omega}=\widetilde{F}_{\overline{\omega}}\cup\widetilde{F}_{\omega}$ denote the set of extensions decomposing the algebraically (closed) symmetric splitting field $\widetilde{F}=\widetilde{F}_R\cup\widetilde{F}_L$.
Let $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}-\omega})$ denote the group of invertible $n\times n$ matrices with entries in $\widetilde{F}_{\overline{\omega}-\omega}$. Then, the algebraic general linear group $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}-\omega})$ has for representation space a vectorial space W of dimension $4n^2$ isomorphic to $(\widetilde{F}_{\overline{\omega}-\omega})^{4n^2}$ and has the Gauss decomposition:

$$g_n(\widetilde{F}_{\overline{\omega}-\omega_j})=(u_n^t(\widetilde{F}_{\overline{\omega}-\omega_j})\times u_n(\widetilde{F}_{\overline{\omega}-\omega_j}))\times d_n(\widetilde{F}_{\overline{\omega}-\omega_j})]$$

for any matrix $g_n(\widetilde{F}_{\overline{\omega}-\omega_j})\in\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}-\omega})$.
On the other hand, let $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_{\omega})$ be the bilinear algebraic semigroup as introduced in section 2.1.3. It has the Gauss bilinear decomposition introduced in proposition 2.1.4.
Then, the linear algebraic group $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}-\omega})$ and the bilinear algebraic semigroup are in one-to-one correspondence under the conditions of proposition 2.1.7.

2.1.7 Proposition

Let $\widetilde{F}_{\overline{\omega}-\omega}=\widetilde{F}_{\overline{\omega}}\cup\widetilde{F}_{\omega}$ *denote the set of extensions decomposing the algebraically closed symmetric splitting field* $\widetilde{F}=\widetilde{F}_R\cup\widetilde{F}_L$.
Let $g_n(\widetilde{F}_{\overline{\omega}-\omega_j})=(u_n^t(\widetilde{F}_{\overline{\omega}-\omega_j})\times u_n(\widetilde{F}_{\overline{\omega}-\omega_j}))\times d_n(\widetilde{F}_{\overline{\omega}-\omega_j})$ *be the Gauss decomposition of the matrix* $g_n(\widetilde{F}_{\overline{\omega}-\omega_j})$ *of the linear algebraic group* $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}-\omega_j})$.
And, let

$$g_n(\widetilde{F}_{\overline{\omega}_j}\times\widetilde{F}_{\omega_j})=[(d_n(\widetilde{F}_{\overline{\omega}_j})\times d_n(\widetilde{F}_{\omega_j})]\times[u_n^t(\widetilde{F}_{\overline{\omega}_j})\times u_n(\widetilde{F}_{\omega_j})]$$

be the Gauss decomposition of the matrix $g_n(\widetilde{F}_{\overline{\omega}_j}\times\widetilde{F}_{\omega_j})$ *of the bilinear algebraic semigroup* $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_{\omega})$.
Then, if we take into account the maps:

a) $u_n(\widetilde{F}_{\overline{\omega}-\omega_j})\xrightarrow{\sim}u_n(\widetilde{F}_{\omega_j})$,

b) $u_n^t(\widetilde{F}_{\overline{\omega}-\omega_j})\xrightarrow{\sim}u_n^t(\widetilde{F}_{\overline{\omega}_j})$,

c) $d_n(\widetilde{F}_{\overline{\omega}-\omega_j})\longrightarrow d_n(\widetilde{F}_{\overline{\omega}_j})\times d_n(\widetilde{F}_{\omega_j})$,

1. *the linear algebraic group* $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}-\omega})$ *is in one-to-one correspondence with the bilinear algebraic semigroup* $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_{\omega})$ *;*

2. *the* $4n^2$*-dimensional representation space* W *of* $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}-\omega})$ *coincides with the* $4n^2$*-dimensional representation space* $\widetilde{M}_R\otimes\widetilde{M}_L$ *of* $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_{\omega})$.

Proof:

1. the maps a) and b) send the unipotent matrices $u_n(\widetilde{F}_{\overline{\omega}-\omega_j})$ and $u_n^t(\widetilde{F}_{\overline{\omega}-\omega_j})$, with entries in the completions of the symmetric splitting field $\widetilde{F}$, respectively in the unipotent matrices $u_n(\widetilde{F}_{\omega_j})$, with entries in the extensions $\widetilde{F}_{\omega_j}$, restricted to the upper half space, and in the transposed unipotent matrices $u_n^t(\widetilde{F}_{\overline{\omega}_j})$, with entries in the extensions $\widetilde{F}_{\overline{\omega}_j}$, restricted to the lower half space.

2. the map c) sends the centralizer $d_n(\widetilde{F}_{\overline{\omega}-\omega_j})$ of $g_n(\widetilde{F}_{\overline{\omega}-\omega_j}) \in \mathrm{GL}_n(\widetilde{F}_{\overline{\omega}-\omega})$ into the "bilinear" centralizer $d_n(\widetilde{F}_{\overline{\omega}_j} \times \widetilde{F}_{\omega_j}) = d_n(\widetilde{F}_{\overline{\omega}_j}) \times d_n(\widetilde{F}_{\omega_j})$ of the matrix $g_n(\widetilde{F}_{\overline{\omega}_j} \times \widetilde{F}_{\omega_j}) \in \mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$.

3. Consequently, the basis of the $4n^2$-dimensional representation space W of the "linear" algebraic group $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}-\omega})$ must correspond to the basis of the $4n^2$-dimensional representation space $\widetilde{M}_R \otimes \widetilde{M}_L$ of the bilinear algebraic semigroup $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$, which implies that $W \simeq \widetilde{M}_R \otimes \widetilde{M}_L$. This means that the linear algebraic group $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}-\omega})$ is in fact in one-to-one correspondence with the bilinear algebraic semigroup $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$. ∎

2.2 Lattices and Hecke algebras

2.2.1 Maximal orders and lattices

Fix the maximal orders $\mathcal{O}_{\widetilde{F},\omega}$ of $\widetilde{F}_{\omega}$, $\mathcal{O}_{\widetilde{F},\overline{\omega}}$ of $\widetilde{F}_{\overline{\omega}}$, $\mathcal{O}_{\widetilde{F}^{nr},\omega}$ of $\widetilde{F}_{\omega}^{nr}$ and $\mathcal{O}_{\widetilde{F}^{nr},\overline{\omega}}$ of $\widetilde{F}_{\overline{\omega}}^{nr}$. Then, the equivalent of the maximal order Λ_{ω} (resp. $\Lambda_{\overline{\omega}}$) in the division semialgebra B_{ω} (resp. $B_{\overline{\omega}}$) is a pseudo-ramified $\mathbb{Z}/N\,\mathbb{Z}$-lattice in the left (resp. right) B_{ω} (resp. $B_{\overline{\omega}}$)-semimodule M_L (resp. M_R). Similarly, the equivalent of the maximal order Λ_{ω}^{nr} (resp. $\Lambda_{\overline{\omega}}^{nr}$) in B_{ω}^{nr} (resp. $B_{\overline{\omega}}^{nr}$) is a (pseudo-)unramified $\mathbb{Z}$-lattice in the (pseudo-)unramified left (resp. right) B_{ω}^{nr} (resp. $B_{\overline{\omega}}^{nr}$)-semimodule M_L^{nr} (resp. M_R^{nr}).

Considering the decomposition of F_{ω} and $F_{\overline{\omega}}$ into pseudo-ramified completions and of F_{ω}^{nr} and $F_{\overline{\omega}}^{nr}$ into pseudo-unramified completions according to section 2.1.2, we have the expected decomposition of the pseudo-ramified $\mathbb{Z}/N\,\mathbb{Z}$-lattice Λ_{ω} (resp. $\Lambda_{\overline{\omega}}$) into:

$$\Lambda_{\omega} = \bigoplus_{j} \bigoplus_{m_j} \Lambda_{\omega_{j,m_j}} \qquad \left(\text{resp.} \quad \Lambda_{\overline{\omega}} = \bigoplus_{j} \bigoplus_{m_j} \Lambda_{\overline{\omega}_{j,m_j}} \right)$$

where $\Lambda_{\omega_{j,m_j}} \simeq \mathcal{O}_{B_{\omega_{j,m_j}}}$ is a (pseudo-)ramified sublattice in the $B_{\omega_{j,m_j}}$-subsemimodule $M_{\omega_{j,m_j}}$.

In the (pseudo-)unramified case, we have the similar decomposition of the pseudo-unramified $\mathbb{Z}$-lattice Λ^{nr}_{ω} (resp. $\Lambda^{nr}_{\overline{\omega}}$) into:

$$\Lambda^{nr}_{\omega}=\underset{j}{\oplus}\underset{m_j}{\oplus}\Lambda^{nr}_{\omega_{j,m_j}}\qquad(\text{resp.}\quad \Lambda^{nr}_{\overline{\omega}}=\underset{j}{\oplus}\underset{m_j}{\oplus}\Lambda^{nr}_{\overline{\omega}_{j,m_j}}\ ).$$

On the other hand, we can fix the isomorphisms:

$$\begin{array}{lcccccc} & \Lambda_{\omega} & \simeq & T_n(\mathcal{O}_{F,\omega}) & (\text{resp.} & \Lambda_{\overline{\omega}} & \simeq & T^t_n(\mathcal{O}_{F,\overline{\omega}})\ ) \\ \text{and} & \Lambda^{nr}_{\omega} & \simeq & T_n(\mathcal{O}_{F^{nr},\omega}) & (\text{resp.} & \Lambda^{nr}_{\overline{\omega}} & \simeq & T^t_n(\mathcal{O}_{F^{nr},\overline{\omega}})\ ) \end{array}$$

for the pseudo-ramified and pseudo-unramified lattices.

2.2.2 Proposition

Let Λ_{ω} and $\Lambda_{\overline{\omega}}$ be pseudo-ramified $\mathbb{Z}\big/N\ \mathbb{Z}$-lattices and Λ^{nr}_{ω} and $\Lambda^{nr}_{\overline{\omega}}$ be the corresponding pseudo-unramified $\mathbb{Z}$-lattices as introduced in section 2.2.1. Then, the pseudo-ramified bilattice $\Lambda_{\overline{\omega}}\otimes\Lambda_{\omega}$ in the $B_{\overline{\omega}}\otimes B_{\omega}$-bisemimodule $M_R\otimes M_L$

- *verifies:*

$$\begin{aligned}\Lambda_{\overline{\omega}}\otimes\Lambda_{\omega}&\simeq T^t_n(\mathcal{O}_{F,\overline{\omega}})\times T_n(\mathcal{O}_{F,\omega})\\ &\simeq \mathrm{GL}_n(\mathcal{O}_{F,\overline{\omega}}\times\mathcal{O}_{F,\omega})\ .\end{aligned}$$

- *has the decomposition:*

$$\Lambda_{\overline{\omega}}\otimes\Lambda_{\omega}=\underset{j}{\oplus}\underset{m_j}{\oplus}(\Lambda_{\overline{\omega}_{j,m_j}}\otimes\Lambda_{\omega_{j,m_j}})$$

  into subbilattices $\Lambda_{\overline{\omega}_{j,m_j}}\otimes\Lambda_{\omega_{j,m_j}}$.

And the pseudo-unramified bilattice $\Lambda^{nr}_{\overline{\omega}}\otimes\Lambda^{nr}_{\omega}$ in the $B^{nr}_{\overline{\omega}}\otimes B^{nr}_{\omega}$-bisemimodule $M^{nr}_R\otimes M^{nr}_L$

- *satisfies:*

$$\begin{aligned}\Lambda^{nr}_{\overline{\omega}}\otimes\Lambda^{nr}_{\omega}&\simeq T^t_n(\mathcal{O}_{F^{nr},\overline{\omega}})\times T_n(\mathcal{O}_{F^{nr},\omega})\\ &\simeq \mathrm{GL}_n(\mathcal{O}_{F^{nr},\overline{\omega}}\times\mathcal{O}_{F^{nr},\omega})\ .\end{aligned}$$

- *has the decomposition:*

$$\Lambda^{nr}_{\overline{\omega}}\otimes\Lambda^{nr}_{\omega}=\underset{j}{\oplus}\underset{m_j}{\oplus}(\Lambda^{nr}_{\overline{\omega}_{j,m_j}}\otimes\Lambda^{nr}_{\omega_{j,m_j}})\ .$$

2.2.3 Proposition

Consider that $\mathrm{GL}_n(\mathcal{O}_{F,\overline{\omega}}\times\mathcal{O}_{F,\omega})$ *has the following Gauss bilinear decomposition:*

$$\mathrm{GL}_n(\mathcal{O}_{F,\overline{\omega}}\times\mathcal{O}_{F,\omega})=[D_n(\mathcal{O}_{F,\omega})\times D_n(\mathcal{O}_{F,\overline{\omega}})]\times[UT_n(\mathcal{O}_{F,\omega})\times UT_n^t(\mathcal{O}_{F,\overline{\omega}})]$$

(see section 2.1.3).
Then, the nilpotent action of $[UT_n(\mathcal{O}_{F,\omega})\times UT_n^t(\mathcal{O}_{F,\overline{\omega}})]$ *on* $[D_n(\mathcal{O}_{F,\omega})\times D_n(\mathcal{O}_{F,\overline{\omega}})]$ *generates the off-diagonal subbilattices* $\Lambda_{\overline{\omega}_{j,m_j}}\otimes\Lambda_{\omega_{j,m_j}}$ *equivalent to the basic subbilatice* $\Lambda_{\overline{\omega}_j,m_0}\otimes\Lambda_{\omega_j,m_0}\equiv\Lambda_{\overline{\omega}_j}\otimes\Lambda_{\omega_j}$ *for every* $1\le j\le r$ *.*

Proof: This is a direct consequence of the generation of equivalent completions $F_{\overline{\omega}_{j,m_j}}\times F_{\omega_{j,m_j}}$ from the basic completion $F_{\overline{\omega}_j}\otimes F_{\omega_j}$ by the action of the nilpotent group element $u_{\omega_j^2,m_j^2}$ as developed in section 1.1.5. ■

2.2.4 Definition: global decomposition groups and Frobenius automorphisms

1. Let $g_n(\mathcal{O}_{\overline{\omega}_j}\times\mathcal{O}_{\omega_j})=[d_n(\mathcal{O}_{\omega_j})\cdot d_n(\mathcal{O}_{\overline{\omega}_j})][u_n(\mathcal{O}_{\omega_j})\cdot u_n(\mathcal{O}_{\omega_j})^t]$ denote the j-th component of $\mathrm{GL}_n(\mathcal{O}_{F,\overline{\omega}}\times\mathcal{O}_{F,\omega})$.

   The nilpotent part $[u_n(\mathcal{O}_{\omega_j})\cdot u_n(\mathcal{O}_{\overline{\omega}_j})^t]$ of $g_n(\mathcal{O}_{\overline{\omega}_j}\times\mathcal{O}_{\omega_j})$ is interpreted in the pseudo-ramified case as the element of the decomposition group of the j-th bisublattice in $(\widetilde{M}_R\otimes\widetilde{M}_L)$. The decomposition group element referring to the (j,m_j)-th bisublattice is denoted $D_{j^2;m_{j^2}}$ and has a representation given by

   $$D_{j^2;m_{j^2}}=u_{n;m_j}(\mathcal{O}_{\omega_j})\cdot u_{n;m_j}(\mathcal{O}_{\overline{\omega}_j})^t\in UT_n(\mathcal{O}_{F,\omega})\times UT_n^t(\mathcal{O}_{F,\overline{\omega}})\ .$$

   $D_{j^2;m_{j^2}}$ acts on the split Cartan subgroup element $\alpha_{n;j^2}=d_n(\mathcal{O}_{\overline{\omega}_j})\cdot d_n(\mathcal{O}_{\omega_j})$ and is unimodular.

   Let $\det(D_{j^2;m_{j^2}}\cdot\alpha_{n;j^2})_{ss}\simeq(\#\omega_j^{2n})\cdot N^{2n}\cdot(m^{(j)})^{2n}=j^{2n}\cdot N^{2n}\cdot(m^{(j)})^{2n}$ be the determinant of the semisimple form of $D_{j^2;m_{j^2}}\cdot\alpha_{n;j^2}$ where:

   - $2n$ is the dimension;
   - the cardinality $\#\omega_j^2=\#\overline{\omega}_j\times\#\omega_j$ of the j-th quadratic place ω_j^2 is equal to $\#\omega_j^2=f_{\overline{\omega}_j}\cdot f_{\omega_j}=j^2$;
   - $(m^{(j)}\cdot N)^{2n}$ is the rank of an irreducible quadratic complex completion according to section 1.1.4.

   Indeed, $\Lambda_\omega\simeq\mathcal{O}_{F,\omega}^{2n}$ and $\Lambda_{\overline{\omega}}\simeq\mathcal{O}_{F,\overline{\omega}}^{2n}$.

Then, the unimodularity of the decomposition group $D_{j^2;m_{j^2}}$ leads to:

$$D_{j^2;m_{j^2}} : \det(\alpha_{2n;j^2}) \xrightarrow[-]{-} \det(D_{j^2;m_{j^2}} \cdot \alpha_{2n;j^2})_{ss} \ .$$

2. in the pseudo-unramified case, i.e. when $N=1$, we have that
    - $D_{j^2;m_{j^2}} \equiv \mathrm{Frob}_{j^2;m_{j^2}} = u_{n;m_j}(\mathcal{O}^{nr}_{\omega_j}) \cdot u_{n;m_j}(\mathcal{O}^{nr}_{\overline{\omega}_j})^t \in UT_n(\mathcal{O}_{F^{nr},\omega}) \times UT^t_n(\mathcal{O}_{F^{nr},\overline{\omega}})$;
    - $\det(\mathrm{Frob}_{j^2;m_{j^2}} \cdot \alpha^{nr}_{2n;j^2}) = j^{2n}$, where $\alpha^{nr}_{2n;j^2} = d_n(\mathcal{O}^{nr}_{\omega_j}) \cdot d_n(\mathcal{O}^{nr}_{\overline{\omega}_j})$;
    - $\mathrm{Frob}_{j^2;m_{j^2}} : \det(\alpha^{nr}_{n;j^2}) \xrightarrow[-]{-} \det(\alpha^{nr}_{n;j^2} \cdot \mathrm{Frob}_{j^2;m_{j^2}})_{ss}$.

2.2.5 Proposition

Let $\mathrm{GL}_n(\mathbb{Z}/N\,\mathbb{Z})^2$ *(resp.* $\mathrm{GL}_n(\mathbb{Z})^2)$ *) be the "pseudo-ramified" (resp. "pseudo-unramified") general bilinear semigroup of matrices of order n with entries in* $\mathbb{Z}/N\,\mathbb{Z}$ *(resp.* $\mathbb{Z}$ *). Then,* $\mathcal{H}_{R\times L}(n)$ *(resp.* $\mathcal{H}^{nr}_{R\times L}(n)$ *) will denote the pseudo-ramified (resp. pseudo-unramified) Hecke bialgebra generated by all the pseudo-ramified (resp. pseudo-unramified) Hecke bioperators* $(T_R(n;r)\otimes T_L(n;r))$ *(resp.* $(T^{nr}_R(n;r)\otimes T^{nr}_L(n;r))$ *) having a representation in the subgroup of matrices of* $\mathrm{GL}_n(\mathbb{Z}/N\,\mathbb{Z})^2)$ *(resp.* $\mathrm{GL}_n(\mathbb{Z})^2)$ *),* $1\le j\le r\le\infty$ *, the index j varying.*

Proof:

1. A left (resp. right) Hecke operator $T_L(n;r)$ (resp. $T_R(n;r)$) is a left (resp. right) correspondence which associates to a left (resp. right) lattice Λ_ω (resp. $\Lambda_{\overline{\omega}}$) the sum of its left (resp. right) sublattices Λ_{ω_j} (resp. $\Lambda_{\overline{\omega}_j}$) of index j and multiplicities $m^{(j)}=\sup(m_j)$:

    $$T_L(n;r)\ \Lambda_\omega = \bigoplus_{j=1}^{r} \bigoplus_{m_j} \Lambda_{\omega_{j,m_j}}$$

    $$\text{(resp.}\quad T_R(n;r)\ \Lambda_{\overline{\omega}} = \bigoplus_{j=1}^{r} \bigoplus_{m_j} \Lambda_{\overline{\omega}_{j,m_j}}\ \text{)}.$$

    A Hecke bioperator $T_R(n;r)\otimes T_L(n;r)$ is then defined by its action on the pseudo-ramified bilattice $\Lambda_\omega\otimes\Lambda_{\overline{\omega}}$

    $$(T_R(n;r)\otimes T_L(n;r))\ (\Lambda_\omega\otimes\Lambda_{\overline{\omega}}) = \bigoplus_{j=1}^{r} \bigoplus_{m_j} (\Lambda_{\overline{\omega}_{j,m_j}} \otimes \Lambda_{\omega_{j,m_j}})\ .$$

According to proposition 2.2.2, we have

$$\Lambda_{\overline{\omega}} \otimes \Lambda_\omega \simeq \mathrm{GL}_n(\mathcal{O}_{F,\overline{\omega}} \times \mathcal{O}_{F,\omega})$$

such that

$$\Lambda_{\overline{\omega}_{j,m_j}} \otimes \Lambda_{\omega_{j,m_j}} \simeq g_n(\mathcal{O}_{F_{\overline{\omega}_{j,m_j}}} \times \mathcal{O}_{F_{\omega_{j,m_j}}}) \in \mathrm{GL}_n(\mathcal{O}_{F,\overline{\omega}} \times \mathcal{O}_{F,\omega})$$

where $\mathcal{O}_{F_{\omega_{j,m_j}}}$ (resp. $\mathcal{O}_{F_{\overline{\omega}_{j,m_j}}}$) is the maximal order $\mathcal{O}_{F,\omega}$ (resp. $\mathcal{O}_{F,\overline{\omega}}$) restricted to the extension $\widetilde{F}_{\omega_{j,m_j}}$ (resp. $\widetilde{F}_{\overline{\omega}_{j,m_j}}$).

Remark that the entries of $g_n(\mathcal{O}_{F_{\overline{\omega}_{j,m_j}}} \times \mathcal{O}_{F_{\omega_{j,m_j}}})$ are integers of $(\widetilde{F}_{\overline{\omega}_{j,m_j}} \times \widetilde{F}_{\omega_{j,m_j}})$ and the eigenvalues of $g_n(\mathcal{O}_{F_{\overline{\omega}_{j,m_j}}} \times \mathcal{O}_{F_{\omega_{j,m_j}}})$ are algebraic integers.

The bisublattice $\Lambda_{\overline{\omega}_{j,m_j}} \otimes \Lambda_{\omega_{j,m_j}}$ corresponds to the coset representative $g_n(\mathcal{O}_{F_{\overline{\omega}_{j,m_j}}} \times \mathcal{O}_{F_{\omega_{j,m_j}}})$ of $\mathrm{GL}_n(\mathcal{O}_{F,\overline{\omega}} \times \mathcal{O}_{F,\omega_j})$, which involves that $g_n(\mathcal{O}_{F_{\overline{\omega}_{j,m_j}}} \times \mathcal{O}_{F_{\omega_{j,m_j}}})$ can be chosen as a coset representative of the tensor product of Hecke operators [Lang].

2. Indeed, the ring of endomorphisms of the $(\mathbb{Z}/N\,\mathbb{Z})^2$-bilattice $\Lambda_{\overline{\omega}} \times \Lambda_\omega$ is generated over $(\mathbb{Z}/N\,\mathbb{Z})^2$ by the products $(T_{j_R} \otimes T_{j_L})$ of Hecke operators T_{j_R} and T_{j_L} for the primes $j \nmid N$ and by the products $(U_{j_R} \otimes U_{j_L})$ of Hecke operators U_{j_R} and U_{j_L} for $j \mid N$ [M-W].

   The coset representative of U_{j_L} , referring to the upper half space, can be chosen to be upper-triangular and given by the integral matrix

$$t_{n;j,m_j} = \begin{pmatrix} j_{1_N} b^1_{2_N} & \cdots & \cdots & b^1_{n_N} \\ & j_{2_N} & \cdots & b^2_{n_N} \\ & & \ddots & \vdots \\ & 0 & & j_{n_N} \end{pmatrix} \qquad \text{of} \quad T_n(\mathcal{O}_{F,\omega}) \subset T_n(\mathbb{Z}/N\,\mathbb{Z})$$

   such that:

   - $j_{n_N} = 0 \bmod N$,
   - $b^1_{2_N} = 0 \bmod N$,
   - $j_{1_N} \times \cdots j_{n_N} \simeq j^n \cdot N^n \cdot (m^{(j)})^n$ such that $(f_{\overline{\omega}_j} \cdot f_{\omega_j})^n \cdot N^{2n} \simeq j^{2n} \cdot N^{2n} \cdot (m^{(j)})^{2n}$ (see definition 2.2.4),

- $t_{n;j,m_j} \equiv t_n(\mathcal{O}_{F_{\omega_{j,m_j}}}) \subset g_n(\mathcal{O}_{F_{\overline{\omega}_{j,m_j}}} \times \mathcal{O}_{F_{\omega_{j,m_j}}})$.

Similarly, the coset representative of U_{j_R} , referring to the lower half space can be chosen to be lower triangular and given by the integral matrix $t^t_{n;j,m_j} \equiv t^t_n(\mathcal{O}_{F_{\overline{\omega}_{j;m_j}}})$ of $T^t_n(\mathcal{O}_{F,\overline{\omega}})$, i.e. by the transposed matrix of $t_{n;j,m_j}$.

So, the coset representative of $(U_{j_R} \times U_{j_L})$ will be

$$g_n(\mathcal{O}_{F_{\overline{\omega}_{j,m_j}}} \times \mathcal{O}_{F_{\omega_{j,m_j}}}) = t^t_{n;j,m_j} \times t_{n;j,m_j} \in \mathrm{GL}_n(\mathcal{O}_{F,\overline{\omega}} \times \mathcal{O}_{F,\omega}) \subset \mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2) \ .$$

Note that $g_n(\mathcal{O}_{F_{\overline{\omega}_{j,m_j}}} \times \mathcal{O}_{F_{\omega_{j,m_j}}})$ has the Gauss decomposition in diagonal and nilpotent part so that the nilpotent part is given by $u_n(\mathcal{O}_{F_{\omega_{j,m_j}}}) \times u^t_n(\mathcal{O}_{F_{\overline{\omega}_{j,m_j}}})$ and constitutes an element of the decomposition group $D_{j^2;m_j^2}$ of the corresponding bisublattice $(\Lambda_{\overline{\omega}_{j,m_j}} \otimes \Lambda_{\omega_{j,m_j}})$ as introduced in definition 2.2.4.

3. The pseudo-unramified Hecke bioperator $T^{nr}_R(n;r) \otimes T^{nr}_L(n;r)$ can also be envisaged throughout its action on the pseudo-unramified bilattice $\Lambda^{nr}_{\overline{\omega}} \otimes \Lambda^{nr}_{\omega}$:

$$T^{nr}_R(n;r) \otimes T^{nr}_L(n;r))\ (\Lambda^{nr}_{\overline{\omega}} \otimes \Lambda^{nr}_{\omega}) = \bigoplus_{j=1}^{r} \bigoplus_{m_j} (\Lambda^{nr}_{\overline{\omega}_{j,m_j}} \otimes \Lambda^{nr}_{\omega_{j,m_j}})$$

such that:

$$(\Lambda^{nr}_{\overline{\omega}_{j,m_j}} \otimes \Lambda^{nr}_{\omega_{j,m_j}}) \simeq g_n(\mathcal{O}_{F^{nr}_{\overline{\omega}_{j,m_j}}} \times \mathcal{O}_{F^{nr}_{\omega_{j,m_j}}}) \in \mathrm{GL}_n(\mathcal{O}_{F^{nr},\overline{\omega}} \times \mathcal{O}_{F^{nr},\omega}) \subset \mathrm{GL}_n(\mathbb{Z}^2) \ .$$

The pseudo-unramified correspondent $U^{nr}_{j_L}$ (resp. $U^{nr}_{j_R}$) of the coset representative of U_{j_L} (resp. U_{j_R}) is an upper (resp. lower) triangular integral matrix $t_n(\mathcal{O}_{F^{nr}_{\omega_{j,m_j}}})$ (resp. $t^t_n(\mathcal{O}_{F^{nr}_{\overline{\omega}_{j,m_j}}})$) of $T_n(\mathcal{O}_{F^{nr},\omega})$ (resp. $T_n(\mathcal{O}_{F^{nr},\overline{\omega}})$) $\subset T_n(\mathbb{Z})$ (resp. $T^t_n(\mathbb{Z})$). ■

2.2.6 Proposition

The Hecke bialgebra $\mathcal{H}_{R\times L}(n)$ *(resp.* $\mathcal{H}^{nr}_{R\times L}(n)$ *) of the endomorphisms of the pseudo-ramified (resp. pseudo-unramified)* $B_{\overline{\omega}} \otimes B_{\omega}$*-bisemimodule* $M_R \otimes M_L$ *(resp.* $B^{nr}_{\overline{\omega}} \otimes B^{nr}_{\omega}$*-bisemimodule* $M^{nr}_R \otimes M^{nr}_L$ *) is generated by the pseudo-ramified (resp. pseudo-unramified) Hecke bioperators* $(T_R(n;r) \otimes T_L(n;r))$ *(resp.* $(T^{nr}_R(n;r) \otimes T^{nr}_L(n;r))$ *) having as representation* $\mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)$ *(resp.* $\mathrm{GL}_n((\mathbb{Z})^2)$ *).*

Proof: as $\Lambda_{\overline{\omega}} \otimes \Lambda_{\omega}$ is a pseudo-ramified bilattice in the $B_{\overline{\omega}} \otimes B_{\omega}$-bisemimodule $M_R \otimes M_L$ and as the decomposition of the $B_{\overline{\omega}} \otimes B_{\omega}$-bisemimodule $M_R \otimes M_L$ into the set of $B_{\overline{\omega}_j} \otimes B_{\omega_j}$-subbisemimodules $M_{\overline{\omega}_j} \otimes M_{\omega_j}$ having multiplicities m_j is obtained through the action of the Hecke bioperator $(T_R(n;r) \otimes T_L(n;r))$, it is immediate to check that $(T_R(n;r) \otimes T_L(n;r))$

has a representation in the bilinear congruence subgroup $\mathrm{GL}_n(\mathbb{Z}/N\,\mathbb{Z})^2)$.
The pseudo-unramified case can be handled similarly. ■

2.3 Lattice semispaces

2.3.1 Definition: pseudo-ramified lattice semispaces

The space $X = \mathrm{GL}_n(\mathbb{R})/\,\mathrm{GL}_n(\mathbb{Z})$ corresponds to the set of lattices of $\mathbb{R}^n$. In this perspective, if $\widetilde{F}_L$ (resp. $\widetilde{F}_R$) denotes a complex symmetric splitting semifield (see section 1.1.4), we shall introduce a left (resp. right) (pseudo-)ramified lattice semispace:

$$X_{S_L} = T_n(\widetilde{F}_L)\Big/T_n(\mathbb{Z}/N\,\mathbb{Z}) \qquad (\text{resp.}\quad X_{S_R} = T^t_n(\widetilde{F}_L)\Big/T^t_n(\mathbb{Z}/N\,\mathbb{Z})\ )$$

isomorphic to the left (resp. right) (pseudo-)ramified B_ω-semimodule M_L (resp. $B_{\overline{\omega}}$-semimodule M_R) such that the left (resp. right) cosets of X_{S_L} (resp. X_{S_R}) are isomorphic to the left (resp. right) pseudo-ramified B_{ω_j}-subsemimodules M_{ω_j} (resp. $B_{\overline{\omega}_j}$-subsemimodules $M_{\overline{\omega}_j}$).
Consequently, we can define a pseudo-ramified lattice bisemispace

$$X_{S_{R\times L}} = \mathrm{GL}_n(\widetilde{F}_R\times\widetilde{F}_L)\Big/\,\mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)$$

isomorphic to the pseudo-ramified $B_{\overline{\omega}}\otimes B_\omega$-bisemimodule $M_R\otimes M_L$.
As the left (resp. right) pseudo-ramified lattice semispace X_{S_L} (resp. X_{S_R}) is defined with respect to the matrix algebra $T_n(\mathbb{Z}/N\,\mathbb{Z})$ (resp. $T^t_n(\mathbb{Z}/N\,\mathbb{Z})$) in one-to-one correspondence with the pseudo-ramified left (resp. right) lattice Λ_ω (resp. $\Lambda_{\overline{\omega}}$), X_{S_L} (resp. X_{S_R}) is a left (resp. right) Hecke lattice semispace.
Similarly, the lattice bisemispace $X_{S_{R\times L}}$ is a pseudo-ramified Hecke lattice bisemispace since $\mathrm{GL}_n(\mathbb{Z}/N\,\mathbb{Z})$ is in one-to-one correspondence with the bilattice $\Lambda_{\overline{\omega}\times\omega} = \Lambda_{\overline{\omega}}\times\Lambda_\omega$.

2.3.2 Definition: pseudo-unramified Hecke lattice semispaces

A left (resp. right) pseudo-unramified lattice semispace:

$$X^{nr}_{S_L} = T_n(\widetilde{F}^{nr}_L)\Big/T_n(\mathbb{Z}) \qquad (\text{resp.}\quad X^{nr}_{S_R} = T^t_n(\widetilde{F}^{nr}_R)\Big/T^t_n(\mathbb{Z})$$

can similarly be introduced; it is a left (resp. right) pseudo-unramified Hecke lattice semispace isomorphic to the left (resp. right) pseudo-unramified B^{nr}_ω-semimodule M^{nr}_L (resp. $B^{nr}_{\overline{\omega}}$-semimodule M^{nr}_R) verifying $M^{nr}_L\subset M_L$ (resp. $M^{nr}_R\subset M_R$).
A pseudo-unramified lattice bisemispace

$$X^{nr}_{S_{R\times L}} = \mathrm{GL}_n(\widetilde{F}^{nr}_R\times\widetilde{F}^{nr}_L)\Big/\,\mathrm{GL}_n((\mathbb{Z}^2))$$

can also be defined: it is a pseudo-unramified Hecke lattice bisemispace isomorphic to the pseudo-unramified $B^{nr}_{\overline{\omega}} \otimes B^{nr}_{\omega}$-bisemimodule $M^{nr}_R \otimes M^{nr}_L$.

2.3.3 Proposition

a) *Any left (resp. right) coset*

$$X^{nr}_{S_L}(\widetilde F^{nr}_{\omega_j}) = t_n(\widetilde F^{nr}_{\omega_j}) \Big/ t_n(\mathcal{O}_{F^{nr}_{\omega_j}}) \qquad \textit{(resp.} \quad X^{nr}_{S_R}(\widetilde F^{nr}_{\overline{\omega}_j}) = t_n(\widetilde F^{nr}_{\overline{\omega}_j})^t \Big/ t_n(\mathcal{O}_{F^{nr}_{\overline{\omega}_j}})^t \ )$$

of $X^{nr}_{S_L}$ *(resp.* $X^{nr}_{S_R}$ *) having a multiplicity one is a left (resp. right) pseudo-unramified Hecke lattice semisubspace having a rank equal to* j^{2n} *if* $f_{\omega_j} = j$.

b) *the product* $X^{nr}_{S_{R\times L}}(\widetilde F^{nr}_{\overline{\omega}_j} \times \widetilde F^{nr}_{\omega_j}) = g_n(\widetilde F^{nr}_{\overline{\omega}_j} \times \widetilde F^{nr}_{\omega_j}) \Big/ g_n(\mathcal{O}_{F^{nr}_{\overline{\omega}_j}} \times \mathcal{O}_{F^{nr}_{\omega_j}})$ *of a right coset* $X^{nr}_{S_R}(\widetilde F^{nr}_{\overline{\omega}_j})$ *by a left coset* $X^{nr}_{S_L}(\widetilde F^{nr}_{\omega_j})$ *having multiplicities one is a pseudo-unramified Hecke lattice bisemisubspace having a rank equal to* j^{2n} .

Proof:

a) The left (resp. right) pseudo-unramified Hecke lattice semisubspace $X^{nr}_{S_L}(\widetilde F^{nr}_{\omega_j})$ (resp. $X^{nr}_{S_R}(\widetilde F^{nr}_{\overline{\omega}_j})$) is isomorphic to the pseudo-unramified left (resp. right) $B^{nr}_{\omega_j}$-subsemimodule $M^{nr}_{\omega_j}$ (resp. $B^{nr}_{\overline{\omega}_j}$-subsemimodule $M^{nr}_{\overline{\omega}_j}$) which has a rank equal to the n-fold product of its global class residue degree $f_{\omega_j} = [F^{nr}_{\omega_j} : k_L] = j$ (resp. $f_{\overline{\omega}_j} = [F^{nr}_{\overline{\omega}_j} : k_R] = j$); so, we have that the rank of $X^{nr}_{S_L}(F^{nr}_{\omega_j})$ (resp. $X^{nr}_{S_R}(F^{nr}_{\overline{\omega}_j})$) is equal to j^{2n} .

b) the pseudo-unramified Hecke lattice bisemisubspace $X^{nr}_{S_{R\times L}}(\widetilde F^{nr}_{\overline{\omega}_j} \times \widetilde F^{nr}_{\omega_j})$ is isomorphic to the pseudo-unramified $B^{nr}_{\overline{\omega}_j} \otimes B^{nr}_{\omega_j}$-subbisemimodule $M^{nr}_{\overline{\omega}_j} \otimes M^{nr}_{\omega_j}$ having a rank equal to j^{2n} . ■

2.3.4 Proposition

Let $f_{\omega_j} = j$ denote the global class residue degree of the j-th place having multiplicity $m^{(j)}$.

a) *The pseudo-unramified* $B^{nr}_{\overline{\omega}} \otimes B^{nr}_{\omega}$*-bisemimodule* $M^{nr}_R \otimes M^{nr}_L$ *has a rank given by:*

$$r^{nr}_{\overline{\omega}\times\omega} = \bigoplus_{j=1}^{r} (f_{\overline{\omega}_j} \cdot m^{(j)})^{2n} (f_{\omega_j} \cdot m^{(j)})^{2n} = \bigoplus_j (j \cdot m^{(j)})^{2n} \ .$$

b) *the pseudo-ramified* $B_{\overline{\omega}} \otimes B_{\omega}$*-bisemimodule* $M_R \otimes M_L$ *has a rank given by:*

$$r_{\overline{\omega}\times\omega} = \bigoplus_{j=1}^{r} N^{2n} (f_{\overline{\omega}_j} \cdot m^{(j)})^{2n} (f_{\omega_j} \cdot m^{(j)})^{2n} = \bigoplus_j N^{2n} (j \cdot m^{(j)})^{2n} \ .$$

2.3.5 Summary

The preceding developments can be summarized as follows:
Let

- $\Lambda_{\overline{\omega}}\otimes\Lambda_\omega$ be a $(\mathbb{Z}/N\,\mathbb{Z})^2$-bilattice (pseudo-ramified);
- $\Lambda^{nr}_{\overline{\omega}}\otimes\Lambda^{nr}_\omega$ be a $\mathbb{Z}^2$-bilattice (pseudo-unramified);
- $M^{nr}_R\otimes M^{nr}_L$ be a $\mathrm{GL}_n(F^{nr}_{\overline{\omega}}\times F^{nr}_\omega)$-bisemimodule (pseudo-unramified) isomorphic to the pseudo-unramified lattice bisemispace $X^{nr}_{S_{R\times L}}$;
- $M_R\otimes M_L$ be a $\mathrm{GL}_n(F_{\overline{\omega}}\times F_\omega)$-bisemimodule (pseudo-ramified) isomorphic to the pseudo-ramified lattice bisemispace $X_{S_{R\times L}}$.

Then, the following commutative diagram is evident:

$$\begin{tikzcd}
\Lambda^{nr}_{\overline{\omega}}\otimes\Lambda^{nr}_\omega \arrow[rr, hook] \arrow[dd] & & \Lambda_{\overline{\omega}}\otimes\Lambda_\omega \arrow[dd] & : & \text{bilattices} \arrow[dd] \\
{} \arrow[r, dashed, "\mathit{pseudo-ramification}"'] & {} & & & \\
M^{nr}_R\otimes M^{nr}_L \arrow[rr, hook] & & M_R\otimes M_L & : & \mathrm{GL}_n(F^{(nr)}_{\overline{\omega}}\times F^{(nr)}_\omega)\text{-bisemimodules}
\end{tikzcd}$$

where

- the vertical arrows stand for injective morphisms from bilattices to $\mathrm{GL}_n(F^{(nr)}_{\overline{\omega}}\times F^{(nr)}_\omega)$-bisemimodules;
- the horizontal arrows stand for pseudo-ramification injective morphisms.

2.4 Representations of global Weil groups

2.4.1 Nonabelian global class field concepts: a summary

1. Let $G^{(2n)}(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_\omega)$, $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_\omega)$ and $\mathrm{GL}(\widetilde{M}_R\otimes\widetilde{M}_L)$ denote respectively the bilinear algebraic semigroup over $(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_\omega)$ which is a bilinear affine semigroup, its bilinear algebraic semigroup of matrices and the group of automorphisms of $(\widetilde{M}_R\otimes\widetilde{M}_L)$ which are isomorphic:

$$G^{(2n)}(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_\omega)\simeq\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_\omega)\simeq\mathrm{GL}(\widetilde{M}_R\otimes\widetilde{M}_L)$$

according to section 2.1.1: this implies the choice of a basis in $\widetilde{M}_R \otimes \widetilde{M}_L$, which is evident from its construction from $(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$.

The diagonal conjugacy classes of $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$, noted $g^{(2n)}_{R\times L}[j,m_j=0]$, then correspond to the $\widetilde{B}_{\overline{\omega}_j} \otimes \widetilde{B}_{\omega_j}$-subbisemimodules $\widetilde{M}_{\overline{\omega}_j} \otimes \widetilde{M}_{\omega_j}$ and their off-diagonal equivalents $g^{(2n)}_{R\times L}[j,m_j]$, $m_j > 0$, related to the multiplicity $m^{(j)} = \sup(m_j)$ of $g^{(2n)}_{R\times L}[j,m_j=0]$, correspond to the $\widetilde{B}_{\overline{\omega}_{j,m_j}} \otimes \widetilde{B}_{\omega_{j,m_j}}$-subbisemimodules $\widetilde{M}_{\overline{\omega}_{j,m_j}} \otimes \widetilde{M}_{\omega_{j,m_j}}$: they are obtained from $(\widetilde{M}_R \otimes \widetilde{M}_L)$ by the action of the Hecke bioperator $T_R(n;r) \otimes T_L(n;r)$.

As $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$ is a smooth reductive bilinear affine semigroup, it can be considered as the n-dimensional analog of the product $(\mathbb{S}^{\,1}_R \times \mathbb{S}^{\,1}_L)$ of the one-dimensional affine complex semigroups $\mathbb{S}^{\,1}_R$ and $\mathbb{S}^{\,1}_L$, introduced in section 2.1.1, we have that:

$$G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega}) \simeq \prod_n (\mathbb{S}^{\,1}_R \times \mathbb{S}^{\,1}_L) \ .$$

Similarly, $\mathrm{GL}(\widetilde{M}_{R_\oplus} \otimes \widetilde{M}_{L_\oplus})$ can be viewed as the n-dimensional correspondent of the product $W^{ab}_{F_R} \times W^{ab}_{F_L}$ of the Weil global groups included in the product $\mathrm{Gal}(F^{ac}_R/k) \times \mathrm{Gal}(F^{ac}_L/k)$ of the Galois groups as introduced in sections 1.1.7 and 1.1.9 which implies that:

$$\mathrm{GL}(\widetilde{M}_{R_\oplus} \otimes \widetilde{M}_{L_\oplus}) \simeq \prod_n (W^{ab}_{F_R} \times W^{ab}_{F_L}) \subset \prod_n (\mathrm{Gal}(F^{ac}_R/k_R) \times \mathrm{Gal}(F^{ac}_L/k_L)) \ .$$

2. Let us now introduce the normal bilinear algebraic semisubgroup $P^{(2n)}(\widetilde{F}_{\overline{\omega}^1} \times \widetilde{F}_{\omega^1})$ of $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$: its j-th class noted $P^{(2n)}(\widetilde{F}_{\overline{\omega}^1_j} \times \widetilde{F}_{\omega^1_j})$ is defined over the product $(\widetilde{F}_{\overline{\omega}^1_j} \times \widetilde{F}_{\omega^1_j})$ of irreducible central extensions $\widetilde{F}_{\overline{\omega}^1_j}$ and $\widetilde{F}_{\omega^1_j}$ having a rank $N \cdot m^{(j)}$, as given in section 1.1.5. So, $P^{(2n)}(\widetilde{F}_{\overline{\omega}^1_j} \times \widetilde{F}_{\omega^1_j})$ has a rank equal to $(N \cdot m^{(j)})^{2n}$. $P^{(2n)}(\widetilde{F}_{\overline{\omega}^1} \times \widetilde{F}_{\omega^1})$ also acts on $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$ by conjugation.

   Recall that the Galois subgroup of the irreducible extension $\widetilde{F}_{\omega^1_j}$ associated to the completion $F_{\omega^1_j}$ is the global inertia subgroup $I_{F_{\omega_j}}$.

   More generally, let

$$\begin{aligned} \widetilde{F}_{\omega^1} &= \{\widetilde{F}_{\omega^1_1}, \cdots, \widetilde{F}_{\omega^1_{j,m_j}}, \cdots, \widetilde{F}_{\omega^1_r}\} \\ (\text{resp.}\quad \widetilde{F}_{\overline{\omega}^1} &= \{\widetilde{F}_{\overline{\omega}^1_1}, \cdots, \widetilde{F}_{\overline{\omega}^1_{j,m_j}}, \cdots, \widetilde{F}_{\overline{\omega}^1_r}\}\ ) \end{aligned}$$

   denote the set of pseudo-ramified irreducible extensions $\widetilde{F}_{\omega^1_j}$ and $\widetilde{F}_{\omega^1_{j,m_j}}$ of $\widetilde{F}_{\omega}$ and $\widetilde{F}_{\overline{\omega}}$ respectively.

Let $G^{(2n)}(\widetilde{F}^{nr}_{\overline{\omega}} \times \widetilde{F}^{nr}_{\omega})$ denote the pseudo-unramified bilinear algebraic semigroup over the product $\widetilde{F}^{nr}_{\overline{\omega}} \times \widetilde{F}^{nr}_{\omega}$ of pseudo-unramified extensions.

Then, $P^{(2n)}(\widetilde{F}_{\overline{\omega}^1} \times \widetilde{F}_{\omega^1})$, which is in fact a bilinear (complex) minimal (not standard) parabolic subsemigroup, can be defined as being the kernel of the map:

$$G^{(2n)}_{F\to F^{nr}} : \quad G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega}) \longrightarrow G^{(2n)}(\widetilde{F}^{nr}_{\overline{\omega}} \times \widetilde{F}^{nr}_{\omega}) \ ;$$

so:

$$P^{(2n)}(\widetilde{F}_{\overline{\omega}^1} \times \widetilde{F}_{\omega^1}) = \operatorname{Ker}(G^{(n)}_{F\to F^{nr}})$$

is the smallest bilinear normal pseudo-ramified subgroup of $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$: it is the minimal parabolic subgroup viewed as the connected component of the identity in $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$ on the Zarisky topology.

3. Referring to the decomposition of the product, right by left, $\widetilde{F}_{\overline{\omega}_j} \times \widetilde{F}_{\omega_j}$ of the basic extensions $\widetilde{F}_{\overline{\omega}_j}$ and $\widetilde{F}_{\omega_j}$ into j' equivalent bisubextensions $\widetilde{F}_{\overline{\omega}^{j'}_j} \times \widetilde{F}_{\omega^{j'}_j}$, $1 \le j' \le j$, as well as the decomposition of the equivalent biextensions $\widetilde{F}_{\overline{\omega}_{j,m_j}} \times \widetilde{F}_{\omega_{j,m_j}}$ into j' equivalent bisubextensions $\widetilde{F}_{\overline{\omega}^{j'}_{j,m_j}} \times \widetilde{F}_{\omega^{j'}_{j,m_j}}$, we have that the j-th diagonal conjugacy class $g^{(2n)}_{R\times L}[j,m_j=0]$ of $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$ will be cut into j' equivalent diagonal conjugacy subclasses $g^{(2n)}_{R\times L}[j,m_j=0,j']$ and that the j-th off-diagonal equivalent conjugacy class $g^{(2n)}_{R\times L}[j,m_j]$, $m_j>0$, will also be cut into j' equivalent off-diagonal conjugacy subclasses $g^{(2n)}_{R\times L}[j,m_j,j']$, $1 \le j' \le j$.

   So, the j-th diagonal conjugacy class $g^{(2n)}_{R\times L}[j,m_j=0]$ and its off-diagonal equivalent $g^{(2n)}_{R\times L}[j,m_j]$, $m_j>0$, is decomposed into a set $\{g^{(2n)}_{R\times L}[j,m_j,j']\}^{j}_{j'=1}$ of j' conjugacy subclasses $g^{(2n)}_{R\times L}[j,m_j,j']$.

Then, we can state the following propositions:

2.4.2 Lemma

Let $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$ be a smooth reductive bilinear affine semigroup and let $P^{(2n)}(\widetilde{F}_{\overline{\omega}^1} \times \widetilde{F}_{\omega^1})$ be its parabolic subgroup viewed as a locally subgroup of $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$.
Then, it is clear that:

1. *at every (bi)extension corresponding to the (bi)place $\overline{\omega}_j \times \omega_j$ of $(F_{\overline{\omega}} \times F_{\omega})$ corresponds a conjugacy class of $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$ whose number of representatives corresponds to the number of equivalent completions of $\overline{\omega}_j \times \omega_j$.*

2. *$G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_\omega)$ acts on $P^{(2n)}(\widetilde F^1_{\overline\omega}\times\widetilde F^1_\omega)$ by conjugation in such a way that the number of conjugates of $P^{(2n)}(\widetilde F_{\overline\omega^1_j}\times\widetilde F_{\omega^1_j})$ in $G^{(2n)}(\widetilde F_{\overline\omega_{j,m_j}}\times\widetilde F_{\omega_{j,m_j}})$ is the index of the normalizer of $P^{(2n)}(\widetilde F_{\overline\omega^1_j}\times\widetilde F_{\omega^1_j})$,*

$$\left|G^{(2n)}(\widetilde F_{\overline\omega_{j,m_j}}\times\widetilde F_{\omega_{j,m_j}}):N_{G^{(2n)}}(P^{(2n)}(\widetilde F_{\overline\omega^1_j}\times\widetilde F_{\omega^1_j}))\right|=j$$

where

- *$G^{(2n)}(\widetilde F_{\overline\omega_{j,m_j}}\times\widetilde F_{\omega_{j,m_j}})$ is $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_\omega)$ restricted to the conjugacy class representative $g^{(2n)}_{R\times L}[j,m_j]$;*
- *$N_{G^{(n)}}(P^{(2n)}(\widetilde F_{\overline\omega^1_j}\times\widetilde F_{\omega^1_j}))$ is the normalizer of $P^{(2n)}(\widetilde F_{\overline\omega^1_j}\times\widetilde F_{\omega^1_j})$ restricted to the j-th irreducible subextension $\widetilde F_{\overline\omega^1_j}\times\widetilde F_{\omega^1_j}$;*
- *j is the global residue degree of the j-th extension class $\widetilde\omega_j$ (or $\widetilde{\overline\omega}_j$).*

Proof:

1. As the bilinear affine semigroup $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_\omega)$ is assumed to be reductive, each conjugacy class representative of $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_\omega)$ is isogeneous to a direct product of tori (or of simple groups). But, taking into account the projective toroidal compactification of the conjugacy classes $g^{(2n)}_{R\times L}[j,m_j]\in G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_\omega)$, introduced in section 3.3, we shall assume that $g^{(2n)}n_{R\times L}[j,m_j]$ is isomorphic to a direct product of tori, which is the product, right by left, of algebraic (semi)tori.

   So, $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_\omega)\simeq\prod\limits_n(\mathbb S^1_R\times\mathbb S^1_L)$, which is the product, right by left, of algebraic (semi)tori,is considered as the n-fold product of a right affine semigroup $\mathbb S^1_R$ localized in the lower half space by its left equivalent $\mathbb S^1_L$ localized in the upper half space.

   But, as $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_\omega)$ is defined over the product $\widetilde F_{\overline\omega}\times\widetilde F_\omega$, and as $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_\omega)$ is assumed to act on $P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})$ by conjugation, a conjugacy class representative $g^{2n}_{R\times L}[j,m_j]\in G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_\omega)$ corresponds to the biextension $\widetilde F_{\overline\omega_{j,m_j}}\times\widetilde F_{\omega_{j,m_j}}$. Then, we have the injective morphism:

   $$Fg:\quad\{\widetilde F_{\overline\omega_{j,m_j}}\times\widetilde F_{\omega_{j,m_j}}\}\longrightarrow\{g^{(2n)}_{R\times L}[j,m_j]$$

   from the set of equivalent representatives $\widetilde F_{\overline\omega_{j,m_j}}\times\widetilde F_{\omega_{j,m_j}}$, associated with all places $\overline\omega_j\times\omega_j$ of $F_R\times F_L$, to the set of corresponding conjugacy class representatives $g^{(2n)}_{R\times L}[j,m_j]$ of $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_\omega)$, such that $\prod\limits_n(\widetilde F_{\overline\omega_{j,m_j}}\times\widetilde F_{\omega_{j,m_j}})\simeq g^{(2n)}_{R\times L}[j,m_j]$.

2. $G^{(2n)}(\widetilde F_{\overline\omega_{j,m_j}}\times \widetilde F_{\omega_{j,m_j}})$ acts on $P^{(2n)}(\widetilde F_{\overline\omega^1_j}\times \widetilde F_{\omega^1_j})$ by conjugation, in the sense that, if $p_{j_R}\in P^{(2n)}(\widetilde F_{\overline\omega^1_j})$, $p_{j_L}\in P^{(2n)}(\widetilde F_{\omega^1_j})$ and $x_{g_{j_R}}\in G^{(2n)}(\widetilde F_{\overline\omega_{j,m_j}})$, $x_{g_{j_L}}\in G^{(2n)}(\widetilde F_{\omega_{j,m_j}})$, then the bielement $x_{g_{j_R}}\cdot x_{g_{j_L}}\in g^{(2n)}_{R\times L}[j,m_j]$ is said to be conjugate of the bielement $x'_{g_{j_R}}\cdot x'_{g_{j_L}}\in g^{(2n)}_{R\times L}[j,m_j]$ if:

$$x'_{g_{j_R}}\cdot x'_{g_{j_L}}=p_{j_R}\cdot x_{g_{j_R}}\cdot x_{g_{j_L}}\cdot p_{j_L}\ .$$

On the other hand, as a extension $\widetilde F_{\omega_{j,m_j}}$ (resp. $\widetilde F_{\overline\omega_{j,m_j}}$) , is cut into j' irreducible equivalent subextensions $\widetilde F_{\omega^{j'}_{j,m_j}}$ (resp. $\widetilde F_{\overline\omega^{j'}_{j,m_j}}$) according to section 1.1.5, the conjugacy class representative $g^{(2n)}_{R\times L}[j,m_j]$ will also be cut into " j' " equivalent conjugacy class subrepresentatives $g^{(2n)}_{R\times L}[j,m_j,j']$, $1\le j'\le j$, with respect to 1.

So, we have the corresponding injective morphism:

$$\begin{aligned}F_{g_{j'}}:\quad \widetilde F_{\overline\omega_{j,m_j}}\times \widetilde F_{\omega_{j,m_j}} &= \{\widetilde F_{\overline\omega^{j'}_{j,m_j}}\times \widetilde F_{\omega^{j'}_{j,m_j}}\}^j_{j'=1}\\ \longrightarrow\quad & g^{(2n)}_{R\times L}[j,m_j]=\{g^{(2n)}_{R\times L}[j,m_j,j']^j_{j'=1}\}\ .\end{aligned}$$

This implies that the conjugation of $G^{(2n)}(\widetilde F_{\overline\omega_{j,m_j}}\times \widetilde F_{\omega_{j,m_j}})$ by $P^{(2n)}(\widetilde F_{\overline\omega^1_j}\times \widetilde F_{\omega^1_j})$ sends the bielement $x_{g_{j_R}}\cdot x_{g_{j_L}}\in g^{(2n)}_{R\times L}[j,m_j,j'']$, $1\le j''\le j$, of the j''-th conjugacy class subrepresentative of $g^{(2n)}_{R\times L}[j,m_j]$ into the bielement $x'_{g_{j_R}}\cdot x'_{g_{j_L}}$ of the j'-th conjugacy class subrepresentative of $g^{(2n)}_{R\times L}[j,m_j]$.

Finally, the number of conjugacy class subrepresentatives of $g^{(2n)}_{R\times L}[j,m_j]$ is exactly the number of conjugates of $P^{(2n)}(\widetilde F_{\overline\omega^1_j}\times \widetilde F_{\omega^1_j})$ in $G^{(2n)}(\widetilde F_{\overline\omega_{j,m_j}}\times \widetilde F_{\omega_{j,m_j}})\equiv g^{(2n)}_{R\times L}[j,m_j]$.

So the index of the normalizer $N_{G^{(2n)}}(P^{(2n)}(\widetilde F_{\overline\omega^1_j}\times \widetilde F_{\omega^1_j}))$ of $P^{(2n)}(\widetilde F_{\overline\omega^1_j}\times \widetilde F_{\omega^1_j})$ in $G^{(2n)}(\widetilde F_{\overline\omega_{j,m_j}}\times \widetilde F_{\omega_{j,m_j}})$ corresponds to the number of conjugacy class subrepresentatives of $G^{(2n)}(\widetilde F_{\overline\omega_{j,m_j}}\times \widetilde F_{\omega_{j,m_j}})$.

Consequently, we have

$$\left|G^{(2n)}(\widetilde F_{\overline\omega_{j,m_j}}\times \widetilde F_{\omega_{j,m_j}}):N_{G^{(2n)}}(P^{(2n)}(\widetilde F_{\overline\omega^1_j}\times \widetilde F_{\omega^1_j}))\right|=j$$

where j is also the global residue degree of the j-th biplace $\overline\omega_j\times\omega_j$ taking into account the inverse morphism $F^{-1}_{g_{j'}}$. ■

2.4.3 Proposition

Let $g^{(2n)}_{R\times L}[j,m_j]$, $m_j\in I\!N$, $m_j\ge 0$, *denote the* j*-th diagonal or off-diagonal conjugacy class representative of the bilinear algebraic semigroup* $G^{(2n)}(\widetilde F_{\overline\omega}\times \widetilde F_\omega)$ *and let* $g^{(2n)}_{R\times L}[j,m_j,j']$, $1\le j'\le j$, *be the* j'*-th conjugacy subclass representative of* $g^{(2n)}_{R\times L}[j,m_j]$.

Let $D^{(2n)}(\widetilde{F}^D_{\overline{\omega}} \times \widetilde{F}^D_{\omega})$ *denote the bilinear algebraic semigroup* $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$ *restricted to the product of the sets:*

$$\widetilde{F}^D_{\overline{\omega}} = \{\widetilde{F}_{\omega_1}, \cdots, \widetilde{F}_{\omega_j}, \cdots, \widetilde{F}_{\omega_r}\} \qquad \text{and} \qquad \widetilde{F}^D_{\omega} = \{\widetilde{F}_{\overline{\omega}_1}, \cdots, \widetilde{F}_{\overline{\omega}_j}, \cdots, \widetilde{F}_{\overline{\omega}_r}\}$$

with $m_j = 0$ *.*
Then:

1. *the* j*-th equivalent off-diagonal conjugacy class representatives* $g^{(2n)}_{R\times L}[j,m_j] \in G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$ *are generated from the diagonal conjugacy class representatives* $g^{(2n)}_{R\times L}[j,m_j=0] \in D^{(2n)}(\widetilde{F}^D_{\overline{\omega}} \times \widetilde{F}^D_{\omega})$ *under the product of the actions* $\mathbb{W}_{\widetilde{F}_{\overline{\omega}_j}} \times \mathbb{W}_{\widetilde{F}_{\omega_j}}$ *of the Weyl groups* $\mathbb{W}_{\widetilde{F}_{\overline{\omega}}}$ *and* $\mathbb{W}_{\widetilde{F}_{\omega}}$ *.*

2. *the Weyl groups* $\mathbb{W}_{\widetilde{F}_{\overline{\omega}}}$ *and* $\mathbb{W}_{\widetilde{F}_{\omega}}$ *acting in a nilpotent way respectively on the* j*-th right and left diagonal conjugacy class representatives* $g^{(2n)}_R[j,m_j=0]$ *and* $g^{(2n)}_L[j,m_j=0] \in g^{(2n)}_{R\times L}[j,m_j=0]$ *are the Coxeter groups* A_{j-1} *(and possibly* B_{j-1} *and* D_{j-1} *).*

Proof:

1. The equivalent off-diagonal conjugacy class representatives $g^{(2n)}_{R\times L}[j,m_j]$, corresponding to the $\widetilde{B}_{\overline{\omega}_{j,m_j}} \otimes \widetilde{B}_{\omega_{j,m_j}}$-subbisemimodules $\widetilde{M}_{\overline{\omega}_{j,m_j}} \otimes \widetilde{M}_{\omega_{j,m_j}}$ according to section 2.4.1, proposition 2.1.4 and definition 2.2.4, are isomorphic to the products, right by left, of $2n$ dimensional maximal semitori as it will be seen below.

   So, the generation of the off-diagonal conjugacy class representatives $g^{(2n)}_{R\times L}[j,m_j]$ from the diagonal conjugacy class representatives $g^{(2n)}_{R\times L}[j,m_j=0]$, which are also isomorphic to the products, right by left, of $2n$-dimensional semitori, is obtained in a classical way under the product $\mathbb{W}_{\widetilde{F}_{\overline{\omega}_j}} \times \mathbb{W}_{\widetilde{F}_{\omega_j}}$ of the Weyl groups.

2. From point 2), it results that the Coxeter group A_{j-1} (and B_{j-1} and D_{j-1}) is directly related to the integer j which corresponds to the number of equivalent conjugacy subclasses $g^{(n)}_{R\times L}[j,m_j=0,j']$ of the diagonal conjugacy class $g^{(n)}_{R\times L}[j,m_j=0]$. ■

2.4.4 Proposition

Let

$$\mathrm{Out}(G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})) = \mathrm{Aut}(G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})\Big/ \mathrm{Int}(G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega}))$$

be the group of Galois automorphisms of the reductive bilinear algebraic semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$ *where* $\mathrm{Int}(G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega}))$ *denotes the group of Galois inner automorphisms.*

Let $\mathrm{Aut}(P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1}))$ *denote the group of the Galois automorphisms of the bilinear parabolic subsemigroup* $P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})$ *.*
Then, it is proved that:

$$\mathrm{Int}(G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_{\omega}))=\mathrm{Aut}(P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1}))$$

which implies that explicit irreducible $2n$*-dimensional representations exist:*

$$\begin{aligned}\mathrm{Rep}^{(2n)}_{I_{F_{R\times L}}}\ :&\quad I_{F_R}\times I_{F_L}\ \longrightarrow\ P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})\ ,\\ \mathrm{Rep}^{(2n)}_{I^D_{F_{R\times L}}}\ :&\quad I^D_{F_R}\times I^D_{F_L}\ \longrightarrow\ P^{(2n)}(\widetilde F^D_{\overline\omega^1}\times\widetilde F^D_{\omega^1})\ ,\end{aligned}$$

from the product, right by left, of inertia subgroups to the corresponding bilinear parabolic subsemigroups where:

- $I_{F_L}:\bigoplus\limits_j I_{F^D_{\omega_j}}\bigoplus\limits_{m_j} I_{F_{\omega_{j,m_j}}}$
- $I^D_{F_L}=\bigoplus\limits_j I_{F^D_{\omega_j}}$ *according to section 1.1.7.*

Proof:

1. The bilinear parabolic subsemigroups $P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})$ and $P^{(2n)}(\widetilde F^D_{\overline\omega^1}\times\widetilde F^D_{\omega^1})$ are defined respectively over the products $\widetilde F_{\overline\omega^1_{j,m_j}}\times\widetilde F_{\omega^1_{j,m_j}}$ and $\widetilde F_{\overline\omega^1_j}\times\widetilde F_{\omega^1_j}$ of irreducible central extensions which constitute the set of fixed points respectively of the products $\widetilde F_{\overline\omega_{j,m_j}}\times\widetilde F_{\omega_{j,m_j}}$ and $\widetilde F_{\overline\omega_j}\times\widetilde F_{\omega_j}$ of the pseudo-ramified extensions of $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_{\omega})$.

   Then, the group of automorphisms $\mathrm{Aut}(P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1}))$ of the reductive bilinear semigroup $P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})$ is the group of automorphisms of the fixed points of $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_{\omega})$.

   Indeed, the map

   $$\sigma_{P^{(2n)}}\ :\quad P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})\ \longrightarrow\ P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})$$

   given by

   $$\sigma_{P^{(2n)}}(x_{p_R}\cdot x_{p_L})=p_R\ x_{p_R}\ x_{p_L}\ p_L=x'_{p_R}\ x'_{p_L}$$

   for all $x_{p_R}\in P^{(2n)}(\widetilde F_{\overline\omega^1})$, $x_{p_L}\in P^{(2n)}(\widetilde F_{\omega^1})$,
   is an automorphism of $P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})$ induced by $p_R\in P^{(2n)}(\widetilde F_{\overline\omega^1})$ and $p_L\in P^{(2n)}(\widetilde F_{\omega^1})$, in such a way that $\sigma_{P^{(2n)}}\in\mathrm{Aut}(P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1}))$.

   Similarly, the map

   $$\sigma^{\mathrm{int}}_{G^{(2n)}}\ :\quad G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_{\omega})\ \longrightarrow\ G^{2n)}(\widetilde F_{\overline\omega}\times\widetilde F_{\omega})$$

given by

$$\sigma^{\rm int}_{G^{(2n)}}(p_R\ x_{g_R}\ x_{g_L}\ p_L)=x_{g_R}\cdot x_{g_L}\ ,$$

for $x_{g_R}\in G^{(2n)}(\widetilde F_{\overline\omega})$ and $x_{g_L}\in G^{(2n)}(\widetilde F_{\omega})$, and verifying $x_{g_R}\cdot x_{g_L}=x'_{p_R}\cdot x'_{p_L}$,
is an inner automorphism of $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_{\omega})$ induced by $p_R\in P^{(2n)}(\widetilde F_{\overline\omega^1})$ and by $p_L\in P^{(2n)}(\widetilde F_{\omega^1})$, in such a way that $\sigma^{\rm int}_{G^{(2n)}}=\sigma_{P^{(2n)}}$ for $\sigma^{\rm int}_{G^{(2n)}}\in{\rm Int}(G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_{\omega}))$.

This leads to:

$${\rm Int}(G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_{\omega}))={\rm Aut}(P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1}))\ .$$

2. On the other hand, the bilinear parabolic subsemigroup

$$P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})=T^t_n(\widetilde F_{\overline\omega^1})\times T_n(\widetilde F_{\omega^1})$$

is constructed from the product of Borel subgroups $T^t_n(\widetilde F_{\overline\omega^1})$ and $T_n(\widetilde F_{\omega^1})$ which are solvable.

And, as it is known [Ser2] that every solvable finite group is Galois, it is clear that ${\rm Aut}(P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1}))$ constitutes the n-dimensional analogue of the product, right by left, of inertia subgroups. Finally, as ${\rm Aut}(P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1}))$ is isomorphic to $P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})$, the bilinear parabolic subsemigroup $P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})$ constitutes a $2n$-dimensional representation ${\rm Rep}^{(2n)}_{I_{F_{R\times L}}}$ of the product $I_{F_R}\times I_{F_L}$ of inertia subgroups in ${\rm Gal}(\widetilde F^{ac}_R/k)\times{\rm Gal}(\widetilde F^{ac}_L/k)$. ■

2.4.5 Proposition

Let $P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})$ be the bilinear parabolic affine subsemigroup of the smooth reductive bilinear affine semigroup $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_{\omega})$. Then, $P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})$ can be considered as the unitary irreducible $2n$-dimensional representation space of the algebraic bilinear semigroup of matrices ${\rm GL}_n(\widetilde F_{\overline\omega}\times\widetilde F_{\omega})$.

Proof: According to proposition 2.4.4,

$${\rm Int}(G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_{\omega}))={\rm Aut}(P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1}))$$

since we have:

$$\sigma^{\rm int}_{G^{(n)}}(p_R\ x_{g_R}\ x_{g_L}\ p_L)=x_{g_R}\cdot x_{g_L}\ ,$$

for $p_R\in P^{(2n)}(\widetilde F_{\overline\omega^1})$, $p_L\in P^{(2n)}(\widetilde F_{\omega^1})$, $x_{g_R}\in G^{(2n)}(\widetilde F_{\overline\omega})$, $x_{g_L}\in G^{(2n)}(\widetilde F_{\omega})$.
So, $P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})$ is the bilinear isotropy subgroup of $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_{\omega})$ since it is the subgroup fixing the bielements of $G^{(2n)}(\widetilde F_{\overline\omega}\times\widetilde F_{\omega})$.
Consequently, $P^{(2n)}(\widetilde F_{\overline\omega^1}\times\widetilde F_{\omega^1})$ is a unitary representation of ${\rm GL}_n(\widetilde F_{\overline\omega}\times\widetilde F_{\omega})$. ■

2.4.6 Proposition

Let $D^{(2n)}(\widetilde{F}^{D}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}^{D}_{\omega_{\oplus}})$ *denote the diagonal bilinear algebraic subsemigroup of* $G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}})$ *. Then, there exist explicit irreducible* $2n$*-dimensional representations:*

$$\begin{aligned}
\mathrm{Rep}^{(2n)}_{\mathrm{Gal}_{\widetilde{F}_{R\times L}}} &: \quad \mathrm{Gal}(\widetilde{F}^{ac}_{R}/k)\times\mathrm{Gal}(\widetilde{F}^{ac}_{L}/k) \longrightarrow G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}})\ ,\\
\mathrm{Rep}^{(2n)}_{\mathrm{Gal}^{D}\,\widetilde{F}_{R\times L}} &: \quad \mathrm{Gal}^{D}(\widetilde{F}^{ac}_{R}/k)\times\mathrm{Gal}^{D}(\widetilde{F}^{ac}_{L}/k) \longrightarrow D^{(2n)}(\widetilde{F}^{D}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}^{D}_{\omega_{\oplus}})
\end{aligned}$$

where $\widetilde{F}^{D}_{\omega_{\oplus}}=\bigoplus\limits_{j}\widetilde{F}_{\omega_j}$ *(resp.* $\widetilde{F}^{D}_{\overline{\varpi}_{\oplus}}=\bigoplus\limits_{j}\widetilde{F}_{\overline{\varpi}_j}$ *) are constructed from* $\widetilde{F}_{\omega}$ *(resp.* $\widetilde{F}_{\overline{\varpi}}$ *) introduced in section 1.1.6.*

Proof: If a group G is quasi split, in the sense that it contains a Borel subgroup, then, the homomorphism

$$\mathrm{Gal}(\overline{\mathbb{Q}}\big/\mathbb{Q}) \longrightarrow \mathrm{Out}(G)=\mathrm{Aut}(G)\big/\mathrm{Int}(G)$$

always exists [Art3].

Transposed into a bilinear context for the bilinear reductive algebraic semigroup $G^{(2n)}(\widetilde{F}_{\overline{\varpi}}\times\widetilde{F}_{\omega})$, this homomorphism becomes:

$$\mathrm{Gal}(\widetilde{F}^{ac}_{R}/k)\times\mathrm{Gal}(\widetilde{F}^{ac}_{L}/k) \longrightarrow \mathrm{Out}(G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}}))$$

where

$$\mathrm{Out}(G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}}))=\mathrm{Aut}(G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}}))\big/\mathrm{Int}(G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}}))\ .$$

Indeed, as $G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}})$ is generated from the product of Borel subgroups which are solvable, it is of Galois type according to proposition 2.4.3. And, $\mathrm{Aut}(G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}}))$ constitutes the $2n$-dimensional analogue of the product $\mathrm{Gal}(\widetilde{F}^{ac}_{R}/k)\times\mathrm{Gal}(\widetilde{F}^{ac}_{L}/k)$ of Galois groups. But, as the bilinear algebraic semigroup $G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}})$ is pseudo-ramified, it is $\mathrm{Out}(G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}}))$ which must be considered as the $2n$-dimensional analogue of $\mathrm{Gal}(\widetilde{F}^{ac}_{R}/k)\times\mathrm{Gal}(\widetilde{F}^{ac}_{L}/k)$ since $\mathrm{Int}(G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}}))=\mathrm{Aut}(P^{(2n)}(\widetilde{F}_{\overline{\varpi}^{1}_{\oplus}}\times\widetilde{F}_{\omega^{1}_{\oplus}}))$ according to proposition 2.4.3.

So,

$$\mathrm{Out}(G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}}))\simeq\prod_{n}(\mathrm{Gal}(\widetilde{F}^{ac}_{R}/k)\times\mathrm{Gal}(\widetilde{F}^{ac}_{L}/k))$$

which implies the commutative diagram:

$$\begin{tikzcd}
\mathrm{Gal}(\widetilde{F}^{ac}_{R}/k)\times\mathrm{Gal}(\widetilde{F}^{ac}_{L}/k) \arrow[r] \arrow[dr] & \mathrm{Out}(G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}})) \arrow[d, "\wr"] \\
 & G^{(2n)}(\widetilde{F}_{\overline{\varpi}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}})
\end{tikzcd}$$

Thus, as $\mathrm{Out}(G^{(2n)}(\widetilde{F}_{\overline{\omega}_\oplus}\times\widetilde{F}_{\omega_\oplus}))$ is isomorphic to $G^{(2n)}(\widetilde{F}_{\overline{\omega}_\oplus}\times\widetilde{F}_{\omega_\oplus})$, the irreducible $2n$-dimensional representation $\mathrm{Rep}^{(2n)}_{\mathrm{Gal}_{\widetilde{F}_{R\times L}}}$ of $\mathrm{Gal}(\widetilde{F}^{ac}_R/k)\times\mathrm{Gal}(\widetilde{F}^{ac}_L/k)$ is given by the reductive bilinear algebraic semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}_\oplus}\times\widetilde{F}_{\omega_\oplus})$. ■

2.4.7 Proposition

Let

$$W^{ab}_{F_R}\times W^{ab}_{F_L}=\mathrm{Gal}(\widetilde{\dot{F}}^{ac}_R/k)\times\mathrm{Gal}(\widetilde{\dot{F}}^{ac}_L/k)\subset\mathrm{Gal}(\widetilde{F}^{ac}_R/k)\times\mathrm{Gal}(\widetilde{F}^{ac}_L/k)$$

be the product of global Weil groups as introduced in definition 1.1.9.
Then, the $2n$-dimensional irreducible representation of the product of global Weil groups is given by:

$$\mathrm{Irr}\ W^{(2n)}_{F_{R\times L}}:\quad W^{ab}_{F_R}\times W^{ab}_{F_L}\longrightarrow G^{(2n)}(\widetilde{F}_{\overline{\omega}_\oplus}\times\widetilde{F}_{\omega_\oplus})\ .$$

Proof: This directly follows from $\mathrm{Rep}^{(2n)}_{\mathrm{Gal}_{\widetilde{F}_{R\times L}}}$ introduced in proposition 2.4.6, implying that $\widetilde{F}_{\overline{\omega}_\oplus}$ and $\widetilde{F}_{\omega_\oplus}$ bear here on extensions characterized by extension degrees $d=0\mod N$. ■

2.5 Trace formulas

2.5.1 Lemma

Let $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$ be a reductive bilinear complete semigroup. As it is solvable, the set of its conjugacy classes forms an increasing sequence:

$$g^{(2n)}_{R\times L}[1,m_1]\subset g^{(2n)}_{R\times L}[2,m_1]\subset\cdots\subset g^{(2n)}_{R\times L}[j,m_j]\subset\cdots\subset g^{(2n)}_{R\times L}[r,m_r]$$

characterized by the increasing ranks of their conjugacy class representatives $g^{(2n)}_{R\times L}[j,m_j]$, $1\le j\le r$.

Proof: According to definition 2.3.1 and proposition 2.1.4, the pseudo-ramified lattice bisemispace

$$X_{S_{R\times L}}=\mathrm{GL}_n(\widetilde{F}_R\times\widetilde{F}_L)\Big/\mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)$$

is homomorphic to the pseudo-ramified $B_{\overline{\omega}_\oplus}\times B_{\omega_\oplus}$-bisemimodule $M_{R_\oplus}\otimes M_{L_\oplus}$ developing (see proposition 2.1.4) as follows:

$$M_{R_\oplus}\otimes M_{L_\oplus}=\mathop{\oplus}_j\mathop{\oplus}_{m_j}\left(M_{\overline{\omega}_{j,m_j}}\otimes M_{\omega_{j,m_j}}\right)\ .$$

As the cosets of $X_{S_{R\times L}}$ correspond to the conjugacy class representatives of $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$ and as $(M_R\otimes M_L)$ constitutes the representation space of $\mathrm{GL}_n(F_{\overline{\omega}}\times F_\omega)$, each conjugacy class representative $g^{(2n)}_{R\times L}[j,m_j]$ is in fact a $\mathrm{GL}_n(F_{\overline{\omega}_{j,m_j}}\times F_{\omega_{j,m_j}})$-subbisemimodule $M_{\overline{\omega}_{j,m_j}}\otimes M_{\omega_{j,m_j}}$ characterized by a rank $r_{\overline{\omega}_j\times\omega_j}=(j\cdot m^{(j)}\cdot N)^{2n}$ according to proposition 2.3.4.
If we consider the m_{j+1}-th representative $g^{(2n)}_{R\times L}[j+1,m_{j+1}]$ of the $(j+1)$-th conjugacy class of $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$, we see that it has a rank given by

$$r_{\overline{\omega}_{j+1}\times\omega_{j+1}}=(j+1)^{2n}\cdot N^{2n}\cdot(m^{(j+1)})^{2n}$$

which is superior to the rank $r_{\overline{\omega}_j\times\omega_j}$ of $g^{(2n)}_{R\times L}[j,m_j]$. ■

2.5.2 Tower of sums of conjugacy class representatives

Remark first that a conjugacy class representative $g^{(2n)}_{R\times L}[j,m_j]$ of $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$ is in fact a bilinear complete subsemigroup noted $G^{(2n)}(F_{\overline{\omega}_{j,m_j}}\times F_{\omega_{j,m_j}})$. Then, we can introduce a tower of embedded bilinear complete subsemigroups on $G^{(2n)}(F_{\overline{\omega}_\oplus}\times F_{\omega_\oplus})$:

$$G^{(2n)}(F_{\overline{\omega}_1}\times F_{\omega_1})\subset\cdots\subset\bigoplus_{j=1}^{j}\bigoplus_{m_j}G^{(2n)}(F_{\overline{\omega}_{j,m_j}}\times F_{\omega_{j,m_j}})\subset\cdots\subset\bigoplus_{j=1}^{r}\bigoplus_{m_j}G^{(2n)}(F_{\overline{\omega}_{j,m_j}}\times F_{\omega_{j,m_j}})$$

corresponding to increasing sums of conjugacy class representatives of $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$.

2.5.3 The bialgebra of bifunctions on $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$

Now, we consider the set of all smooth continuous bifunctions on $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$ which is a bilinear complete semigroup. Consequently, we have to envisage the set $\widehat{G}^{(2n)}_R(F_{\overline{\omega}})$ of smooth continuous functions $\phi^{(2n)}_{G_R}(x_{g_R})$, $x_{g_R}\in G^{(2n)}(F_{\overline{\omega}})$, on $G^{(2n)}(F_{\overline{\omega}})\simeq T^t_n(F_{\overline{\omega}})$ and localized in the lower half space as well as the corresponding set $\widehat{G}^{(2n)}_L(F_\omega)$ of smooth continuous functions $\phi^{(2n)}_{G_L}(x_{g_L})$, $x_{g_L}\in G^{(2n)}(F_\omega)$, on $G^{(2n)}(F_\omega)\simeq T_n(F_\omega)$ and localized in the upper half space in such a way that we have on $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$ the product $\widehat{G}^{(2n)}(F_{\overline{\omega}})\times\widehat{G}^{(2n)}(F_\omega)$ of the tensor products $\phi^{(2n)}_{G_R}(x_{g_R})\otimes\phi^{(2n)}_{G_L}(x_{g_L})$ of the smooth continuous functions $\phi^{(2n)}_{G_R}(x_{g_R})$ and $\phi^{(2n)}_{G_L}(x_{g_L})$.
$\widehat{G}^{(2n)}_R(F_{\overline{\omega}})$ (resp. $\widehat{G}^{(2n)}_L(F_\omega)$) is the coordinate (semi)ring $k[G^{(2n)}(F_{\overline{\omega}})]$ (resp. $k[G^{(2n)}(F_\omega)]$) of $G^{(2n)}(F_{\overline{\omega}})$ (resp. $G^{(2n)}(F_{\overline{\omega}})$).
But, as $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$ is partitioned into conjugacy classes, we can consider bifunctions on the conjugacy class representatives of $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$.
So, let $\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}})\otimes\phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}})$ denote a bifunction on $G^{(2n)}(F_{\overline{\omega}_{j,m_j}}\times F_{\omega_{j,m_j}})$. Then, the set of smooth continuous bifunctions $\{\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}})\otimes\phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}})\}_{j=1}^{r}$ on $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$ forms a bialgebra

$$\widehat{G}^{(2n)}(F_{\overline{\omega}}\times F_\omega)=\widehat{G}^{(2n)}(F_{\overline{\omega}})\otimes\widehat{G}^{(2n)}(F_\omega)$$

where $\widehat{G}^{(2n)}(F_{\overline{\omega}})$ is the coalgebra of $\widehat{G}^{(2n)}(F_\omega)$.

2.5.4 The bialgebras $L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}}\times F_\omega))$ and $L^2_{L\times L}(G^{(2n)}(F_\omega\times F_\omega))$

The algebra (resp. coalgebra) $\widehat{G}^{(2n)}(F_\omega)$ (resp. $\widehat{G}^{(2n)}(F_{\overline{\omega}})$) is the set of all continuous complex-valued measurable functions on $G^{(2n)}(F_\omega)$ (resp. $G^{(2n)}(F_{\overline{\omega}})$) satisfying

$$\int_{G^{(2n)}(F_\omega)} \left|\phi^{(2n)}_{G_L}(x_{g_L})\right| \ dx_{g_L} < \infty \ ,$$

$$\text{(resp.} \quad \int_{G^{(2n)}(F_{\overline{\omega}})} \left|\phi^{(2n)}_{G_R}(x_{g_R})\right| \ dx_{g_R} < \infty \ ),$$

with respect to a unique Haar measure on $G^{(2n)}(F_\omega)$ (resp. $G^{(2n)}(F_{\overline{\omega}})$): it is noted $L^1_L(G^{(2n)}(F_\omega))$ (resp. $L^1_R(G^{(2n)}(F_{\overline{\omega}}))$).
The bialgebra $\widehat{G}^{(2n)}(F_{\overline{\omega}}\times F_\omega)$ of all continuous complex-valued measurable bifunctions $\phi^{(2n)}_{G_R}(x_{g_R})\otimes\phi^{(2n)}_{G_L}(x_{g_L})$ on $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$ satisfying

$$\int_{G^{(2n)}(F_{\overline{\omega}}\otimes F_\omega)} \left|\phi^{(2n)}_{G_R}(x_{g_R})\otimes\phi^{(2n)}_{G_L}(x_{g_L})\right| \ dx_{g_R}\ dx_{g_L} < \infty$$

is also noted $L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}}\times F_\omega))$.
On the other hand, let

$$\begin{aligned} i_{G_{R\to L}} \quad L^1_R(G^{(2n)}(F_{\overline{\omega}})) &\longrightarrow L^1_L(G^{(2n)}(F_\omega)) \\ \phi^{(2n)}_{G_R}(x_{g_R}) &\longrightarrow \phi^{(2n)}_{G_L}(x_{g_L}) \end{aligned}$$

be the involution mapping each continuous function $\phi^{(2n)}_{G_R}(x_{g_R}) \in L^1_R(G^{(2n)}(F_{\overline{\omega}}))$, restricted to the lower half space, to the corresponding continuous function $\phi^{(2n)}_{G_L}(x_{g_L}) \in L^1_L(G^{(2n)}(F_\omega))$, restricted to the upper half space.
Then, the involution:

$$\begin{aligned} i_{G_{R\to L}} : \quad & L^1_R(G^{(2n)}(F_{\overline{\omega}}))\times L^1_L(G^{(2n)}(F_\omega)) = L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}}\times F_\omega)) \\ & \qquad \longrightarrow L^2_{L\times L}(G^{(2n)}(F_\omega\times F_\omega)) \\ & \phi^{(2n)}_{G_R}(x_{g_R})\otimes\phi^{(2n)}_{G_L}(x_{g_L}) \longrightarrow \phi^{(2n)}_{G_L}(x_{g_L})\otimes\phi^{(2n)}_{G_L}(x_{g_L}) \end{aligned}$$

has the property that the continuous bifunctions $\phi^{(2n)}_{G_R}(x_{g_R})\otimes\phi^{(2n)}_{G_L}(x_{g_L}) \in L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}}\times F_\omega))$ are transformed into the bifunctions $\phi^{(2n)}_{G_L}(x_{g_L})\otimes\phi^{(2n)}_{G_L}(x_{g_L}) \in L^2_{L\times L}(G^{(2n)}(F_\omega\times F_\omega))$ satisfying

$$\int_{G^{(2n)}(F_\omega\times F_\omega)} \left|\phi^{(2n)}_{G_L}(x_{g_L})\right|^2 \ dx_{g_L} < \infty \ .$$

Thus, $L^2_{L\times L}(G^{(2n)}(F_\omega \times F_\omega))$ is the standard space of square integrable continuous functions on $G^{(2n)}(F_\omega \times F_\omega)$ restricted to the upper half space and defined with respect to a fixed Haar measure.

2.5.5 Proposition

The bialgebra $\widehat{G}^{(2n)}(F_{\overline{\omega}} \times F_\omega)$ of smooth continuous complex-valued measurable bifunctions on the locally compact bilinear complete semigroup $G^{(2n)}(F_{\overline{\omega}} \times F_\omega)$ is composed of the increasing sequence:

$$\phi^{(2n)}_{G_{1_R}}(x_{g_{1_R}}) \otimes \phi^{(2n)}_{G_{1_L}}(x_{g_{1_L}}) \subset \cdots \subset \phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}}) \otimes \phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}}) \subset \cdots \subset \phi^{(2n)}_{G_{r_R}}(x_{g_{r_R}}) \otimes \phi^{(2n)}_{G_{r_L}}(x_{g_{r_L}})$$

of these bifunctions in such a way that the j-th bifunction $\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}}) \otimes \phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}})$ is defined on the closed bilinear subsemigroup $G^{(2n)}(F_{\overline{\omega}_{j,m_j}} \times F_{\omega_{j,m_j}}) \equiv g^{(2n)}_{R\times L}[j,m_j]$ corresponding to the j-th conjugacy class of $G^{(2n)}(F_{\overline{\omega}} \times F_\omega)$.

Sketch of proof : This results from lemma 2.5.1 and from section 2.5.3. ■

2.5.6 Proposition

The bialgebra $\widehat{G}^{(2n)}(F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus}) \equiv L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus}))$ of bifunctions on $G^{(2n)}(F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus})$ is composed of a tower:

$$\begin{aligned}\phi^{(2n)}_{G_{1_R}}(x_{g_{1_R}}) \otimes \phi^{(2n)}_{G_{1_L}}(x_{g_{1_L}}) \subset \cdots \subset \bigoplus_{j=1}^{j} \left(\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}}) \otimes \phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}})\right) \\ \subset \cdots \subset \bigoplus_{j=1}^{r} \left(\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}}) \otimes \phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}})\right)\end{aligned}$$

of the partial sums of these bifunctions $\bigoplus_{j=1}^{j} \left(\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}}) \otimes \phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}})\right)$ on the corresponding sums of conjugacy class representatives of $G^{(2n)}(F_{\overline{\omega}} \times F_\omega)$.

Sketch of proof : This is a consequence of the introduction of a tower of sums of conjugacy class representatives of $G^{(2n)}(F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus})$ given in section 2.5.2.

2.5.7 The bialgebra $L^{1-1}_{R\times L}(G^{(2n)}(F^{nr}_{\overline{\omega}}\times F^{nr}_{\omega}))$ of bifunctions on the pseudo-unramified bilinear complete semigroup $G^{(2n)}(F^{nr}_{\overline{\omega}}\times F^{nr}_{\omega})$

Similarly, on the set of the conjugacy class representatives of the pseudo-unramified bilinear complete semigroup $G^{(2n)}(F^{nr}_{\overline{\omega}}\times F^{nr}_{\omega})$, an increasing sequence

$$\phi^{(2n)}_{G^{nr}_{1_R}}(x^{nr}_{g_{1_R}})\otimes\phi^{(2n)}_{G^{nr}_{1_L}}(x^{nr}_{g_{1_L}})\subset\cdots\subset\phi^{(2n)}_{G^{nr}_{j_R}}(x^{nr}_{g_{j_R}})\otimes\phi^{(2n)}_{G^{nr}_{j_L}}(x^{nr}_{g_{j_L}})$$

$$\subset\cdots\subset\phi^{(2n)}_{G^{nr}_{r_R}}(x^{nr}_{g_{r_R}})\otimes\phi^{(2n)}_{G^{nr}_{r_L}}(x^{nr}_{g_{r_L}})$$

of smooth continuous complex-valued measurable bifunctions can be introduced in such a way that they satisfy:

$$\int_{G^{(2n)}(F^{nr}_{\overline{\omega}}\times F^{nr}_{\omega})}\left|\phi^{(2n)}_{G^{nr}_{j_R}}(x^{nr}_{g_{j_R}})\otimes\phi^{(2n)}_{G^{nr}_{j_L}}(x^{nr}_{g_{j_L}})\right|dx^{nr}_{g_{j_R}}\ dx^{nr}_{g_{j_L}}<\infty\ .$$

Then, the bialgebra of these bifunctions on $G^{(2n)}(F^{nr}_{\overline{\omega}}\times F^{nr}_{\omega})$ is written $\widehat{G}^{(2n)}(F^{nr}_{\overline{\omega}}\times F^{nr}_{\omega})$ or $L^{1-1}_{R\times L}(G^{(2n)}(F^{nr}_{\overline{\omega}}\times F^{nr}_{\omega}))$.

2.5.8 The bialgebra $L^{1-1}_{R\times L}(P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1}))$ of bifunctions on the complete parabolic semigroup $P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1})$

As $P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1})$ is defined over the products $F_{\overline{\omega}^1_{j,m_j}}\times F_{\omega^1_{j,m_j}}$ of the irreducible completions of the completions $F_{\overline{\omega}_{j,m_j}}\times F_{\omega_{j,m_j}}$ at all infinite places $\overline{\omega}_j\times\omega_j$ of $F_R\times F_L$, a bialgebra $L^{1-1}_{R\times L}(P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1}))$ of smooth continuous complex-valued measurable bifunctions $\phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}})\otimes\phi^{(2n)}_{P_{j_L}}(x_{p_{j_L}})$, $1\le j\le r$, on $P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1})$ can be introduced in such a way that:

$$\int_{P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1})}\left|\phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}})\otimes\phi^{(2n)}_{P_{j_L}}(x_{p_{j_L}})\right|dx_{p_{j_R}}\ dx_{p_{j_L}}<\infty\ .$$

Each bifunction $\phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}})\otimes\phi^{(2n)}_{P_{j_L}}(x_{p_{j_L}})$ is defined on the irreducible equivalence class representative $P^{(2n)}(F_{\overline{\omega}^1_{j,m_j}}\times F_{\omega^1_{j,m_j}})$ characterized by a rank $r_{P_{\overline{\omega}_j\times\omega_j}}=(N\cdot m^{(j)})^{2n}$.
The normalized rank $r_{/N_{P_{\overline{\omega}_j\times\omega_j}}}$ of $P^{(2n)}(F_{\overline{\omega}^1_{j,m_j}}\times F_{\omega^1_{j,m_j}})$ is defined by:

$$r_{/N_{P_{\overline{\omega}_j\times\omega_j}}}=r_{P_{\overline{\omega}_j\times\omega_j}}\Big/(N\cdot m^{(j)})^{2n}=1\ .$$

A character on $G^{(2n)}(F_{\overline{\omega}}\times F_{\omega})$ is then defined to be a bifunction $\chi_{j_{R\times L}}=\phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}})\otimes\phi^{(2n)}_{P_{j_L}}(x_{p_{j_L}})$ from the irreducible equivalence class representative $P^{(2n)}(F_{\overline{\omega}^1_{j,m_j}}\times F_{\omega^1_{j,m_j}})$ of $G^{(2n)}(F_{\overline{\omega}}\times F_{\omega})$, having as normalized rank $r_{/N_{P_{\overline{\omega}_j\times\omega_j}}}$, into $\mathbb{C}$.
And, the group of characters $\chi_{j_{R\times L}}$ of $P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1})$ is written $X(P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1}),\mathbb{C})$.

2.5.9 Two types of trace formulas: an introduction

Two kinds of trace formulas will now be developed [Kot1, Art1]. The first one is an adaptation of the Selberg-Arthur trace formula [Sel, Art2] of an operator associated with the quotient of a semisimple Lie group and a discrete subgroup in such a way that this operator corresponds to a unitary representation of the semisimple Lie group $G(\mathbb{A})$. The adaptation considered here concerns the trace of the operator associated with the bilinear parabolic semigroup $P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1})$ envisaged as the unitary representation of the complete bilinear semigroup $\mathrm{GL}_n(F_{\overline{\omega}} \times F_{\omega})$. This operator acts by convolution on the bialgebra $L^{1-1}_{R\times L}(G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega}))$ of bifunctions on the pseudo-unramified bilinear complete semigroup $G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega})$. As a result, this operator decomposes according to the unitary conjugacy classes of the pseudo-ramified bilinear algebraic semigroup $G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})$ and the corresponding trace formula occurs in the bialgebra $L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus}))$ of bifunctions on $G^{(2n)}(F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus})$. Notice that the complete trace formula referring to the set of irreducible representations of $G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})$ will be envisaged in chapter 4.
Remark that a trace formula, being an operation of bilinear type since it refers to the matrix representation of an operator, is well adapted to the bilinear case considered here and, especially, to the second kind of trace formulas occurring directly in the bialgebra $L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus}))$ of bifunctions on the pseudo-ramified complete bilinear semigroup $G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})$. This latter case then leads to the Plancherel formula.
Consequently, the two types of trace formulas correspond to each other.

Let us now start with the first trace formula referring to the Arthur-Selbeg trace formula.

2.5.10 Unitary representation of $\mathrm{GL}_n(F_{\overline{\omega}} \times F_{\omega})$

Let $\mathrm{GL}_n(\mathbb{Z}^2)$ be the discrete subgroup of the pseudo-unramified bilinear complete semigroup $\mathrm{GL}_n(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega})$. We then consider the bialgebra $L^{1-1}_{R\times L}(\mathrm{GL}_n(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega}) / \mathrm{GL}_n(\mathbb{Z}^2))$ of bifunctions $\phi^{(2n)}_{G^{nr}_{j_R}}(x^{nr}_{g_{j_R}}) \otimes \phi^{(2n)}_{G^{nr}_{j_L}}(x^{nr}_{g_{j_L}})$ on the conjugacy class representatives of $\mathrm{GL}_n(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega})$ according to section 2.5.7.
The conjugacy class representatives $\mathrm{GL}_n(F^{nr}_{\overline{\omega}_{j,m_j}} \times F^{nr}_{\omega_{j,m_j}})$ correspond to the cosets of $\mathrm{GL}_n(F^{nr}_{R} \times F^{nr}_{L}) / \mathrm{GL}_n(\mathbb{Z}^2)$ in such a way that the space of these cosets is locally compact.
On the other hand, let $\mathrm{GL}_n(F_{\overline{\omega}} \times F_{\omega})$ be the reductive pseudo-ramified complete bilinear semigroup of matrices and let $G^{(2n)}(F_{\overline{\omega}} \times F_{\omega}) = M_R \otimes M_L$ be its representation space, i.e. a bilinear semigroup. Then, a unitary representation of $\mathrm{GL}_n(F_{\overline{\omega}} \times F_{\omega})$ in this bilinear

semigroup $G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})$ is any unitary mapping:

$$\begin{aligned} U_{P^{(2n)}} : \quad P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1}) &\longrightarrow R(P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1})) \\ P^{(2n)}(F_{\overline{\omega}^1_{j,m_j}} \times F_{\omega^1_{j,m_j}}) &\longrightarrow R(P^{(2n)}(F_{\overline{\omega}^1_{j,m_j}} \times F^1_{\omega_{j,m_j}})) \end{aligned}$$

from the bilinear parabolic semigroup $P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1})$ to the unitary operator $R(P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1}))$ according to proposition 2.4.5.

2.5.11 Kernel of the integral operator restricted to the (j, m_j)-th conjugacy class representative of $P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1})$

Let $\phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}}) \otimes \phi^{(2n)}_{P_{j_L}}(x_{p_{j_L}}) \in L^{1-1}_{R\times L}(P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1}))$ be the bifunction on the j-th irreducible equivalence class representative of $P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1})$ and let $\phi^{(2n)}_{G^{nr}_{j_R}}(x^{nr}_{g_{j_R}}) \otimes \phi^{(2n)}_{G^{nr}_{j_L}}(x^{nr}_{g_{j_L}}) \in L^{1-1}_{R\times L}(G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega}))$ be the bifunction on the corresponding j-th equivalence class representative of the pseudo-unramified complete bilinear semigroup $G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega})$.
Then, the unitary operator $R(P^{(2n)}(F^1_{\overline{\omega}} \times F^1_{\omega}))$ acts by convolution on every bifunction $\phi^{(2n)}_{G^{nr}_{j_R}}(x^{nr}_{g_{j_R}}) \otimes \phi^{(2n)}_{G^{nr}_{j_L}}(x^{nr}_{g_{j_L}})$ on the pseudo-unramified bilinear complete semigroup $G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega})$ according to:

$$\begin{aligned} & R(P^{(2n)}(F_{\overline{\omega}^1_{j,m_j}} \times F_{\omega^1_{j,m_j}})) \cdot (\phi^{(2n)}_{G^{nr}_{j_R}}(x^{nr}_{g_{j_R}}) \otimes \phi^{(2n)}_{G^{nr}_{j_L}}(x^{nr}_{g_{j_L}})) \\ &= \int_{G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})} (\phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}}) \otimes \phi^{(2n)}_{P_{j_L}}(x_{P_{j_L}})) \\ &\quad \times (\phi^{(2n)}_{G^{nr}_{j_R}}(x^{nr}_{g_{j_R}} \cdot x_{p_{j_R}}) \otimes \phi^{(2n)}_{G^{nr}_{j_L}}(x^{nr}_{g_{j_L}} \cdot x_{p_{j_L}}))\ dx_{p_{j_R}}\ dx_{p_{j_L}} \\ &= \int_{G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})} (\phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}} \cdot (x^{nr}_{g_{j_R}})^{-1})) \otimes \phi^{(2n)}_{P_{j_L}}(x_{P_{j_L}} \cdot (x^{nr}_{g_{j_L}})^{-1}))) \\ &\quad \times (\phi^{(2n)}_{G^{nr}_{j_R}}(x_{p_{j_R}}) \otimes \phi^{(2n)}_{G^{nr}_{j_L}}(x_{p_{j_L}}))\ dx_{p_{j_R}}\ dx_{p_{j_L}} \\ &= \int_{G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})} (N_{P_{j_R}}(x_{p_{j_R}}, x^{nr}_{g_{j_R}}) \otimes N_{P_{j_L}}(x_{p_{j_L}}, x^{nr}_{g_{j_L}})) \\ &\quad \times (\phi^{(2n)}_{G^{nr}_{j_R}}(x_{p_{j_R}}) \otimes \phi^{(2n)}_{G^{nr}_{j_L}}(x_{p_{j_L}}))\ dx_{p_{j_R}}\ dx_{p_{j_L}} \end{aligned}$$

where

$$N_{P_{j_R}}(x_{p_{j_R}}, x^{nr}_{g_{j_R}}) = \phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}} \cdot (x^{nr}_{g_{j_R}})^{-1})$$

$$(\text{resp.} \quad N_{P_{j_L}}(x_{p_{j_L}}, x^{nr}_{g_{j_L}}) = \phi^{(2n)}_{P_{j_L}}\ (x_{p_{j_L}} \cdot (x^{nr}_{g_{j_L}})^{-1})\ )$$

is the right (resp. left) kernel of the integral operator

$$R_{P^{(2n)}(F_{\overline{\omega}^1})}(\phi^{(2n)}_{P_{j_R}}) = \int_{P^{(2n)}} R_{P^{(2n)}(F_{\overline{\omega}^1})}(x_{p_{j_R}})\ \phi^{(2n)}_{P_{j_R}}\ (x_{p_{j_R}})\ dx_{p_{j_R}}$$

$$(\text{resp.} \quad R_{P^{(2n)}(F_{\omega^1})}(\phi^{(2n)}_{P_{j_L}}) = \int_{P^{(2n)}} R_{P^{(2n)}(F_{\omega^1})}(x_{p_{j_L}})\ \phi^{(2n)}_{P_{j_L}}\ (x_{p_{j_L}})\ dx_{p_{j_L}}\ )$$

restricted to the (j, m_j)-th conjugacy class representative of $P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1})$.

2.5.12 Proposition

The unitary operator $R(P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1}))$ *corresponding to the bilinear parabolic semigroup* $P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1})$ *acts by convolution on the bifunction(s)* $(\phi^{(2n)}_{G^{nr}_R}(x^{nr}_{g_R}) \otimes \phi^{(2n)}_{G^{nr}_L}(x^{nr}_{p_L})$ *on the pseudo-unramified bilinear complete semigroup* $G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega})$ *according to*

$$\begin{aligned}
& R(P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1}))(\phi^{(2n)}_{G^{nr}_R}(x^{nr}_{g_R}) \otimes \phi^{(2n)}_{G^{nr}_L}(x^{nr}_{p_L})) \\
& = \int_{G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})} (N_{P_R}(x_{p_R}, x^{nr}_{g_R}) \otimes N_{P_L}(x_{p_L}, x^{nr}_{g_L}))(\phi^{(2n)}_{G^{nr}_R}(x_{p_R}) \otimes \phi^{(2n)}_{G^{nr}_L}(x_{p_L}))\ dx_{p_R}\ dx_{p_L} \\
& = \bigoplus_{j,m_j} \int_{G^{(2n)}(F_{\overline{\omega}} \times F_{\omega}) \big/ \mathrm{GL}_n(\mathbb{Z} \big/ N\mathbb{Z})^2)} (N_{P_{j_R}}(x_{p_{j_R}}, x^{nr}_{g_{j_R}}) \otimes N_{P_{j_L}}(x_{p_{j_L}}, x^{nr}_{g_{j_L}})) \\
& \qquad \times (\phi^{(2n)}_{G^{nr}_{j_R}}(x_{p_{j_R}}) \otimes \phi^{(2n)}_{G^{nr}_{j_L}}(x_{p_{j_L}}))\ dx_{p_{j_R}}\ dx_{p_{j_L}}
\end{aligned}$$

where

- $\phi^{(2n)}_{G^{nr}_R}(x_{p_R}) \otimes \phi^{(2n)}_{G^{nr}_L}(x_{p_L})) = \bigoplus\limits_{j} \bigoplus\limits_{m_j} (\phi^{(2n)}_{G^{nr}_{j_R}}(x_{p_{j_R}}) \otimes \phi^{(2n)}_{G^{nr}_{j_L}}(x_{p_{j_L}}))$;
- *the (bi)kernel of the integral (bi)operator* $R_{P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1})}(\phi^{(2n)}_{P_R} \otimes \phi^{(2n)}_{P_L})$ *decomposes:*

  $$N_{P_R}(x_{p_R}, x^{nr}_{g_R}) \otimes N_{P_L}(x_{p_L}, x^{nr}_{g_L}) = \bigoplus_{j,m_j} (N_{P_{j_R}}(x_{p_{j_R}}, x^{nr}_{g_{j_R}}) \otimes N_{P_{j_L}}(x_{p_{j_L}}, x^{nr}_{g_{j_L}})$$

  according to the conjugacy class representatives of $P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1})$.

Proof:

1. By this way, the (bi)kernel $(N_{P_R}(x_{p_R},x^{nr}_{g_R})\otimes N_{P_L}(x_{p_L},x^{nr}_{g_L}))$ is separable since it decomposes according to the conjugacy class representatives of $P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1})$ which corresponds to the cosets of $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)\big/\operatorname{GL}_n((\mathbb{Z}\,/N\ \mathbb{Z}\,)^2)$.

2. The integral $\int_{G^{(2n)}(F_{\overline{\omega}}\times F_\omega)}(N_{P_R}(x_{p_R},x^{nr}_{g_R})\otimes N_{P_L}(x_{p_L},x^{nr}_{g_L}))(\phi^{(2n)}_{G^{nr}_R}(x_{p_R})\otimes\phi^{(2n)}_{G^{nr}_L}(x_{p_L}))$ $dx_{p_R}\,dx_{p_L}$, decomposing according to the conjugacy class representatives of $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$, converges since the integral $\int_{G^{(2n)}(F_{\overline{\omega}}\times F_\omega)\big/\operatorname{GL}_n((\mathbb{Z}\,/N\mathbb{Z}\,)^2)}(N_{P_{j_R}}(x_{p_{j_R}},x^{nr}_{g_{j_R}})\otimes N_{P_{j_L}}(x_{p_{j_L}},x^{nr}_{g_{j_L}}))(\phi^{(2n)}_{G^{nr}_{j_R}}(x_{p_{j_R}})\otimes\phi^{(2n)}_{G^{nr}_{j_L}}(x_{p_{j_L}}))\ dx_{p_{j_R}}\ dx_{p_{j_L}}$ restricted to the (j,m_j)-th conjugacy class representative is bounded. ■

2.5.13 Proposition

The trace of the unitary operator $R(P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1}))$ *can be expressed as a sum of integrals of* $\phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}}\ (x^{nr}_{g_{j_R}})^{-1})\otimes_D\phi^{(2n)}_{P_{j_L}}(x_{p_{j_L}}\ (x^{nr}_{g_{j_L}})^{-1})$ *according to:*

$$\operatorname{tr}(R_{P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1})}(\phi^{(2n)}_{P_R}\otimes\phi^{(2n)}_{P_L}))$$

$$=\bigoplus_{j,m_j}\ \operatorname{vol}(P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1})\int_{P^{(2n)}}\phi^{(2n)}_{P_{j_R}}\ (x_{p_{j_R}}\ (x^{nr}_{g_{j_R}})^{-1})\otimes_D\phi^{(2n)}_{P_{j_L}}\ (x_{p_{j_L}}\ (x^{nr}_{g_{j_L}})^{-1})\ dx_{p_{j_R}}\ dx_{p_{j_L}}$$

$$=\bigoplus_{j,m_j} R_{P^{(2n)}(F_{\overline{\omega}^1_{j,mj}}\times F_{\omega^1_{j,m_j}})}(\phi^{(2n)}_{P_{j_R}},\phi^{(2n)}_{P_{j_L}})$$

where $(\cdot,\cdot)$ *is a bilinear form.*

Proof: According to proposition 2.5.12, the kernel $N_{P_R}(x_{p_R},x^{nr}_{g_R})\otimes N_{P_L}(x_{p_L},x^{nr}_{g_L})$ of the integral operator $R(P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1}))$ corresponds precisely to the action of the unitary parabolic bilinear semigroup $P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1})$ on the bifunction $\phi^{(2n)}_{G^{nr}_R}(x_{p_R})\otimes\phi^{(2n)}_{G^{nr}_L}(x_{p_L})$ on the pseudo-unramified complete bilinear semigroup $G^{(2n)}(F^{nr}_{\overline{\omega}}\times F^{nr}_\omega)$.
As a result, we have the thesis which is an adaptation of the Arthur's trace formula for reductive linear groups [Art2]. ■

2.5.14 Trace formula for the operator $R(G^{(2n)}(F_{\overline{\omega}} \times F_{\omega}))$

A representation of the pseudo-ramified bilinear complete semigroup $G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})$ is any mapping

$$\begin{aligned} R_{G^{(2n)}} : \quad G^{(2n)}(F_{\overline{\omega}} \times F_{\omega}) &\longrightarrow R(G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})) \\ G^{(2n)}(F_{\overline{\omega}_{j,m_j}} \times F_{\omega_{j,m_j}}) &\longrightarrow R(G^{(2n)}(F_{\overline{\omega}_{j,m_j}} \times F_{\omega_{j,m_j}})) \end{aligned}$$

from $G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})$ to the operator $R(G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})) \simeq \mathrm{GL}_n(F_{\overline{\omega}} \times F_{\omega})$ in such a way that the matrix $\mathrm{GL}_n(F_{\overline{\omega}_{j,m_j}} \times F_{\omega_{j,m_j}})$ corresponds to the conjugacy class representative $g^{(2n)}_{R\times L}[j,m_j] \equiv G^{(2n)}(F_{\overline{\omega}_{j,m_j}} \times F_{\omega_{j,m_j}})$ of the pseudo-ramified bilinear semigroup $G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})$.

Then, a trace formula of the operator $R(G^{(2n)}(F_{\overline{\omega}} \times F_{\omega}))$ can be reached from the trace of the unitary parabolic operator $R(P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1}))$ by considering the analytic continuation action of the (bi)kernel $N_{P_R}(x_{p_R}, x^{nr}_{g_R}) \otimes N_{P_L}(x_{p_L}, x^{nr}_{g_L})$ according to the following proposition.

2.5.15 Proposition

The trace of the unitary parabolic operator $R(P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1}))$ *can be extended to the trace of the pseudo-ramified bilinear complete semigroup* $R(G^{(2n)}(F_{\overline{\omega}} \times F_{\omega}))$ *if the action of the (bi)kernel* $N_{P_R}(x_{p_R}, x^{nr}_{g_R}) \otimes N_{P_L}(x_{p_L}, x^{nr}_{g_L})$ *of* $R(P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1}))$ *on the (pseudo-)unramified bifunction* $\phi^{(2n)}_{G^{nr}_R}(x_{p_R}) \otimes \phi^{(2n)}_{G^{nr}_L}(x_{p_L})$ *consists in the analytic continuation of this bifunction in such a way that the (pseudo-)ramified bifunction* $\phi^{(2n)}_{G_R}(x_{g_R}) \otimes \phi^{(2n)}_{G_L}(x_{g_L}) \in L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}} \times F_{\omega}))$ *on the pseudo-ramified bilinear complete semigroup* $G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})$ *verifies:*

$$\phi^{(2n)}_{G_R}(x_{g_R}) \otimes \phi^{(2n)}_{G_L}(x_{p_L}) = (N_{P_R}(x_{p_R}, x^{nr}_{g_R}) \otimes N_{P_L}(x_{p_L}, x^{nr}_{g_L}))(\phi^{(2n)}_{G^{nr}_R}(x^{nr}_{p_R}) \otimes \phi^{(2n)}_{G^{nr}_L}(x^{nr}_{p_L})) \ .$$

So, we get the following trace formula:

$$\begin{aligned} &\mathrm{tr}(R_{G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})}(\phi^{(2n)}_{G_R} \otimes \phi^{(2n)}_{G_L})) \\ &\qquad = \bigoplus_{j,m_j} \mathrm{vol}(G^{(2n)}(F_{\overline{\omega}_{\oplus}} \times F_{\omega_{\oplus}})) \int_{G^{(2n)}} \phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}}) \otimes_D \phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}}) \, dx_{g_{j_R}} \, dx_{g_{j_L}} \end{aligned}$$

where the sum ranges over the diagonal part of the conjugacy class representatives of $G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})$ *.*

Proof: The action of the bikernel

$$N_{P_R}(x_{p_R}, x^{nr}_{g_R}) \otimes N_{P_L}(x_{p_L}, x^{nr}_{g_L}) = \bigoplus_{j,m_j} (\phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}} \ (x^{nr}_{g_{j_R}})^{-1}) \otimes \phi^{(2n)}_{P_{j_L}}(x_{p_{j_L}} \ (x^{nr}_{g_{j_L}})^{-1}))$$

on the (pseudo-)unramified bifunction

$$\phi^{(2n)}_{G^{nr}_R}(x_{p_R}) \otimes \phi^{(2n)}_{G^{nr}_L}(x_{p_L}) = \mathop{\oplus}_{j,m_j} (\phi^{(2n)}_{G^{nr}_{j_R}}(x_{p_{j_R}}) \otimes \phi^{(2n)}_{G^{nr}_{j_L}}(x_{p_{j_L}})) \ ,$$

defined on the pseudo-unramified complete bilinear semigroup $G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega})$, consists in the pseudo-ramification of this bifunction according to:

$$I^{nr\to r}_{R\times L} : \quad \phi^{(2n)}_{G^{nr}_R}(x_{p_R}) \otimes \phi^{(2n)}_{G^{nr}_L}(x_{p_L}) \longrightarrow \phi^{(2n)}_{G_R}(x_{g_R}) \otimes \phi^{(2n)}_{G_L}(x_{g_L})$$

where $I^{nr\to r}_{R\times L}$ is an injective morphism to the (pseudo-)ramified bifunction $\phi^{(2n)}_{G_R}(x_{g_R}) \otimes \phi^{(2n)}_{G_L}(x_{g_L})$ on the pseudo-ramified bilinear complete semigroup $G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})$. Consequently, the injective morphism of pseudo-ramification $I^{nr\to r}_{R\times L}$ of the pseudo-unramified bifunction $\phi^{(2n)}_{G^{nr}_R}(x_{p_R}) \otimes \phi^{(2n)}_{G^{nr}_L}(x_{p_L})$ corresponds to an analytic continuation of this bifunction since the injective morphism $I^{nr\to r}_{R\times L}$ corresponds to the inverse of the map:

$$G^{(2n)}_{F\to F^{nr}} : \quad G^{(2n)}(F_{\overline{\omega}} \times F_{\omega}) \longrightarrow G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega})$$

introduced in section 2.4.1.
The volume $\mathrm{vol}(G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega}))$ of the pseudo-unramified complete bilinear semigroup $G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega})$ with respect to a fixed Haar measure is then inflated by the volume $\mathrm{vol}(P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1}))$ of the kernel $\mathrm{Ker}(G^{(2n)}_{F\to F^{nr}}) = P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1})$ of $G^{(2n)}_{F\to F^{nr}}$ into the volume $\mathrm{vol}(G^{(2n)}(F_{\overline{\omega}} \times F_{\omega}))$.
And, the trace formula $\mathrm{tr}(R_{G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})}(\phi^{(2n)}_{G_R} \otimes \phi^{(2n)}_{G_L}))$ then follows from the developments of propositions 2.5.12 and 2.5.13. ■

2.5.16 Corollary

Let $L^{1-1}_{R\times L}(G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega}))$ *denote the bialgebra of (pseudo-)unramified bifunctions on the (pseudo-)unramified bilinear complete semigroup* $G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega})$ *and let* $L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}} \times F_{\omega}))$ *be the bialgebra of (pseudo-)ramified bifunctions on the pseudo-ramified bilinear complete semigroup* $G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})$ *.*
Then, the injective morphism:

$$\begin{aligned} I^{nr\to r}_{R\times L} : \quad L^{1-1}_{R\times L}(G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega})) &\longrightarrow L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}} \times F_{\omega})) \\ \phi^{(2n)}_{G^{nr}_R}(x^{nr}_{g_R}) \otimes \phi^{(2n)}_{G^{nr}_L}(x^{nr}_{g_L}) &\longrightarrow \phi^{(2n)}_{G_R}(x_{g_R}) \otimes \phi^{(2n)}_{G_L}(x_{g_L}) \end{aligned}$$

corresponds to a pseudo-ramification morphism of the bialgebra $L^{1-1}_{R\times L}(G^{(2n)}(F^{nr}_{F_{\overline{\omega}}} \times F^{nr}_{F_{\omega}}))$ *.*

2.5.17 Proposition

Let the bifunction

$$\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}}) \otimes \phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}}) \in L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}} \times F_\omega))$$

be developed according to:

$$\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}}) \otimes \phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}}) = (\phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}}) \otimes \phi^{(2n)}_{P_{j_L}}(x_{p_{j_L}})) \times (\phi^{(2n)}_{G^{nr}_{j_R}}(x^{nr}_{g_{j_R}}) \otimes \phi^{(2n)}_{G^{nr}_{j_L}}(x^{nr}_{g_{j_L}})) \ .$$

Then, the following trace formula

$$\begin{aligned}
&\mathrm{Tr}(R_{G^{(2n)}(F_{\overline{\omega}}\times F_\omega)}(\phi^{(2n)}_{G_R} \otimes \phi^{(2n)}_{G_L}))\\
&\quad = \bigoplus_{j,m_j} \left[\mathrm{vol}(P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1})) \int_{P^{(2n)}} \phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}}) \otimes_D \phi^{(2n)}_{P_{j_L}}(x_{p_{j_L}})\ dx_{p_{j_R}}\ dx_{p_{j_L}}\right]\\
&\qquad\quad \times \left[\mathrm{vol}(G^{(2n)}(F^{nr}_{\overline{\omega}} \times F^{nr}_{\omega})) \int_{G^{(2n)}} \phi^{(2n)}_{G^{nr}_{j_R}}(x^{nr}_{g_{j_R}}) \otimes_D \phi^{(2n)}_{G^{nr}_{j_L}}(x^{nr}_{g_{j_L}})\ dx^{nr}_{g_{j_R}}\ dx^{nr}_{g_{j_L}}\right]
\end{aligned}$$

follows and relies on the Lefschetz trace formula.

Proof: The considered trace formula results from the development of the bifunction $\phi^{(2n)}_{G_{j_R}}(x_{j_R}) \otimes \phi^{(2n)}_{G_{j_L}}(x_{j_L})$ given in proposition 2.5.15 and relies on the Lefschetz trace formula because $\phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}}) \otimes \phi^{(2n)}_{P_{j_L}}(x_{p_{j_L}})$ is defined on the bilinear parabolic semigroup $P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1})$ which is the isotropy subgroup of $G^{(2n)}(F_{\overline{\omega}} \times F_\omega)$ fixing the bielements of $G^{(2n)}(F_{\overline{\omega}} \times F_\omega)$ (see proposition 2.4.5 and [Gro2]). ■

2.5.18 Proposition

Let $\mathrm{tr}(R_{P^{(2n)}(F^1_{\overline{\omega}}\times F^1_\omega)}(\phi^{(2n)}_{P_R} \otimes \phi^{(2n)}_{P_L}))$ *be given by:*

$$\begin{aligned}
&\mathrm{tr}(R_{P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1})}(\phi^{(2n)}_{P_R} \otimes \phi^{(2n)}_{P_L}))\\
&\quad = \bigoplus_j \bigoplus_{m_j} \mathrm{vol}(P^{(2n)}(F_{\overline{\omega}^1} \times F_{\omega^1})) \int_{P^{(2n)}} \phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}}) \otimes_D \phi^{(2n)}_{P_{j_L}}(x_{p_{j_L}})\ dx_{p_{j_R}}\ dx_{p_{j_L}}\\
&\quad = \bigoplus_j \bigoplus_{m_j} R_{P^{(2n)}(F_{\overline{\omega}^1_{j,m_j}} \times F_{\omega^1_{j,m_j}})}(\phi^{(2n)}_{P_{j_R}}, \phi^{(2n)}_{P_{j_L}})
\end{aligned}$$

according to proposition 2.5.13.

Then, the trace formula $\mathrm{Tr}(R_{G^{(2n)}(F_{\overline{\varpi}}\times F_\omega)}(\phi^{(2n)}_{G_R}\otimes\phi^{(2n)}_{G_L}))$ *of proposition 2.5.17, relied on the Lefschetz trace formula, can be developed as follows:*

$$\begin{aligned}&\mathrm{Tr}(R_{G^{(2n)}(F_{\overline{\varpi}}\times F_\omega)}(\phi^{(2n)}_{G_R}\otimes\phi^{(2n)}_{G_L}))\\ &\qquad=\mathrm{Tr}(R_{P^{(2n)}(F_{\overline{\varpi}^1}\times F_{\omega^1})}(\phi^{(2n)}_{P_R}\otimes\phi^{(2n)}_{P_L}))\times\mathrm{Tr}(R_{G^{(2n)}(F^{nr}_{\overline{\varpi}}\times F^{nr}_\omega)}(\phi^{(2n)}_{G^{nr}_R}\otimes\phi^{(2n)}_{G^{nr}_L}))\end{aligned}$$

where

$$\begin{aligned}&\mathrm{Tr}(R_{G^{(2n)}(F^{nr}_{\overline{\varpi}}\times F^{nr}_\omega)}(\phi^{(2n)}_{G^{nr}_R}\otimes\phi^{(2n)}_{G^{nr}_L}))\\ &\qquad=\underset{j,m_j}{\oplus}\mathrm{vol}(G^{(2n)}(F^{nr}_{\overline{\varpi}}\times F^{nr}_\omega))\int_{G^{(2n)}}\phi^{(2n)}_{G^{nr}_{j_R}}(x^{nr}_{g_{j_R}})\otimes_D\phi^{(2n)}_{G^{nr}_{j_L}}(x^{nr}_{g_{j_L}})\ dx^{nr}_{g_{j_R}}\ dx^{nr}_{g_{j_L}}\end{aligned}$$

in such a way that $R_{P^{(2n)}(F_{\overline{\varpi}^1_{j,m_j}}\times F_{\omega^1_{j,m_j}})}(\phi^{(2n)}_{P_{j_R}}\otimes\phi^{(2n)}_{P_{j_L}})$ *is an invariant of* $\mathrm{Tr}(R_{G^{(2n)}(F_{\overline{\varpi}}\times F_\omega)}(\phi^{(2n)}_{G_R}\otimes\phi^{(2n)}_{G_L}))$.

Proof: The parabolic bifunctions $\phi^{(2n)}_{P_{j_R}}(x_{p_{j_R}})\otimes\phi^{(2n)}_{P_{j_L}}(x_{p_{j_L}})$, $1\le j\le r$, are in one-to-one correspondence and are defined on the bilinear parabolic semigroup $P^{(2n)}(F_{\overline{\varpi}^1}\times F_{\omega^1})$, being the isotropy subgroup of $G^{(2n)}(F_{\overline{\varpi}}\times F_\omega)$: so, the $R_{P^{(2n)}(F_{\overline{\varpi}^1_{j,m_j}}\times F_{\omega^1_{j,m_j}})}(\phi^{(2n)}_{P_{j_R}}\otimes\phi^{(2n)}_{P_{j_L}})$, $1\le j\le r$, are the invariants of $\mathrm{Tr}(R_{G^{(2n)}(F_{\overline{\varpi}}\times F_\omega)}(\phi^{(2n)}_{G_R}\otimes\phi^{(2n)}_{G_L}))$. ■

2.5.19 Remark

If $P^{(2n)}(F_{\overline{\varpi}^1}\times F_{\omega^1})$ is considered as the unitary representation of the bilinear semigroup $G^{(2n)}(F_{\overline{\varpi}}\times F_\omega)$, then the trace formula $\mathrm{Tr}(R_{G^{(2n)}(F_{\overline{\varpi}}\times F_\omega)}(\phi^{(2n)}_{G_R}\otimes\phi^{(2n)}_{G_L}))$ of proposition 2.5.17 refers to the Arthur's trace formula.
But, if $P^{(2n)}(F_{\overline{\varpi}^1}\times F_{\omega^1})$ is viewed as the isotropy subgroup of $G^{(2n)}(F_{\overline{\varpi}}\times F_\omega)$, then $\mathrm{tr}(R_{G^{(2n)}(F_{\overline{\varpi}}\times F_\omega)}(\phi^{(2n)}_{G_R}\otimes\phi^{(2n)}_{G_L}))$ relies on the Lefschetz trace formula.

2.5.20 Second trace formula for the operator $R(G^{(2n)}(F_{\overline{\varpi}}\times F_\omega))$

A trace formula of the operator $R(G^{(2n)}(F_{\overline{\varpi}}\times F_\omega))$ can be developed directly from the set of bifunctions $\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}})\otimes\phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}})\in L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\varpi}}\times F_\omega))$ on the conjugacy class representatives of $G^{(2n)}(F_{\overline{\varpi}}\times F_\omega)$ and not from an extension of the trace formula of the unitary parabolic operator $R(P^{(2n)}(F_{\overline{\varpi}^1}\times F_{\omega^1}))$.
As the bifunction $\phi^{(2n)}_{G_R}(x_{g_R})\otimes\phi^{(2n)}_{G_L}(x_{g_L})$, defined on the pseudo-ramified bilinear complete semigroup $G^{(2n)}(F_{\overline{\varpi}_\oplus}\times F_{\omega_\oplus})$, develops

$$\phi^{(2n)}_{G_R}(x_{g_R})\otimes\phi^{(2n)}_{G_L}(x_{g_L})=\underset{j,m_j}{\oplus}\left(\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}})\otimes\phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}})\right)$$

according to the bifunctions $\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}})\otimes\phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}})$ on the conjugacy class representatives $g^{(2n)}_{R\times L}[j,m_j]$ of $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$, which correspond to the cosets of $\mathrm{GL}_n(F_R\times F_L)\big/\mathrm{GL}_n((\mathbb{Z}\big/N\ \mathbb{Z})^2)$, we get clearly the searched trace formula:

$$\begin{aligned}&\operatorname{tr}(R_{\mathrm{GL}_n(F_{\overline{\omega}}\times F_\omega)\big/\mathrm{GL}_n((\mathbb{Z}\big/N\ \mathbb{Z})^2)}(\phi^{(2n)}_{G_R}(x_{g_R})\otimes\phi^{(2n)}_{G_L}(x_{g_L}))\\&\quad=\bigoplus_{j,m_j}\operatorname{vol}(G^{(2n)}(F_{\overline{\omega}}\times F_\omega))\int_{G^{(2n)}}\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}})\otimes_D\phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}})\ dx_{g_{j_R}}\ dx_{g_{j_L}}\ .\end{aligned}$$

This trace formula of $R(G^{(2n)}(F_{\overline{\omega}}\times F_\omega))$ corresponds to the trace formula obtained in proposition 2.5.15 from the trace of $R(P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1}))$. It results from the action of the Hecke bialgebra $\mathcal{H}_{R\times L}(n)$ on the pseudo-ramified bilinear semigroup $G^{(2n)}(F_{\overline{\omega}}\times F_\omega)$ since it is generated by the Hecke bioperators $(T_R(n;r)\otimes T_L(n;r))$ having as representation $\mathrm{GL}_n((\mathbb{Z}\big/N\ \mathbb{Z})^2)$ according to proposition 2.2.6.

This trace formula will be proved in chapter 3 to correspond to the Plancherel formula if the considered pseudo-ramified bilinear algebraic semigroup has undergone a projective toroidal isomorphism of compactification.

3 Langlands global correspondences for irreducible representations of GL(n)

Introductive keys

In chapter 2, nonabelian global class field concepts were introduced, i.e. essentially the construction of an algebraic bilinear semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_{\omega})$ constituting the $2n$-dimensional irreducible representation $\mathrm{Irr}\, W^{(2n)}_{F_{R\times L}}(W^{ab}_{F_R}\times W^{ab}_{F_L})$ of the product, right by left, of global Weil groups.
It was shown that $G^{(2n)}(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_{\omega})$ decomposes into diagonal and off-diagonal r conjugacy (bi)classes whose equivalent representatives m_r are noted $g^{(2n)}_{R\times L}[j,m_j]$, $1\le j\le r$. The equivalent representative m_j of the j-th left (resp. right) class of $T^{(2n)}(\widetilde{F}_{\omega})$ (resp. $T^{(2n)}(\widetilde{F}_{\overline{\omega}})^t\subset G^{(2n)}(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_{\omega})$ is noted $g^{(2n)}_{L}[j,m_j]$ (resp. $g^{(2n)}_{R}[j,m_j]$).
The algebraic representation space of the algebraic bilinear semigroup of matrices $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_{\omega})$ is noted $\mathrm{Repsp}(\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_{\omega}))$ and is precisely the algebraic bilinear semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_{\omega})$ isomorphic to the $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_{\omega})$-bisemimodule $(\widetilde{M}_R\otimes\widetilde{M}_L)$.
Then, the content of this chapter will concern:

a) **the introduction of a general bilinear cohomology** defined as a contravariant bifunctor H^* from smooth abstract (resp. algebraic) bisemivarieties together with a natural transformation $n_{H^*\to H^{[*,*]}}$ from H^* to the associated the de Rham bilinear cohomology $H^{[*,*]}$;

b) the Fulton-McPherson compactification of bilinear algebraic semigroups;

c) a toroidal compactification of the lattice bisemispace $X_{S_{R\times L}}$;

d) a double coset decomposition of $\mathrm{GL}_n(F_{\overline{\omega}}\times F_{\omega})$ leading to the equivalent of a bilinear Shimura variety $\overline{S}^{P_n}_{K_n}$;

e) the Langlands global correspondence on $\mathrm{Repsp}(\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}_{\oplus}}\times\widetilde{F}_{\omega_{\oplus}}))$ based upon the study of the ring of regular (bi)functions on $\mathrm{Repsp}(\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_{\omega}))$ and the corresponding Langlands “real” correspondence.

3.1 Compactifications of bilinear algebraic semigroups and global holomorphic correspondences

3.1.1 The compactification of bilinear algebraic semigroups

The compactification of the bilinear algebraic semigroup $G^{(n)}(\widetilde F^+_{\overline v} \times \widetilde F^+_v)$ over the product, right by left, of real pseudo-ramified extension semirings leads to the compactified bilinear algebraic semigroup $G^{(n)}(F^+_{\overline v} \times F^+_v)$, taking into account the inclusions

$$\begin{tikzcd}
G^{(n)}(F^+_{\overline v} \times F^+_v) \arrow[r, hook] \arrow[d, "\wr"] & G^{(2n)}(F_{\overline\omega} \times F_\omega) \arrow[d, "\wr"] \\
G^{(n)}(\widetilde F^+_{\overline v} \times \widetilde F^+_v) \arrow[r, hook] & G^{(2n)}(\widetilde F_{\overline\omega} \times \widetilde F_\omega)
\end{tikzcd}$$

of the "real" bilinear algebraic semigroups into the corresponding "complex" bilinear algebraic semigroups as justified in section 3.5.1.

We show that this problem is closely associated to the generation of a smooth bilinear general bisemivariety $\tau_\sigma^{(n)}(F^+_{\overline v} \times F^+_v)$ characterized by polyhedral convex bicones $\sigma^{(n)}_{R\times L}(F^+_{\overline v_{j_\delta,m_{j_\delta}}} \times F^+_{v_{j_\delta,m_{j_\delta}}})$.

And, this bilinear general bisemivariety $\tau_\sigma^{(n)}(F^+_{\overline v} \times F^+_v)$ is in one-to-one correspondence with the bilinear general "toroidal" bisemivariety $\tau_T^{(n)}(F^{+,T}_{\overline v} \times F^{+,T}_v)$ in such a way that the polyhedral convex bicones $\sigma^{(n)}_{R\times L}(F^+_{\overline v_{j_\delta,m_{j_\delta}}} \times F^+_{v_{j_\delta,m_{j_\delta}}})$ are transformed into products, right by left, of n-dimensional real semitori $T^n_R[j_\delta, m_{j\delta}] \times T^n_L[j_\delta, m_{j\delta}] \in G^{(n)}(F^{+,T}_{\overline v} \times F^{+,T}_v)$.

3.1.2 An adaptation of the Fulton-McPherson compactification

The set of left (resp. right) real pseudo-ramified completions

$$F^+_v = \{F^+_{v_{1_\delta}}, \cdots, F^+_{v_{j_\delta,m_{j_\delta}}}, \cdots, F^+_{v_{r_\delta}}\} \qquad (\text{resp.} \quad F^+_{\overline v} = \{F^+_{\overline v_{1_\delta}}, \cdots, F^+_{\overline v_{j_\delta,m_{j_\delta}}}, \cdots, F^+_{\overline v_{r_\delta}}\} \ )$$

introduced in section 1.1.5, proceeds from the set of left (resp. right) real pseudo-ramified Galois extensions of k :

$$\widetilde F^+_v = \{\widetilde F^+_{v_{1_\delta}}, \cdots, \widetilde F^+_{v_{j_\delta,m_{j_\delta}}}, \cdots, \widetilde F^+_{v_{r_\delta}}\} \qquad (\text{resp.} \quad \widetilde F^+_{\overline v} = \{\widetilde F^+_{\overline v_{1_\delta}}, \cdots, \widetilde F^+_{\overline v_{j_\delta,m_{j_\delta}}}, \cdots, \widetilde F^+_{\overline v_{r_\delta}}\} \ )$$

(see sections 1.1.4 and 1.1.7).

Let $G^{(n)}(\widetilde F^+_{\overline v} \times \widetilde F^+_v)$ denote the non-compact real bilinear algebraic semigroup with entries in $(\widetilde F^+_{\overline v} \times \widetilde F^+_v)$.

Several types of compactification can be envisaged for it: we shall introduce here an adaptation of the compactification of W. Fulton and R. McPherson [F-M], which allows to consider a compactification by unit blocks consisting in irreducible real extensions $\widetilde F_{v^{j'_\delta}_{j_\delta}}$, $1\le j'_\delta\le j_\delta$, of rank N in one-to-one correspondence with the irreducible real completions $F_{v^{j'_\delta}_{j_\delta}}$, as defined in section 1.1.5.

Let $\{\tilde g^{(n)}_{R\times L}[j_\delta,m_{j_\delta}]\}^t_{j_\delta=1}$ denote the set of conjugacy class representatives of the real non compact bilinear algebraic semigroup $G^{(n)}(\widetilde F^+_{\overline v}\times\widetilde F^+_v)$ and let

$$\begin{aligned}\{\tilde g^{(n)}_L[j_\delta,m_{j_\delta}]\}^t_{j_\delta=1}&\subset G^{(n)}(\widetilde{F^+_v})\simeq T_n(\widetilde F^+_v)\\ \text{(resp.}\quad \{\tilde g^{(n)}_R[j_\delta,m_{j_\delta}]\}^t_{j_\delta=1}&\subset G^{(n)}(\widetilde{F^+_{\overline v}})\simeq T^t_n(\widetilde F^+_{\overline v})\ )\end{aligned}$$

be the corresponding set of left (resp. right) conjugacy class representatives of the real non compact linear algebraic semigroup $G^{(n)}(\widetilde{F^+_v})$ (resp. $G^{(n)}(\widetilde{F^+_{\overline v}})$).

Referring to proposition 2.3.4, it is easy to see that the rank $r_{v_{j_\delta}}$ (resp. $r_{\overline v_{j_\delta}}$)of the conjugacy class representative $\tilde g^{(n)}_L[j_\delta,m_{j_\delta}]$ (resp. $\tilde g^{(n)}_R[j_\delta,m_{j_\delta}]$) is given by:

$$r_{v_{j_\delta}}=N^n\cdot j^n\qquad\text{(resp.}\quad r_{\overline v_{j_\delta}}=N^n\cdot j^n\ )$$

where:

- N is the rank of the real irreducible extensions $\widetilde F_{v^{j'_\delta}_{j_\delta}}$,
- j is the global class residue degree of $\widetilde F_{v_{j_\delta}}$.

If n_F is the number of non-units of $\widetilde F_{v^{j'_\delta}_{j_\delta}}$, then:

- the number of points of $\tilde g^{(n)}_L[j_\delta,m_{j_\delta}]$ (resp. $\tilde g^{(n)}_R[j_\delta,m_{j_\delta}]$) is:

$$\begin{aligned}n_{\tilde g^{(n)}_L[j_\delta,m_{j_\delta}]}&=j^n\cdot N^n\cdot n^n_F\\ \text{(resp.}\quad n_{\tilde g^{(n)}_R[j_\delta,m_{j_\delta}]}&=j^n\cdot N^n\cdot n^n_F\ ),\end{aligned}$$

- the number of irreducible subsets $\widetilde F_{v^{j'_\delta}_{j_\delta}}(n)$ of $\tilde g^{(n)}_L[j_\delta,m_{j_\delta}]$ (resp. $\tilde g^{(n)}_R[j_\delta,m_{j_\delta}]$) is:

$$n_{\widetilde F_{v^{j'_\delta}_{j_\delta}}}=j^n\qquad\text{(resp.}\quad n_{\widetilde F_{\overline v_{j_\delta}}}=j^n\ ).$$

The compactification of the conjugacy class representatives $\tilde{g}_L^{(n)}[j_\delta, m_{j_\delta}]$ and $\tilde{g}_R^{(n)}[j_\delta, m_{j_\delta}]$ is realized symmetrically respectively in the upper and in the lower half spaces according to the method developed by W. Fulton and R. McPherson in [F-M]: this leads to the following propositions.

3.1.3 Proposition

At each irreducible subset $\widetilde{F}_{v^{j'_\delta}_{j_\delta}}(n) \in \tilde{g}_L^{(n)}[j_\delta, m_{j_\delta}]$ *(resp.* $\widetilde{F}_{\overline{v}^{j'_\delta}_{j_\delta}}(n) \in \tilde{g}_R^{(n)}[j_\delta, m_{j_\delta}]$ *) at* $N \cdot n_F$ *elements, corresponds a non-singular irreducible compact completion* $F_{v^{j'_\delta}_{j_\delta}}(n)$ *(resp.* $F_{\overline{v}^{j'_\delta}_{j_\delta}}(n)$ *), i.e. a closed irreducible compact one-dimensional subset of* $g_L^{(n)}[j_\delta, m_{j_\delta}] \in G^{(n)}(F_v^+)$ *(resp.* $g_R^{(n)}[j_\delta, m_{j_\delta}] \in G^{(n)}(F_{\overline{v}}^+)$ *) such that:*

- *a) the union of the irreducible completions* $F_{v^{j'_\delta}_{j_\delta}}(n)$ *(resp.* $F_{\overline{v}^{j'_\delta}_{j_\delta}}(n)$ *) is* $g_L^{(n)}[j_\delta, m_{j_\delta}]$ *(resp.* $g_R^{(n)}[j_\delta, m_{j_\delta}]$ *);*
- *b) these irreducible completions meet transversally.*

Proof:

1. As the conjugacy class representatives $\widetilde{g}_L^{(n)}[j_\delta, m_{j_\delta}]$ (resp. $\widetilde{g}_R^{(n)}[j_\delta, m_{j_\delta}]$) are built from irreducible subsets $\widetilde{F}_{v^{j'_\delta}_{j_\delta}}(n)$ (resp. $\widetilde{F}_{\overline{v}^{j'_\delta}_{j_\delta}}(n)$), it is normal that the compactification acts on these unit blocks.

2. A set of blowups is then envisaged on the $N \cdot n_F$ elements of the irreducible subsets $\widetilde{F}_{v^{j'_\delta}_{j_\delta}}(n)$ and $\widetilde{F}_{\overline{v}^{j'_\delta}_{j_\delta}}(n)$in such a way that the $N \cdot n_F$ points of $\widetilde{F}_{v^{j'_\delta}_{j_\delta}}(n)$ form an irreducible divisor (or completion) $F_{v^{j'_\delta}_{j_\delta}}(n)$ cent red on a point in the upper half space and that the $N \cdot n_F$ points of $\widetilde{F}_{\overline{v}^{j'_\delta}_{j_\delta}}(n)$ form a symmetric divisor $D(F_{\overline{v}^{j'_\delta}_{j_\delta}}(n))$ in the lower half space centered on a symmetric point.

   A sequence of maximum $N \cdot n_F$ blowups [F-M] is envisaged such that the $N \cdot n_F$ points become together a point and generate a completion (i.e. a closed irreducible compact one-dimensional subset) which constitutes a limiting configuration in the compactification of $\widetilde{F}_{v^{j'_\delta}_{j_\delta}}(n)$ (resp. $\widetilde{F}_{\overline{v}^{j'_\delta}_{j_\delta}}(n)$), in the sense that equivalent compactifications of $\widetilde{F}_{v^{j'_\delta}_{j_\delta}}(n)$ (resp. $\widetilde{F}_{\overline{v}^{j'_\delta}_{j_\delta}}(n)$) exist (they are called degenerated in [F-M]).

3. More concretely:

- a first irreducible subset $\widetilde{F}_{v^{1'_\delta}_{j_\delta}}(n)$ is compactified by blowups of its $N \cdot n_F$ points into the completion $F_{v^{1'_\delta}_{j_\delta}}(n)$;
- a second irreducible subset $\widetilde{F}_{v^{2'_\delta}_{j_\delta}}(n)$ is compactified in the completion $F_{v^{2'_\delta}_{j_\delta}}(n)$ by blowing up its $(N \cdot n_F)$ points in a compact neighbourhood of $F_{v^{1'_\delta}_{j_\delta}}(n)$ in such a way that $F_{v^{2'_\delta}_{j_\delta}}(n)$ be transversal to $F_{v^{1'_\delta}_{j_\delta}}(n)$;
- it is proceeded in that manner for all irreducible subsets of $\tilde{g}_L^{(n)}[j_\delta, m_{j_\delta}]$ in such a way that the compactification of these gives a polyhedral convex cone $g_L^{(n)}[j_\delta, m_{j_\delta}]$ with j_δ^n completions or closed one-dimensional compact subsets meeting transversally (or having normal crossings);
- to the polyhedral convex cone $g_L^{(n)}[j_\delta, m_{j_\delta}]$ localized in the upper half space corresponds a symmetric polyhedral convex cone $g_R^{(n)}[j_\delta, m_{j_\delta}]$ localized in the lower half space. ■

3.1.4 Definitions

- A **left (resp. right) polyhedral convex cone** $\sigma_L^{(n)}(F^+_{v_{j_\delta},m_{j_\delta}})$ (resp. $\sigma_R^{(n)}(F^+_{\overline{v}_{j_\delta},m_{j_\delta}})$) is a conjugacy class representative $g_L^{(n)}[j_\delta, m_{j_\delta}]$ (resp. $g_R^{(n)}[j_\delta, m_{j_\delta}]$) of the linear complete algebraic semigroup $G^{(n)}(F_v^+)$ (resp. $G^{(n)}(F_{\overline{v}}^+)$) on the $(\mathbb{Z}/N\,\mathbb{Z})$-lattice Λ_v (resp. $\Lambda_{\overline{v}}$) (corresponding to the $(\mathbb{Z}/N\,\mathbb{Z})$-lattice Λ_ω (resp. $\Lambda_{\overline{\omega}}$) introduced in section 2.2 in such a way that $g_L^{(n)}[j_\delta, m_{j_\delta}]$ (resp. $g_R^{(n)}[j_\delta, m_{j_\delta}]$) be included into the subsemigroup $g_L^{(2n)}[j, m_j]$ (resp. $g_R^{(2n)}[j, m_j]$) localized in the upper (resp. lower) half space) [Dan].

- A **polyhedral convex bicone** $\sigma_{R\times L}^{(n)}(F^+_{\overline{v}_{j_\delta},m_{j_\delta}} \times F^+_{v_{j_\delta},m_{j_\delta}})$ is a conjugacy class representative $g_{R\times L}^{(n)}[j_\delta, m_{j_\delta}]$ of the bilinear complete semigroup $G^{(n)}(F_{\overline{v}}^+ \times F_v^+)$ on the $(\mathbb{Z}/N\,\mathbb{Z})^2$-bilattice $\Lambda_{\overline{v}} \otimes \Lambda_v$.

- A **smooth linear general semivariety** [K-K-M-S] $\tau_\sigma^{(n)}(F_v^+)$ (resp. $\tau_\sigma^{(n)}(F_{\overline{v}}^+)$) is a linear complete algebraic semigroup $G^{(n)}(F_v^+)$ (resp. $G^{(n)}(F_{\overline{v}}^+)$) composed of the family $\{g_L^{(n)}[j_\delta, m_{j_\delta}]\}$ (resp. $\{g_R^{(n)}[j_\delta, m_{j_\delta}]\}$) of disjoint conjugacy class representatives together with a collection of charts from real conjugacy class representatives $g_L^{(n)}[j_\delta, m_{j_\delta}]$ (resp. $g_R^{(n)}[j_\delta, m_{j_\delta}]$) to their complex equivalents $g_L^{(2n)}[j, m_j]$

(resp. $g_R^{(2n)}[j,m_j]$):

$$\begin{array}{rrcl} & cz^j : \ \{g_L^{(n)}[j_\delta,m_{j_\delta}]\}_{m_{j_\delta}} & \longrightarrow & g_L^{(2n)}[j,m_j] \\ (\text{resp.} & cz^{*j} : \ \{g_R^{(n)}[j_\delta,m_{j_\delta}]\}_{m_{j_\delta}} & \longrightarrow & g_R^{(2n)}[j,m_j] \ ) \end{array}$$

where cz^j (resp. cz^{*j}) are coordinate functions on the corresponding conjugacy class representatives.

The cz^j (resp. cz^{*j}) can be represented as Laurent monomials in functions of the complex variables $z_1^j,\cdots,z_n^j$ (resp. $z_1^{*j},\cdots,z_n^{*j}$).

- A **compactified smooth linear general complex semivariety** $\tau_{c_\sigma}^{(2n)}(F_\omega)$ (resp. $\tau_{c_\sigma}^{(2n)}(F_{\overline{\omega}})$) is obtained from $\tau_\sigma^{(n)}(F_v^+)$ (resp. $\tau_\sigma^{(n)}(F_{\overline{v}}^+)$) by gluing together the conjugacy class representatives of $G^{(2n)}(F_\omega)$ (resp. $G^{(2n)}(F_{\overline{\omega}})$) in such a way that, if

  $$\begin{array}{rrcl} & f_{\omega_{j,m_j}}(z^j) : \ g_L^{(2n)}[j,m_j] & \longrightarrow & \mathbb{C} \\ (\text{resp.} & f_{\overline{\omega}_{j,m_j}}(z^{*j}) : \ g_R^{(2n)}[j,m_j] & \longrightarrow & \mathbb{C}^* \ ) \end{array}$$

  is a function on $g_L^{(2n)}[j,m_j]$ (resp. $g_R^{(2n)}[j,m_j]$),
  then a Laurent polynomial

  $$f_\omega(z) : \quad G^{(2n)}(F_\omega) \longrightarrow \mathbb{C}$$

  $$(\text{resp.} \quad f_{\overline{\omega}}(z^*) : \quad G^{(2n)}(F_{\overline{\omega}}) \longrightarrow \mathbb{C}^* \ )$$

  given by:

  $$\begin{array}{rl} & f_\omega(z) = \sum\limits_{j,m_j=1}^{r} f_{\omega_{j,m_j}}(z^j) \ , \quad 1 \le j \le r \le \infty \ , \\ (\text{resp.} & f_{\overline{\omega}}(z^*) = \sum\limits_{j,m_j=1}^{r} f_{\overline{\omega}_{j,m_j}}(z^{*j}) \ ) \end{array}$$

  can be introduced on $\tau_{c_\sigma}^{(2n)}(F_{\omega_\oplus})$ (resp. $\tau_{c_\sigma}^{(2n)}(F_{\overline{\omega}_\oplus})$) as a(n) (in)finite linear combination of Laurent monomials corresponding to the left (resp. right) polyhedral convex cones $\sigma_L^{(n)}(F^+_{v_{j_\delta,m_{j_\delta}}})$ (resp. $\sigma_R^{(n)}(F^+_{\overline{v}_{j_\delta,m_{j_\delta}}})$).

3.1.5 Proposition

Let $\tau_{c_\sigma}^{(2n)}(F_\omega)$ (resp. $\tau_{c_\sigma}^{(2n)}(F_{\overline{\omega}})$) be a compactified smooth linear general semivariety whose conjugacy class representatives $g_L^{(2n)}[j,m_j]$ (resp. $g_R^{(2n)}[j,m_j]$) are glued together and on which the functions $f_\omega(z^j)$ (resp. $f_{\overline{\omega}}(z^{*j})$) are considered.

Then, on the compactified semivariety $\tau^{(2n)}_{c_\sigma}(F_\omega)$ *(resp.* $\tau^{(2n)}_{c_\sigma}(F_{\overline{\omega}})$ *), the (differentiable) function* $f_\omega(z)$ *(resp.* $f_{\overline{\omega}}(z^*)$ *), defined in a neighbourhood of a point* z_0 *(resp.* z_0^* *) of* $\mathbb{C}^n$ *, is holomorphic at* z_0 *(resp.* z_0^* *) if we have the following multiple power series development:*

$$f_\omega(z)=\sum_{j=1}^{\infty}\sum_{m_j}c_{j,m_j}\ (z'_1-z_{0_1})^j\cdots(z'_n-z_{0_n})^j\ ,$$

$$\textit{(resp.}\quad f_{\overline{\omega}}(z^*)=\sum_{j=1}^{\infty}\sum_{m_j}c^*_{j,m_j}\ (z'^*_1-z'^*_{0_1})^j\cdots(z'^*_n-z'^*_{0_n})^j\ )$$

where $z'_1,\cdots,z'_n$ *are (functions of) complex variables, on unitary closed supports taking into account that:*

a) to each conjugacy class representative $g^{(2n)}_L[j,m_j]\in\tau^{(2n)}_{c_\sigma}(F_\omega)$ *(resp.* $g^{(2n)}_R[j,m_j]\in\tau^{(2n)}_{c_\sigma}(F_{\overline{\omega}})$ *) corresponds a term of the multiple power series development of* $f_\omega(z)$ *(resp.* $f_{\overline{\omega}}(z^*)$ *). These terms have the same structure by construction of* $g^{(2n)}_L[j,m_j]$ *(resp.* $g^{(2n)}_R[j,m_j]$ *),* $1\le j\le r\le\infty$ *.*

b) the coefficients c_{j,m_j} *(resp.* c^*_{j,m_j} *) have an inflation action from functions on the conjugacy class representatives* $P^{(2n)}(F_{\omega^1_j})$ *(resp.* $P^{(2n)}(F_{\overline{\omega}^1_j})$ *) of the "complex" parabolic subsemigroup to the corresponding functions on the conjugacy class representatives* $g^{(2n)}_L[j,m_j]$ *(resp.* $g^{(2n)}_R[j,m_j]$ *) of the algebraic semigroup* $G^{(2n)}(F_\omega)$ *(resp.* $G^{(2n)}(F_{\overline{\omega}})$ *).*

c) $j\to\infty$ *.*

Proof:

a) According to definitions 3.1.4, the coordinate functions cz^j (resp. cz^{*j}) on the conjugacy classes of $\tau^{(2n)}_{c_\sigma}(F_\omega)$ (resp. $\tau^{(2n)}_{c_\sigma}(F_{\overline{\omega}})$) can be represented as Laurent monomials. So, each term of the multiple power series development of $f_\omega(z)$ (resp. $f_{\overline{\omega}}(z^*)$) corresponds to a conjugacy class representative $g^{(2n)}_L[j,m_j]$ (resp. $g^{(2n)}_R[j,m_j]$).

b) If $j\to\infty$, the number of conjugacy classes of $\tau^{(2n)}_{c_\sigma}(F_\omega)$ (resp. $\tau^{(2n)}_{c_\sigma}(F_{\overline{\omega}})$) must tend to infinity in order that the multiple power series, converging to z in some neighbourhood of z_0 , be equal there to $f_\omega(z)$ (resp. $f_{\overline{\omega}}(z^*)$). ■

3.1.6 Proposition

The coefficients c_{j,m_j} of the holomorphic function $f_\omega(z)$ (resp. $f_{\overline{\omega}}(z^*)$) on the compactified algebraic semivariety $\tau^{(2n)}_{c_\sigma}(F_\omega)$ (resp. $\tau^{(2n)}_{c_\sigma}(F_{\overline{\omega}})$) are given by the products:

$$\lambda^{\frac{1}{2}}(2n,j,m_j)=\prod_{d=1}^{2n}\lambda_d^{\frac{1}{2}}(2n,j,m_j)$$

of the square roots of the eigenvalues $\lambda_d(2n,j,m_j)$ of the (j,m_j)-th coset representatives $U_{j,m_{j_R}}\times U_{j,m_{j_L}}$ of the product $T_R(n;t)\otimes T_L(n;t)$ of the Hecke operators.

Proof: As the holomorphic function $f_\omega(z)$ (resp. $f_{\overline{\omega}}(z^*)$) decomposes into a multiple power series whose terms are (sub)functions on the conjugacy class representatives $g^{(2n)}_L[j,m_j]$ (resp. $g^{(2n)}_R[j,m_j]$) of the semivariety $\tau^{(2n)}_{c_\sigma}(F_\omega)$ (resp. $\tau^{(2n)}_{c_\sigma}(F_{\overline{\omega}})$), $f_\omega(z)$ (resp. $f_{\overline{\omega}}(z^*)$) corresponds to an endomorphism of $\tau^{(2n)}_{c_\sigma}(F_\omega)$ (resp. $\tau^{(2n)}_{c_\sigma}(F^+_{\overline{\omega}})$) into itself.
On the other hand, as the semivariety $\tau^{(2n)}_{c_\sigma}(F_\omega)$ (resp. $\tau^{(2n)}_{c_\sigma}(F_{\overline{\omega}})$) is defined on the $(\mathbb{Z}/N\,\mathbb{Z})$-lattice Λ_ω (resp. $\Lambda_{\overline{\omega}}$) (see definitions 3.1.4), on which acts the Hecke operator $T_L(n;t)$ (resp. $T_R(n;t)$) whose product $T_R(n;t)\otimes T_L(n;t)$ has a representation in the subgroup of matrices $\mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)$, the product of the square roots of the eigenvalues $\lambda(2n,j,m_j)$ of the (j,m_j)-th coset representative $U_{j,m_{j_R}}\times U_{j,m_{j_L}}$ of $T_R(n;t)\otimes T_L(n;t)$ will naturally constitute the searched coefficient c_{j,m_j} of the holomorphic function $f_\omega(z)$: this results from proposition 2.2.5. ■

3.1.7 Laurent polynomial on the general bisemivariety

Let $G^{(n)}(\widetilde{F}^+_{\overline{v}}\times\widetilde{F}^+_{v})$ denote the non-compact real bilinear algebraic semigroup and let $G^{(n)}(F^+_{\overline{v}}\times F^+_{\overline{v}})$ be its locally compact equivalent obtained by the Fulton-McPherson compactification as introduced in proposition 3.1.4. To the set of polyhedral convex bicones $\sigma^{(n)}_{R\times L}(F^+_{\overline{v}_{j_\delta,m_{j_\delta}}}\times F^+_{v_{j_\delta,m_{j_\delta}}})\equiv g^{(n)}_{R\times L}[j_\delta,m_{j_\delta}]$ of $G^{(n)}(F^+_{\overline{v}}\times F^+_{v})$ corresponds the smooth general bisemivariety $\tau^{(n)}_{\sigma}(F^+_{\overline{v}}\times F^+_{v})$ which leads to its compactified complex equivalent $\tau^{(2n)}_{c_\sigma}(F_{\overline{\omega}}\times F_\omega)$ obtained by gluing together the conjucacy class representatives of $G^{(n)}(F^+_{\overline{v}}\times F^+_{v})$. And, a Laurent (bi)polynomial

$$f_{\overline{\omega}}(z^*)\otimes_D f_\omega(z)=\sum_{j=1}^{r}\sum_{m_j}c_{j,m_j}\cdot c^*_{j,m_j}\;z^{*j}\cdot z^j$$

can be introduced on $\tau^{(2n)}_{c_\sigma}(F_{\overline{\omega}_\oplus}\times F_{\omega_\oplus})$.

3.1.8 Proposition

On the compactified complex bisemivariety $\tau^{(2n)}_{c_\sigma}(F_{\overline{\omega}_\oplus}\times F_{\omega_\oplus})$, the bifunction $f_{\overline{\omega}}(z^*)\otimes f_\omega(z)$, defined in the neighbourhood of a bipoint $(z_0^*\times z_0)$ of $\mathbb{C}^n\times\mathbb{C}^n$, is holomorphic at $(z_0^*\times z_0)$

if we have the multiple power series development:

$$f_{\overline{\omega}}(z^*) \otimes_{(D)} f_\omega(z) = \sum_{j=1}^{\infty} \sum_{m_j} c^*_{j,m_j}\ c_{j,m_j}\ (z_1^{*'} z'_1 - z^*_{0_1} z_{0_1})^j \cdots (z_n^{*'} z'_n - z^*_{0_n} z_{0_n})^j$$

verifying:

a) to each conjugacy class representative $g^{(2n)}_{R\times L}[j,m_j] \in \tau^{(2n)}_{c_\sigma}(F_{\overline{\omega}} \times F_\omega)$ corresponds a term of $f_{\overline{\omega}}(z^*) \otimes f_\omega(z)$;

b) $j \to \infty$;

c) each term of $f_{\overline{\omega}}(z^*) \otimes_{(D)} f_\omega(z)$ generates a function subspace of dimension j^{2n} .

Proof: This is an adaptation of proposition 3.1.6 to the bilinear case. ■

3.1.9 Corollary

The holomorphic bifunction $f_{\overline{\omega}}(z^*) \otimes_{(D)} f_\omega(z)$ on the compactified algebraic bisemivariety $\tau^{(2n)}_{c_\sigma}(F_{\overline{\omega}_\oplus} \times F_{\omega_\oplus})$ constitutes an irreducible holomorphic representation $\mathrm{Irr\,hol}(\mathrm{GL}_n(F_{\overline{\omega}} \times F_\omega))$ of the bilinear complete semigroup $G^{(2n)}(F_{\overline{\omega}} \times F_\omega) \approx \mathrm{GL}_n(F_{\overline{\omega}} \times F_\omega)$.

3.1.10 Power series development on the real compactified semigroups

If the inclusion $G^{(n)}(F^+_{\overline{v}} \times F^+_v \hookrightarrow G^{(2n)}(F_{\overline{\omega}} \times F_\omega)$ of the real compactified bilinear complete semigroup $G^{(n)}(F^+_{\overline{v}} \times F^+_v)$ into its complex equivalent $G^{(2n)}(F_{\overline{\omega}} \times F_\omega)$ is not taken into account, a power series development can be envisaged on $G^{(n)}(F^+_v) = \{g^{(n)}_L[j_\delta, m_{j_\delta}]\}$ (resp. $G^{(n)}(F^+_{\overline{v}}) = \{g^{(n)}_R[j_\delta, m_{j_\delta}]\}$).
Consider the compactified smooth linear general real semivariety $\tau^{(n)}_{c_\sigma}(F^+_v)$ (resp. $\tau^{(n)}_{c_\sigma}(F^+_{\overline{v}})$) obtained from $\tau^{(n)}_\sigma(F^+_v)$ (resp. $\tau^{(n)}_\sigma(F^+_{\overline{v}})$) by gluing together the conjugacy class representatives of $G^{(n)}(F^+_v)$ (resp. $G^{(n)}(F^+_{\overline{v}})$) in such a way that:

$$\begin{aligned} f_{v_{j_\delta,m_{j_\delta}}}(x^{j_\delta}) : \quad & g^{(n)}_L[j_\delta, m_{j_\delta}] \longrightarrow I\!R\ , \qquad x = (x_1,\dots,x_n) \in I\!R^n\ , \\ (\text{resp.}\quad f_{\overline{v}_{j_\delta,m_{j_\delta}}}(-x^{j_\delta}) : \quad & g^{(n)}_R[j_\delta, m_{j_\delta}] \longrightarrow I\!R\ ) \end{aligned}$$

be a function on $g^{(n)}_L[j_\delta, m_{j_\delta}]$ (resp. $g^{(n)}_R[j_\delta, m_{j_\delta}]$).

3.1.11 Proposition

On the compactified real semivariety $\tau^{(n)}_{c_\sigma}(F^+_v)$ *(resp.* $\tau^{(n)}_{c_\sigma}(F^+_{\overline{v}})$ *), the (differentiable) function* $f_v(x)$ *(resp.* $f_{\overline{v}}(x)$*) has a power series development if it can be expressed according to*

$$\begin{aligned} f_v(x) &= \sum_{j_\delta} \sum_{m_{j_\delta}} c_{j_\delta,m_{j_\delta}} (x')^{j_\delta} \\ \text{(resp.} \quad f_{\overline{v}}(-x) &= \sum_{j_\delta} \sum_{m_{j_\delta}} \overline{c}_{j_\delta,m_{j_\delta}} (-x')^{j_\delta} \ ), \end{aligned}$$

where x' *is a function of the real variables* $x = (x_1, \dots, x_n)$*.*
This implies that:

$$\begin{aligned} f_{v_{j_\delta,m_{j_\delta}}}(x^{j_\delta}) &= c_{j_\delta,m_{j_\delta}} (x')^{j_\delta} \\ \text{(resp.} \quad f_{\overline{v}_{j_\delta,m_{j_\delta}}}(-x^{j_\delta}) &= \overline{c}_{j_\delta,m_{j_\delta}} (-x')^{j_\delta} \ ). \end{aligned}$$

Proof: The power series development of $f_v(x)$ (resp. $f_{\overline{v}}(-x)$) on the complete algebraic semigroup $G^{(n)}(F^+_v)$ (resp. $G^{(n)}(F^+_{\overline{v}})$) is rather "natural" since its conjugacy class representatives have the same structure by construction. So, the terms of $f_v(x)$ (resp. $f_{\overline{v}}(-x)$) on the conjugacy class representatives have the same functional structure. ■

3.1.12 Corollary

The power series bifunction $f_{\overline{v}}(-x) \otimes_{(D)} f_v(x)$ *on the compactified algebraic real bisemivariety* $\tau^{(n)}_{c_\sigma}(F^+_{\overline{v}_\oplus} \times F^+_{v_\oplus})$ *constitutes an irreducible functional representation space* $\operatorname{Irr}\operatorname{Repsp}(\mathrm{GL}_n(F^+_{\overline{v}} \times F^+_v))$ *of the bilinear complete semigroup* $G^{(n)}(F^+_{\overline{v}} \times F^+_v)$*.*

3.1.13 Proposition

Let $\widetilde{F}^+_{v_\oplus} = \bigoplus\limits_{j_\delta} \widetilde{F}^+_{v_{j_\delta}} \bigoplus\limits_{m_{j_\delta}} \widetilde{F}^+_{v_{j_\delta,m_{j_\delta}}}$ *(resp.* $\widetilde{F}^+_{\overline{v}_\oplus} = \bigoplus\limits_{j_\delta} \widetilde{F}^+_{\overline{v}_{j_\delta}} \bigoplus\limits_{m_{j_\delta}} \widetilde{F}^+_{\overline{v}_{j_\delta,m_{j_\delta}}}$ *) denote the sum of the left (resp. right) real pseudo-ramified extensions of* k *. Then, the non compact real bilinear algebraic semigroup* $G^{(n)}(\widetilde{F}^+_{\overline{v}_\oplus} \times \widetilde{F}^+_{v_\oplus})$ *, associated with* $G^{(n)}(F^+_{\overline{v}} \times F^+_v)$ *, constitutes a* n*-dimensional non compact irreducible representation of the product* $W^{ab}_{F^+_R} \times W^{ab}_{F^+_L}$ *of global Weil groups according to:*

$$\operatorname{Irr}\operatorname{Rep} W^{(n)}_{F^+_R \times F^+_L} : \quad W^{ab}_{F^+_R} \times W^{ab}_{F^+_L} \longrightarrow G^{(n)}(\widetilde{F}^+_{\overline{v}_\oplus} \times \widetilde{F}^+_{v_\oplus}) \ .$$

Proof: This is an adaptation of proposition 2.4.7 to the real case. ■

3.1.14 Proposition

On the non compact real bilinear algebraic semigroup $G^{(n)}(\widetilde F^+_{\overline{v}_\oplus}\times \widetilde F^+_{v_\oplus})$ *there exists the global functional correspondence:*

$$\begin{tikzcd}
\operatorname{Irr}\operatorname{Rep} W^{(n)}_{F^+_R\times F^+_L}(W^{ab}_{F^+_R}\times W^{ab}_{F^+_L}) \arrow[rr] \arrow[d, equal] & & \operatorname{Irr}\operatorname{Repsp}(G^{(n)}(F^+_{\overline{v}_\oplus}\times F^+_{v_\oplus})) \arrow[d, equal] \\
G^{(n)}(\widetilde F^+_{\overline{v}_\oplus}\times \widetilde F^+_{v_\oplus}) \arrow[dr] & & f_{\overline{v}}(-x)\otimes f_v(x) \\
& G^{(n)}(F^+_{\overline{v}_\oplus}\times F^+_{v_\oplus}) \arrow[ur] &
\end{tikzcd}$$

- *from the sum of the products, right by left, of the equivalence classes of the irreducible n-dimensional non-compact representation* $\operatorname{Irr}\operatorname{Rep} W^{(n)}_{F^+_R\times F^+_L}(W^{ab}_{F^+_R}\times W^{ab}_{F^+_L})$ *of the product* $W^{ab}_{F^+_R}\times W^{ab}_{F^+_L}$ *of global Weil groups, given by the non-compact real bilinear algebraic semigroup* $G^{(n)}(\widetilde F^+_{\overline{v}_\oplus}\times \widetilde F^+_{v_\oplus})$
- *to the sum of the products, right by left, of the equivalence classes of the irreducible representation space* $\operatorname{Irr}\operatorname{Repsp}(G^{(n)}(F^+_{\overline{v}}\times F^+_v))$ *of* $G^{(n)}(F^+_{\overline{v}}\times F^+_v)$ *given by the power series bifunction* $f_{\overline{v}}(-x)\otimes f_v(x)$ *on the compactified algebraic bisemivariety* $\tau^{(n)}_{c_\sigma}(F^+_{\overline{v}_\oplus}\times F^+_{v_\oplus})$ *or on the compact bilinear algebraic semigroup* $G^{(n)}(F^+_{\overline{v}}\times F^+_v)$ *.*

Proof:

1. It has been seen in proposition 3.1.13 that $\operatorname{Irr}\operatorname{Rep} W^{(n)}_{F^+_R\times F^+_L}(W^{ab}_{F^+_R}\times W^{ab}_{F^+_L}) = G^{(n)}(\widetilde F^+_{\overline{v}_\oplus}\times \widetilde F^+_{v_\oplus})$ where $G^{(n)}(\widetilde F^+_{\overline{v}_\oplus}\times \widetilde F^+_{v_\oplus})$ is the non-compact real bilinear algebraic semigroup with entries in $\widetilde F^+_{\overline{v}_\oplus}\times \widetilde F^+_{v_\oplus}$.
2. The Fulton-McPherson compactification by blowups transforms $G^{(n)}(\widetilde F^+_{\overline{v}}\times \widetilde F^+_v)$ into the compact real bilinear algebraic semigroup $G^{(n)}(F^+_{\overline{v}}\times F^+_v)$.
3. The power series development of $f_{\overline{v}}(-x)\otimes f_v(x)$ on $G^{(n)}(F^+_{\overline{v}_\oplus}\times F^+_{v_\oplus})$ constitutes an irreducible functional representation space of $G^{(n)}(F^+_{\overline{v}}\times F^+_v)$ according to corollary 3.1.12. ■

3.2 General bilinear cohomology theory

3.2.1 Introduction of bilinear cohomology

As the envisaged mathematical structures are bilinear, we have to introduce a bilinear cohomology in one-to-one correspondence with its linear equivalent.
The existence of a bilinear cohomology is justified by:

- the cycle map;
- the Tannakian category of representations of affine groups schemes.

The considered bilinear cohomology will be "general" in the sense that it will be a cohomology theory on abstract bisemivarieties in one-to-one correspondence with the associated affine algebraic bisemivarieties covering those [Hart].
The affine algebraic bisemivarieties will be affine bilinear algebraic semigroups $\boldsymbol{G^{(n)}(\widetilde F^+_{\overline v}\times\widetilde F^+_v)}$ embedded isomorphically in their complex equivalents $\boldsymbol{G^{(n)}(\widetilde F_{\overline\omega}\times \widetilde F_\omega)}$ and covering the complete algebraic bilinear semigroups $\boldsymbol{G^{(n)}(F^+_{\overline v}\times F^+_v)}$.
Note that the conjugacy classes of $G^{(n)}(F^+_{\overline v}\times F^+_v)$ are isomorphic to these of $G^{(n)}(\widetilde F^+_{\overline v}\times\widetilde F^+_v)$ by a suitable compactification described in section 3.1.

The following proposition will define and justify **the general bilinear cohomology theory on smooth abstract (resp. algebraic) bisemivarieties** $G^{(n)}(F^+_{\overline v}\times F^+_v)$ (resp. $G^{(n)}(\widetilde F^+_{\overline v}\times\widetilde F^+_v)$) **as the (graded) right derived bifunctor $\boldsymbol{H^{2*}(G^{(n)}(F^+_{\overline v}\times F^+_v),\mathrm{FREPSP}(\mathrm{GL}_{2*}(F^+_{\overline v}\times F^+_v)))}$ (resp. $\boldsymbol{H^{2*}(G^{(n)}(\widetilde F^+_{\overline v}\times\widetilde F^+_v),\mathrm{FREPSP}(\mathrm{GL}_{2*}(\widetilde F^+_{\overline v}\times \widetilde F^+_v)))}$** of the functional representation spaces of the complete (resp. algebraic) bilinear parabolic subsemigroups $P^{2*}(F^+_{\overline v^1}\times F^+_{v^1})$ (resp. $P^{2*}(\widetilde F^+_{\overline v^1}\times\widetilde F^+_{v^1})$) having trivial actions where $\mathrm{FREPSP}(\mathrm{GL}_{2*}(\widetilde F^+_{\overline v}\times\widetilde F^+_v)$ denotes the functional representation spaces of the graded bilinear algebraic semigroups

$$\mathrm{GL}_{2*}(\widetilde F^+_{\overline v}\times\widetilde F^+_v)=\bigoplus_{i\le n}\mathrm{GL}_{2i}(\widetilde F^+_{\overline v}\times\widetilde F^+_v)\ .$$

In connection with the bilinear cohomology semigroup $H^{2*}(G^{(n)}(F^+_{\overline v}\times F^+_v),\mathrm{FREPSP}(\mathrm{GL}_{2*}(F^+_{\overline v}\times F^+_v)))$ on the smooth abstract bisemivariety $G^{(n)}(F^+_{\overline v}\times F^+_v)$, there exists a morphism

$$\begin{aligned}\mathrm{H}^{\,2*\to[*,*]}:\quad & H^{2*}(G^{(n)}(F^+_{\overline v}\times F^+_v),\mathrm{FREPSP}(\mathrm{GL}_{2*}(F^+_{\overline v}\times F^+_v)))\\ &\qquad\longrightarrow H^{[*,*]}(G^{(n)}_d(F^+_{\overline v}\times F^+_v),\Omega^{*+*}_{G^{(n)}_d(F^+_{\overline v}\times F^+_v)})\end{aligned}$$

into the bilinear cohomology of the de Rham type $H^{[*,*]}(G^{(n)}_d(F^+_{\overline v}\times F^+_v),\Omega^{*+*}_{G^{(n)}_d(F^+_{\overline v}\times F^+_v)})$ where:

- the dimension $*$ on the left in $[*,*]$ is a covariant cohomological dimension and the dimension $*$ on the right is a contravariant cohomological dimension;
- $G^{(n)}_d(F^+_{\overline{v}}\times F^+_v)$ is a differentiable smooth abstract bisemivariety;
- $\Omega^{*+*}_{G^{(n)}_d(F^+_{\overline{v}}\times F^+_v)}=\bigoplus\limits_i\Omega^{i+i}_{G^{(n)}_d(F^+_{\overline{v}}\times F^+_v)}$ are the graded bisemisheaves of differential $(i+i)$-forms on $G^{(n)}_d(F^+_{\overline{v}}\times F^+_v)$, i.e. the wedge product of right semisheaves of differential $(i,0)$-forms by left semisheaves of differential $(0,i)$-forms.

3.2.2 Proposition (General bilinear cohomology)

A general bilinear cohomology theory is a contravariant bifunctor

$$\begin{aligned}H^{2*}:\quad&\{\textit{smooth abstract (resp. algebraic) bisemivarieties}\quad G^{(n)}(F^+_{\overline{v}}\times F^+_v)\\&\qquad\textit{(resp.}\ \ G^{(n)}(\widetilde F^+_{\overline{v}}\times \widetilde F^+_v)\ )\}\\&\longrightarrow\{\textit{graded (functional) representation spaces of the complete}\\&\qquad\textit{(resp. algebraic) bilinear semigroups}\ \ \mathrm{GL}_{2*}(F^+_{\overline{v}}\times F^+_v)\\&\qquad\textit{(resp.}\ \ \mathrm{GL}_{2*}(\widetilde F^+_{\overline{v}}\times \widetilde F^+_v)\ )\}\end{aligned}$$

together with a natural transformation:

$$\begin{aligned}n_{H^{2*}\to H^{[*,*]}}:\quad&H^{2*}(G^{(n)}(F^+_{\overline{v}}\times F^+_v),\mathrm{FREPSP}(\mathrm{GL}_{2*}(F^+_{\overline{v}}\times F^+_v)))\\&\textit{(resp.}\ \ H^{2*}(G^{(n)}(\widetilde F^+_{\overline{v}}\times \widetilde F^+_v),\mathrm{FREPSP}(\mathrm{GL}_{2*}(\widetilde F^+_{\overline{v}}\times \widetilde F^+_v)))\ )\\&\longrightarrow H^{[*,*]}(G^{(n)}_d(F^+_{\overline{v}}\times F^+_v),\Omega^{*+*}_{G^{(n)}_d(F^+_{\overline{v}}\times F^+_v)})\end{aligned}$$

into the bilinear cohomology of the de Rham type of dimension $*+*$.

Let $H^{2i}(G^{(n)}(F^+_{\overline{v}}\times F^+_v),\mathrm{FREPSP}(\mathrm{GL}_{2i}(F^+_{\overline{v}}\times F^+_v)))$ *be the general bilinear cohomology semigroup of dimension* $2i$.
Let $H^{2i}(G^{(n)}(\widetilde F^+_{\overline{v}-v}))$ *be the associated linear cohomology group where* $\widetilde F^+_{\overline{v}-v}=\widetilde F^+_{\overline{v}}\cup\widetilde F^+_v$ *is the set of algebraic extensions corresponding to* $F^+_{\overline{v}}\times F^+_v$.

The linear cohomology $H^{2i}(G^{(n)}(\widetilde F^+_{\overline{v}-v}))$ *is characterized by the cycle map:*

$$\gamma^i_{G^{(n)}_{\overline{v}-v}}:\quad \mathcal{Z}^i(G^{(n)}(\widetilde F^+_{\overline{v}-v}))\longrightarrow H^{2i}(G^{(n)}(\widetilde F^+_{\overline{v}-v}))$$

where $\mathcal{Z}^i(G^{(n)}(\widetilde F^+_{\overline{v}-v}))$ *is the group of algebraic cycles of codimension* i *over the linear algebraic group* $G^{(n)}(\widetilde F^+_{\overline{v}-v})$.

The bilinear cohomology $H^{2i}(G^{(n)}(F^+_{\overline{v}}\times F^+_v),\mathrm{FREPSP}(\mathrm{GL}_{2i}(F^+_{\overline{v}}\times F^+_v))$ *is characterized by the bisemicycle map*

$$\gamma^i_{G^{(n)}_{\overline{v}\times v}}:\quad \not{Z}^{\,i}(G^{(n)}(F^+_{\overline{v}}\times F^+_v))\longrightarrow H^{2i}(G^{(n)}(F^+_{\overline{v}}\times F^+_v)),\mathrm{FREPSP}(\mathrm{GL}_{2i}(F^+_{\overline{v}}\times F^+_v))$$

where $\not{Z}^{\,i}(G^{(n)}(F^+_{\overline{v}}\times F^+_v))$ *is the bilinear semigroup of compactified bisemicycles of codimension i on the bilinear complete semigroup* $G^{(n)}(F^+_{\overline{v}}\times F^+_v)$ *in such a way that* $FREPSP(\mathrm{GL}_{2i}(F^+_{\overline{v}}\times F^+_v))$ *be embedded isomorphically in its complex equivalent* $FREPSP(\mathrm{GL}_{2i}(F^+_{\overline{\omega}}\times F^+_\omega))$ *: this is at the origin of this cycle map.*
The associated the de Rham bilinear cohomology is characterized by the bisemicycle map:

$$\gamma^{[i,i]}_{G^{(n)}_{\overline{v}-v}}:\quad \not{Z}^{\,[i,i]}(G^{(n)}_d(F^+_{\overline{v}}\times F^+_v))\longrightarrow H^{[i,i]}(G^{n)}_d(F^+_{\overline{v}}\times F^+_v),\Omega^{*+*}_{G^{(n)}_d(F^+_{\overline{v}}\times F^+_v)}$$

where $\not{Z}^{\,[i,i]}(G^{(n)}_d(F^+_{\overline{v}}\times F^+_v))$ *is the bilinear semigroup of compactified bisemicycles of codimension $2i=i+i$ on the bilinear complete semigroup* $G^{(n)}_d(F^+_{\overline{v}}\times F^+_v)$ *.*

The correspondence between the linear cohomology and the associated bilinear cohomologies is given by the commutative diagram*:*

$$\begin{tikzcd}
\textbf{Linear} & \textbf{Bilinear} & \textbf{Bilinear} \\
\not{Z}^{\,i}(G^{(n)}(\widetilde{F}^+_{\overline{v}-v})) \arrow[r] \arrow[d, "\gamma^i_{G^{(n)}_{\overline{v}-v}}"] & \not{Z}^{\,[i,i]}(G^{(n)}_d(F^+_{\overline{v}}\times F^+_v)) \arrow[r] \arrow[d, "\wr\ \gamma^{[i,i]}_{G^{(n)}_{\overline{v}-v}}"] & \not{Z}^{\,i}(G^{(n)}(F^+_{\overline{v}}\times F^+_v)) \arrow[d, "\wr\ \gamma^i_{G^{(n)}_{\overline{v}\times v}}"] \\
H^{2i}(G^{(n)}(\widetilde{F}^+_{\overline{v}-v})) \arrow[r, "\sim"] & H^{[i,i]}(G^{(n)}_d(F^+_{\overline{v}}\times F^+_v), \arrow[r] & H^{2i}(G^{(n)}(F^+_{\overline{v}}\times F^+_v), \\
 & \Omega^{i+i}_{G^{(n)}_d(F^+_{\overline{v}}\times F^+_v))} & \mathrm{FREPSP}(\mathrm{GL}_{2i}(F^+_{\overline{v}}\times F^+_v)))
\end{tikzcd}$$

Proof: Remark first that the general bilinear cohomology theory is induced from:

a) The cycle map $\gamma^i_{G^{(n)}_{\overline{v}\times v}}$ which is related to the linear cycle map $\gamma^i_{G^{(n)}_{\overline{v}-v}}$ and to the bilinear the de Rham cycle map $\gamma^{[i,i]}_{G^{(n)}_{\overline{v}-v}}$.

b) the Tannakian category of representations of affine group schemes [Del4].

The commutative diagram of this proposition clearly shows that the linear cycle map $\gamma^i_{G^{(n)}_{\overline{v}-v}}$ proceeds from the bilinear cycle map of the de Rham type $\gamma^{[i,i]}_{G^{(n)}_{\overline{v}-v}}$ and from the isomorphism:

$$H^{2i}(G^{(n)}(\widetilde{F}^+_{\overline{v}-v}))\xrightarrow{\ \sim\ }H^{[i,i]}(G^{(n)}_d(F^+_{\overline{v}}\times F^+_v),\Omega^{i+i}_{G^{(n)}_d(F^+_{\overline{v}}\times F^+_v)})\ .$$

Note that **the bilinear cohomology of the de Rham type** $H^{[i,i]}(G_d^{(n)}(F_{\overline{v}}^+ \times F_v^+)$ **has dimension** $\boldsymbol{2i = i + i}$ (as the corresponding bilinear semigroup of compactified bisemicycles $\not{Z}^{[i,i]}(G_d^{(n)}(F_{\overline{v}}^+ \times F_v^+))$ from the consideration of bisemisheaves of differential $(i+i)$-forms **while the general bilinear cohomology** $H^{2i}(G^{(n)}F_{\overline{v}}^+ \times F_v^+), \mathrm{FREPSP}(\mathrm{GL}_{2i}(F_{\overline{v}}^+ \times F_v^+))$ **has dimension "$\boldsymbol{2i}$" due to its "diagonal" isomorphic embedding into** $H^{2i}(G^{(n)}(F_{\overline{\omega}} \times F_\omega), \mathrm{FREPSP}(\mathrm{GL}_{2i}(F_{\overline{\omega}} \times F_\omega))$. ■

3.2.3 Proposition

The general bilinear cohomology is a general cohomology theory in the sense that:

- *it is a motivic (bilinear) cohomology theory or a Weil (bilinear) cohomology theory;*
- *it is directly related to the standard cohomology theories like the singular, de Rham and Betti cohomologies.*

Proof:

1. The general bilinear cohomology $H^{2*}(G^{(n)}(F_{\overline{v}}^+ \times F_v^+)), \mathrm{FREPSP}(\mathrm{GL}_{2*}(F_{\overline{v}}^+ \times F_v^+))$ is directly related to:
   - the Betti cohomology because the complete (algebraic) bilinear semigroup $G^{(n)}(F_{\overline{v}}^+ \times F_v^+)$ is covered by the affine bilinear algebraic semigroup $G^{(n)}(\widetilde{F}_{\overline{v}}^+ \times \widetilde{F}_v^+)$ which is embedded isomorphically in its complex equivalent $G^{(n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$.
     Thus, we have an embedding $\sigma_{R\times L} : F_{\overline{v}}^+ \times F_v^+ \hookrightarrow \mathbb{C} \times \mathbb{C}$.
     This allows to define a bilinear Betti cohomology which must be in one-to-one correspondence with the classical linear Betti cohomology according to proposition 3.2.2.
   - the de Rham cohomology if the considered complete (algebraic) bilinear semigroup is differentiable, i.e. is $G_d^{(n)}(F_{\overline{v}}^+ \times F_v^+)$.
     The obtained de Rham cohomology $H^{[*,*]}(G_d^{(n)}(F_{\overline{v}}^+ \times F_v^+), \Omega^{*+*}_{G_d^{(n)}(F_{\overline{v}}^+ \times F_v^+)})$ is bilinear, but, by the commutative diagram of proposition 3.2.2, it is in one-to-one correspondence with the classical linear de Rham cohomology.
   - the singular bilinear cohomology which is in one-to-one correspondence with the de Rham bilinear cohomology.
2. The general bilinear cohomology is a motivic or Weil (bilinear) cohomology theory if it verifies the standard conjectures on algebraic (bi)cycles as developed in the following proposition. ■

3.2.4 Proposition

The general bilinear cohomology theory, defined by the contravariant bifunctor H^{2*} *and the natural transformation* $n_{H^{2*}\to H^{[*,*]}}$ *, is characterized by:*

1. *Hodge bisemicycles* $H^{i+i}(G^{(n)}(F_{\overline{\omega}}\times F_{\omega}),\mathrm{FREPSP}(\mathrm{GL}_{i+i}(F^{+}_{\overline{v}}\times F^{+}_{v}))$ *(resp.* $H^{i+i}(G^{(n)}(\widetilde{F}^{+}_{\overline{\omega}}\times \widetilde{F}^{+}_{\omega}),\mathrm{FREPSP}(\mathrm{GL}_{i+i}(\widetilde{F}^{+}_{\overline{v}}\times \widetilde{F}^{+}_{v}))$ *),* $2i=i+i$ *, on abstract (resp. algebraic) complex bisemivarieties into "real" functional representation spaces of* $\mathrm{GL}_{i+i}(F^{+}_{\overline{v}}\times F^{+}_{v})$ *in such a way that there is a bifiltration* $F^{p}_{R\times L}$ *on the right and left cohomology semigroups of* $H^{2i}(G^{(n)}(\bullet\times\bullet),-)$ *given by:*

   $$F^{p}_{R\times L}H^{2i}(G^{(n)}(\bullet\times\bullet),-)=\bigoplus_{i=p+q}H^{2(p+q)}(G^{(n)}(\bullet\times\bullet),-)\ .$$

2. *a Künneth standard conjecture implying that the projectors*

   $$H^{2*}(G^{(n)}(\bullet\times\bullet),-)\longrightarrow H^{2i}(G^{(n)}(\bullet\times\bullet),-)$$

   are induced by algebraic bisemicycles $\mathrm{CY}^{i}(G^{(n)}(\widetilde{F}^{+}_{\overline{v}}\times \widetilde{F}^{+}_{v}))\subset \mathcal{Z}^{i}(G^{(n)}(\widetilde{F}^{+}_{\overline{v}}\times \widetilde{F}^{+}_{v}))$ *decomposing into rational subbisemicycles according to the conjugacy class representatives of* $\mathrm{GL}_{2i}(\widetilde{F}^{+}_{\overline{v}}\times \widetilde{F}^{+}_{v})$ *.*

3. *a Künneth biisomorphism:*

   $$\begin{aligned}&H^{2*}(G^{(n)}(F^{+}_{\overline{v}}),\mathrm{FREPSP}(\mathrm{GL}_{2*}(F^{+}_{\overline{v}})))\otimes_{F^{+}_{\overline{v}}\times F^{+}_{v}}H^{2*}(G^{(n)}(F^{+}_{v}),\mathrm{FREPSP}(\mathrm{GL}_{2*}(F^{+}_{v})))\\ &\qquad\longrightarrow H^{2*}(G^{(n)}(F^{+}_{\overline{v}}\times F^{+}_{v}),\mathrm{FREPSP}(\mathrm{GL}_{2*}(F^{+}_{\overline{v}}\times F^{+}_{v})))\end{aligned}$$

   associated with the exact sequence:

   $$\begin{aligned}0\longrightarrow&\sum_{i=p+q}H^{2i}(G^{(n)}(F^{+}_{\overline{v}}\times F^{+}_{v}),\mathrm{FREPSP}(\mathrm{GL}_{2p+2q}(F^{+}_{\overline{v}}\times F^{+}_{v})))\\ &\xrightarrow{\mathcal{H}_{i-0}}\sum_{i=p+q}H^{2i}(G^{(n)}(F^{+}_{\overline{v}}\times F^{+}_{v}),\mathrm{FREPSP}(\mathrm{GL}_{2p+2q}(F^{+}_{\overline{v}}\times F^{+}_{v})))\\ &\qquad+H^{0}(G^{(n)}(F^{+}_{\overline{v}}\times F^{+}_{v}),\mathrm{FREPSP}(\mathrm{GL}_{1}(F^{+}_{\overline{v}}\times F^{+}_{v})))\\ &\qquad\longrightarrow H^{0}(G^{(n)}(F^{+}_{\overline{v}}\times F^{+}_{v}),\mathrm{FREPSP}(\mathrm{GL}_{1}(F^{+}_{\overline{v}}\times F^{+}_{v})))\longrightarrow 0\end{aligned}$$

   in such a way that

   $$\begin{aligned}&H^{2p}(G^{(n)}(F^{+}_{\overline{v}}\times F^{+}_{v}),\mathrm{FREPSP}(\mathrm{GL}_{2p}(F^{+}_{\overline{v}}\times F^{+}_{v})))\\ &\qquad\otimes H^{2i-2p}(G^{(n)}(F^{+}_{\overline{v}}\times F^{+}_{v}),\mathrm{FREPSP}(\mathrm{GL}_{2i-2p}(F^{+}_{\overline{v}}\times F^{+}_{v})))\\ &\qquad\qquad\longrightarrow H^{0}(G^{(n)}(F^{+}_{\overline{v}}\times F^{+}_{v}),\mathrm{FREPSP}(\mathrm{GL}_{1}(F^{+}_{\overline{v}}\times F^{+}_{v})))\end{aligned}$$

   is the bilinear version of the intersection cohomology according to the Goresky-Mac Pherson approach [G-MP] *and referring to the Poincare duality.*

Proof:

1. The Hodge bisemicycles $H^{i+i}(G^{(n)}(F_{\overline{\omega}} \times F_\omega), \mathrm{FREPSP}(\mathrm{GL}_{i+i}(F^+_{\overline{v}} \times F^+_v))$ are justified because $G^{(n)}(F_{\overline{\omega}} \times F_\omega)$ is an abstract complex bisemivariety embedding its real equivalent $G^{(n)}(F^+_{\overline{v}} \times F^+_v)$. So, $G^{(n)}(F_{\overline{\omega}} \times F_\omega)$ is a Kähler bisemivariety on which a Hodge decomposition [D-M-O-S]an be performed.

   Since abstract bisemivarieties are concerned, the Hodge decomposition must apply on the right and left semivarieties $G^{(n)}(F_{\overline{\omega}})$ and $G^{(n)}(F_\omega)$ of $G^{(n)}(F_{\overline{\omega}} \times F_\omega)$ and Hodge bisemicycles follow themselves.

   Similarly, a Hodge bifiltration on right and left (abstract) bisemivarieties of $G^{(n)}(F^+_{\overline{v}} \times F^+_v)$ is justified.

2. The Künneth standard conjecture dealing with the projectors $H^{2*}(G^{(n)}(\bullet \times \bullet), -) \to H^{2i}(G^{(n)}(\bullet \times \bullet), -)$ are induced from algebraic bisemicycles $\mathrm{CY}^i(G^{(n)}(\widetilde{F}^+_{\overline{v}} \times \widetilde{F}^+_v))$ because the concerned cohomologies are bilinear cohomologies of abstract bisemivarieties.

   On the other hand, these projectors refer to mappings interpreted as inverse quantum deformations of Galois representations as introduced in [Pie6] and applied in [Pie2].

3. The considered Künneth biisomorphism takes into account the intersection cohomology which is a pairing between the $2p$-th and $2(i-p)$-th bilinear cohomologies in the 0-th bilinear cohomology $H^0(G^{(n)}(\bullet \times \bullet), -)$ yielding a 0-dimensional bisemicycle because the intersection is composed of a finite set of bipoints. ■

3.3 Borel-Serre toroidal compactification

3.3.1 General bilinear toroidal bisemivariety

A following step is the transformation of the bilinear general toric bisemivariety $\tau_\sigma^{(n)}(\widetilde{F}^+_{\overline{v}} \times \widetilde{F}^+_v)$ (or $\tau_{c_\sigma}^{(n)}(\widetilde{F}^+_{\overline{v}} \times \widetilde{F}^+_v)$) into the bilinear general toroidal bisemivariety $\tau_T^{(n)}(F^{+,T}_{\overline{v}} \times F^{+,T}_v)$. This can be realized by considering the projective emergent isomorphism $\gamma^c_R \times \gamma^c_L$ from the bilinear algebraic semigroup $G^{(n)}(\widetilde{F}^+_{\overline{v}} \times \widetilde{F}^+_v)$, corresponding to $\tau_\sigma^{(n)}(\widetilde{F}^+_{\overline{v}} \times \widetilde{F}^+_v)$ or to $\tau_{c_\sigma}^{(n)}(\widetilde{F}^+_{\overline{v}} \times \widetilde{F}^+_v)$, to its toroidal equivalent $G^{(n)}(F^{+,T}_{\overline{v}} \times F^{+,T}_v)$, corresponding to $\tau_T^{(n)}(F^{+,T}_{\overline{v}} \times F^{+,T}_v)$ as considered in the following section: in fact, the projective emergent isomorphism $\gamma^c_R \times \gamma^c_L$ will be more precisely developed for the complex bilinear algebraic semigroup $G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$, taking into account the inclusion

$$G^{(n)}(\widetilde{F}^+_{\overline{v}} \times \widetilde{F}^+_v) \hookrightarrow G^{(2n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega) \; .$$

A toroidal compactification will been envisaged for the lattice bisemispace $X_{S_{R\times L}}$: it will correspond to the Borel-Serre toroidal compactification which will be decomposed into a two step sequence which will lead to the equivalent of a Shimura (bisemi)variety. This compactification is in fact a projective mapping of $X_{S_{R\times L}}$, generating products of pairs of embedded complex semitori (which are the only complex compact Lie (semi)groups).

3.3.2 Definition

The left (resp. right) toroidal projective emergent (iso)morphism $\gamma_L^C : X_{S_L} \to \overline{X}_{S_L}$ (resp. $\gamma_R^C : X_{S_R} \to \overline{X}_{S_R}$) can be decomposed into the two steps sequence [Pie1]:

a) the points $P_{a_{L[j,m_j]}} \in \tilde{g}_L^{(2n)}[j,m_j]$ (resp. $P_{a_{R[j,m_j]}} \in \tilde{g}_R^{(2n)}[j,m_j]$) are mapped onto the origin of $\widetilde{F}_\omega^{2n}$ (resp. $\widetilde{F}_{\overline{\omega}}^{2n}$).

b) these points $P_{a_{L[j,m_j]}}$ (resp. $P_{a_{R[j,m_j]}}$) are then projected symmetrically from the origin of $\widetilde{F}_\omega^{n}$ (resp. $\widetilde{F}_{\overline{\omega}}^{n}$) into a connected compact complete semivariety which is a complex semitorus $T_L^{2n}[j,m_j]$ (resp. $T_R^{2n}[j,m_j]$) in $(F_\omega^T)^n$ (resp. $(F_{\overline{\omega}}^T)^n$) where $F_\omega^T = \{F_{\omega_1}^T,\cdots,F_{\omega_{j,m_j}}^T,\cdots,F_{\omega_r}^T\}$ (resp. $F_{\overline{\omega}}^T = \{F_{\overline{\omega}_1}^T,\cdots,F_{\overline{\omega}_{j,m_j}}^T,\cdots,F_{\overline{\omega}_r}^T\}$) is the set of toroidal (compactified) completions $F_{\omega_{j,m_j}}^T$ (resp. $F_{\overline{\omega}_{j,m_j}}^T$) corresponding to $F_{\omega_{j,m_j}}$(resp. $F_{\overline{\omega}_{j,m_j}}$).

This left (resp. right) projective emergent morphism γ_L^C (resp. γ_R^C) is an isomorphism because it is characterized by its representatives $h_{a_{L[j,m_j]}}$ (resp. $h_{a_{R[j,m_j]}}$) given by the triple

$$\begin{aligned} & h_{a_{L[j,m_j]}} = \{P_{a_{L[j,m_j]}}, r(P_{a_{L[j,m_j]}}), \gamma_{a_{L[j,m_j]}}\} \\ \text{(resp.} \quad & h_{a_{R[j,m_j]}} = \{P_{a_{R[j,m_j]}}, r(P_{a_{R[j,m_j]}}), \gamma_{a_{R[j,m_j]}}\} \text{)} \end{aligned}$$

where:

- $r(P_{a_{L[j,m_j]}})$ (resp. $r(P_{a_{R[j,m_j]}})$) is the euclidian distance, from the origin, of a point $P_{a_{L[j,m_j]}}$ (resp. $P_{a_{R[j,m_j]}}$) projected into the $2n$-dimensional semitorus $T^{2n}_{L[j,m_j]}$ (resp. $T^{2n}_{R[j,m_j]}$);

- $\gamma_{a_{L[j,m_j]}}$ (resp. $\gamma_{a_{R[j,m_j]}}$) is a one-to-one correspondence between the point $P_{a_{L[j,m_j]}}$ (resp. $P_{a_{R[j,m_j]}}$) and its projective localization given by $r(P_{a_{L[j,m_j]}})$ (resp. $r(P_{a_{R[j,m_j]}})$).

3.3.3 Proposition

The projective emergent isomorphism $\gamma^C_{R\times L} : X_{S_{R\times L}} \to \overline{X}_{S_{R\times L}}$ *, mapping the pseudo-ramified lattice bisemispace* $X_{S_{R\times L}} = \mathrm{GL}_n(\widetilde{F}_R\times\widetilde{F}_L)\Big/ \mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)$ *into the corresponding toroidal compactified lattice bisemispace:* $\overline{X}_{S_{R\times L}} = \mathrm{GL}_n(F^T_R \times F^T_L\Big/ \mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)$ *, is such that* $X_{S_{R\times L}}$ *may be viewed as the interior of its compactified corresponding* $\overline{X}_{S_{R\times L}}$ *in the sense of the Borel-Serre compactification.*

Proof: according to definition 3.3.2, the projective emergent isomorphism sends all equivalent representatives of conjugacy classes into their toroidal compactified equivalents $g^{(2n)}_{T_{R\times L}}[j,m_j]$ consisting in products of n-dimensional complex semitori $T^{2n}_R[j,m_j]\times T^{2n}_L[j,m_j]$.
The inclusion $X_{S_{R\times L}} \hookrightarrow \overline{X}_{S_{R\times L}}$ is a homotopy equivalence so that $X_{S_{R\times L}}$ may be considered as the interior of $\overline{X}_{S_{R\times L}}$. ■

3.3.4 Double coset decomposition

A double coset decomposition of the bilinear toroidal semigroup $\mathrm{GL}_n(F^T_R \times F^T_L)$ gives rise to the compactified bisemispace:

$$\overline{S}^{P_n}_{K_n} = P_n(F^T_{\overline{\omega}^1} \times F^T_{\omega^1}) \setminus \mathrm{GL}_n(F^T_R \times F^T_L\Big/ \mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)$$

where $P_n(F^T_{\overline{\omega}^1} \times F^T_{\omega^1})$ is the bilinear parabolic subsemigroup of matrices associated with the complete (algebraic) bilinear parabolic subsemigroup $P^{(2n)}(F^T_{\overline{\omega}^1} \times F^T_{\omega^1})$ introduced in section 2.4.1.
$P_n(F^T_{\overline{\omega}^1} \times F^T_{\omega^1})$ has the Gauss decomposition as developed in the following section.
Remark that

$$\mathrm{GL}_n(F^T_R \times F^T_L)\Big/ \mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2) \simeq P_n(F^T_{\overline{\omega}^1} \times F^T_{\omega^1}) \setminus \mathrm{GL}_n(F^T_R \times F^T_L)$$

since $\mathrm{GL}_n(F^T_R \times F^T_L)\Big/ \mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)$ corresponds to the set of lattices of $(F^T_{\overline{\omega}})^n \times (F^T_{\omega})^n$ [Bor1], [Del3].

3.3.5 Definition: Levi and Gauss decompositions for (complex) parabolic subgroups

$P_n(F^T_{\omega^1})$ (resp. $P_n(F^T_{\overline{\omega}^1})$) has the standard Levi decomposition:

$$P_n(F^T_{\omega^1}) = \mathrm{Z}_n(F^T_{\omega^1}) \cdot U_n(F^T_{\omega^1}) \qquad (\text{resp.} \quad P_n(F^T_{\overline{\omega}^1}) = \mathrm{Z}_n(F^T_{\overline{\omega}^1}) \cdot U_n(F^T_{\overline{\omega}^1})\ )$$

where:

- $\mathrm{Z}_n(F^T_{\omega^1})$ (resp. $\mathrm{Z}_n(F^T_{\overline{\omega}^1})$) is the left (resp. right) centralizer of $F^T_{\omega^1}$ (resp. $F^T_{\overline{\omega}^1}$) in $T_n(F^T_{\omega^1})$ (resp. $T_n(F^T_{\overline{\omega}^1})^t \subset \mathrm{GL}_n(F^T_{\overline{\omega}} \times F^T_{\omega})$ and is represented by diagonal matrices $d_n(F^T_{\omega^1})$ (resp. $d_n(F^T_{\overline{\omega}^1})$);
- $U_n(F^T_{\omega^1})$ (resp. $U_n(F^T_{\overline{\omega}^1})$) is the left (resp. right) unipotent radical of $P_n(F^T_{\omega^1})$ (resp. $P_n(F^T_{\overline{\omega}^1})$) and is represented by upper (resp. lower) unitriangular matrices $u_n(F^T_{\omega^1})$ (resp. $u_n(F^T_{\overline{\omega}^1})^t$).

Similarly, the complex bilinear parabolic subsemigroup $P_n(F^T_{\overline{\omega}^1} \times F^T_{\omega^1}) \equiv P_n(F^T_{\overline{\omega}^1}) \times P_n(F^T_{\omega^1})$) has the Gauss decomposition:

$$P_n(F^T_{\overline{\omega}^1} \times F^T_{\omega^1}) = (D_n(F^T_{\overline{\omega}^1}) \cdot D_n(F^T_{\omega^1}))(UT_n(F^T_{\overline{\omega}^1})^t \cdot UT_n(F^T_{\omega^1}))$$

according to proposition 2.1.4. $P_n(F^T_{\overline{\omega}^1} \times F^T_{\omega^1})$ is a bilinear normal subgroup of $\mathrm{GL}_n(F^T_{\overline{\omega}} \times F^T_{\omega})$ because it is represented by the product of the subgroups of diagonal matrices, isomorphic to maximal (semi)tori, by the subgroups of unipotent matrices [Bor2].

3.3.6 Representation spaces of bilinear algebraic semigroups

a) The double coset decomposition of $\mathrm{GL}_n(F^T_R \times F^T_L)$ is a compactified bisemispace $\overline{S}^{P_n}_{K_n}$ which corresponds to the representation space $\mathrm{Repsp}(\mathrm{GL}_n(F^T_{\overline{\omega}} \times F^T_{\omega}))$ of $\mathrm{GL}_n(F^T_{\overline{\omega}} \times F^T_{\omega})$. So we have

$$\overline{S}^{P_n}_{K_n} \simeq \mathrm{Repsp}(\mathrm{GL}_n(F^T_{\overline{\omega}} \times F^T_{\omega})) = G^{(2n)}(F^T_{\overline{\omega}} \times F^T_{\omega})$$

since $G^{(2n)}(F^T_{\overline{\omega}} \times F^T_{\omega})$ decomposes into conjugacy classes having equivalent representatives $g^{(2n)}_{T_{R\times L}}[j, m_j]$.

b) Considering the Gauss decomposition of $\mathrm{GL}_n(F^T_{\overline{\omega}} \times F^T_{\omega})$ and of $P_n(F^T_{\overline{\omega}^1} \times F^T_{\omega^1})$, we can introduce the double coset decomposition of the diagonal part of $\mathrm{GL}_n(F^T_R \times F^T_L)$ which is a compactified diagonal bisemispace noted $\overline{S}^{Z[P_n]}_{Z[K_n]}$ and given by:

$$\overline{S}^{Z[P_n]}_{Z[K_n]} = D_n(F^T_{\overline{\omega}^1} \times F^T_{\omega^1}) \setminus D_n(F^T_R \times F^T_L) \Big/ D_n((\mathbb{Z}/N\,\mathbb{Z})^2)$$

where:

- $D_n(F^T_{\overline{\omega}^1} \times F^T_{\omega^1}) \subset P_n(F^T_{\overline{\omega}^1} \times F^T_{\omega^1})$ following definition 3.3.5;
- $D_n(F^T_R \times F^T_L) \subset \mathrm{GL}_n(F^T_R \times F^T_L)$.

So, as in a), we can state:

$$\overline{S}^{Z[P_n]}_{Z[K_n]} \simeq \mathrm{Repsp}(D_n(F^T_{\overline{\omega}} \times F^T_{\omega})) = D^{(n)}(F^T_{\overline{\omega}} \times F^T_{\omega})$$

with evident notations.

c) In the following sections, bilinear cohomology semigroups of $\overline{S}^{P_n}_{K_n}$ will be considered with coefficients in complete algebraic representation spaces $\mathrm{Repsp}(\mathrm{GL}_n(F^T_{\overline{\omega}} \times F^T_{\omega}))$ of $\mathrm{GL}_n(F^T_{\overline{\omega}} \times F^T_{\omega})$, $1 \le n \le \infty$.

These representation spaces $\mathrm{Repsp}(\mathrm{GL}_n(F^T_{\overline{\omega}} \times F^T_{\omega}))$ are $\mathrm{GL}_n(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus})$-bisemimodules $M_{T_{R_\oplus}} \otimes M_{T_{L_\oplus}}$ (see proposition 2.1.4). Now, generally, the **coefficients of the (bilinear) cohomology are envisaged in (bisemi)sheaf of rings.** In this purpose, a (semi)sheaf of rings $\widehat{M}^{2i}_{T_R}$ (resp. $\widehat{M}^{2i}_{T_L}$) of complex-valued continuous functions $\phi^{(2i)}_{G^T_{j_R}}(x_{g^T_{j_R}})$ (resp. $\phi^{(2i)}_{G^T_{j_L}}(x_{g^T_{j_L}})$) on the basic conjugacy class representatives $g^{(2i)}_{T_R}[j, m_j = 0]$ (resp. $g^{(2i)}_{T_L}[j, m_j = 0]$) of $\mathrm{Repsp}(T^t_i(F^T_{\overline{\omega}}))$ (resp. $\mathrm{Repsp}(T_i(F^T_{\omega}))$) can thus be envisaged.

On $M^{2i}_{T_R} \otimes M^{2i}_{T_L}$, it is then the bisemisheaf of rings $\widehat{M}^{2i}_{T_R} \otimes \widehat{M}^{2i}_{T_L}$ which has to be taken into account (in this context, see also section 3.4.10).

So, the developments dealing **with bilinear cohomology with coefficients** in $\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})$-bisemimodules $M^{2i}_{T_R} \otimes M^{2i}_{T_L}$, $1 \le i \le \infty$, can be easily transposed to those with coefficients **in bisemisheaves of rings $\widehat{\boldsymbol{M}}^{\boldsymbol{2i}}_{\boldsymbol{T_R}} \otimes \widehat{\boldsymbol{M}}^{\boldsymbol{2i}}_{\boldsymbol{T_L}}$ over $\boldsymbol{M}^{\boldsymbol{2i}}_{\boldsymbol{T_R}} \otimes \boldsymbol{M}^{\boldsymbol{2i}}_{\boldsymbol{T_L}}$** .

3.3.7 Definition: Weil algebra of the bilinear nilpotent Lie algebra

Let

$$p_{G \to \mathrm{Z}[G]} : \quad \mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})) \longrightarrow \mathrm{Repsp}(D_i(F^T_{\overline{\omega}} \times F^T_{\omega}))$$

denote the projective mapping of the smooth principal bundle H^{2i} given by the triple $(\mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})), \mathrm{Repsp}(D_i(F^T_{\overline{\omega}} \times F^T_{\omega})), p_{G \to \mathrm{Z}[G]})$ and having as fiber $\mathcal{F}^{(2i)}_{R \times L} = UT_i(F^T_{\overline{\omega}})^t \times UT_i(F^T_{\omega})$ the product of the unipotent complete subgroups $UT_i(F^T_{\overline{\omega}})^t$ and $UT_i(F^T_{\omega})$.
Let $\mathrm{Lie}(\mathcal{F}^{(2i)}_{R \times L}) = \mathcal{U}_i(F^T_{\overline{\omega}}) \times \mathcal{U}_i(F^T_{\omega})$ be the Lie algebra of the fiber $\mathcal{F}^{(2i)}_{R \times L}$.
The exterior algebra $A(\mathrm{Lie}(\mathcal{F}^{(2i)}_{R \times L}))$ on $\mathrm{Lie}(\mathcal{F}^{(2i)}_{R \times L})$ is the algebra of products of differential forms of all degrees. $\wedge(\mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})))$ will denote the graded differential algebra of differential forms of $\mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega}))$ and $\wedge(\mathrm{Repsp}(D_i(F^T_{\overline{\omega}} \times F^T_{\omega})))$ will denote the graded differential algebra of differential forms of $\mathrm{Repsp}(D_i(F^T_{\overline{\omega}} \times F^T_{\omega}))$.
It is evident that:

$$\wedge(\mathrm{Repsp}(D_i(F^T_{\overline{\omega}} \times F^T_{\omega}))) \subset \mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega}))) \ .$$

Let $S(\mathrm{Lie}(\mathcal{F}^{(2i)}_{R \times L}))$ denote the symmetric algebra on $\mathrm{Lie}(\mathcal{F}^{(2i)}_{R \times L})$ which can be identified to the algebra of symmetric multilinear forms on $\mathrm{Lie}(\mathcal{F}^{(2i)}_{R \times L})$.

Then, the Weil algebra of the Lie algebra $\mathrm{Lie}(\mathcal{F}^{(2i)}_{R\times L})$ is by definition the graded algebra [Car]:

$$W(\mathrm{Lie}(\mathcal{F}^{(2i)}_{R\times L})) = A(\mathrm{Lie}(\mathcal{F}^{(2i)}_{R\times L})) \times S(\mathrm{Lie}(\mathcal{F}^{(2i)}_{R\times L}))\ .$$

3.3.8 Definition

A connection on the fibered space $\mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega}))$ consists in a bilinear mapping $f_{R\times L}$ of $A^1(\mathrm{Lie}(\mathcal{F}^{(2i)}_{R\times L}))$ in the subspace of the elements of degree one of the algebra $\wedge(\mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})))$.

$f_{R\times L}$ can be extended as a homomorphism:

$$f'_{R\times L} : \quad A(\mathrm{Lie}(\mathcal{F}^{(2i)}_{R\times L})) \longrightarrow \wedge\,(\mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})))\ .$$

On the other hand, according to [G-H-V] and [Car], the connection $f_{R\times L}$ generates a homomorphism from $I_s(\mathrm{Lie}(\mathcal{F}^{(2i)}_{R\times L}))$, which is the subalgebra of invariant elements of $S(\mathrm{Lie}(\mathcal{F}^{(2i)}_{R\times L}))$ identified to the algebra of symmetric multilinear forms $\vee(\mathrm{Lie}(\mathcal{F}^{(2i)}_{R\times L}))$ on $\mathrm{Lie}(\mathcal{F}^{(2i)}_{R\times L})$, into $H^{2n}(\wedge(\mathrm{Repsp}(D_i(F^T_{\overline{\omega}} \times F^T_{\omega}))))$: this is the Chern-Weil homomorphism

$$h_{R\times L} : \quad \vee(\mathrm{Lie}(\mathcal{F}^{(2i)}_{R\times L})) \longrightarrow H^{2i}(\wedge(\mathrm{Repsp}(D_i(F^T_{\overline{\omega}} \times F^T_{\omega}))))$$

were $H^{2i}(\cdot)$ defines a Hodge bilinear (i,i) class also written $i+i$ in section 3.2.

3.3.9 Proposition

Let $(M^{2i}_{T_R} \otimes M^{2i}_{T_L})$ *be the representation space of* $\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})$.

Then, the existence of a biconnection associated with the homomorphism $\mathcal{I}_s(\mathrm{Lie}(\mathcal{F}^{(2i)}_{R\times L})) \to \wedge(\mathrm{Repsp}(D_i(F^T_{\overline{\omega}} \times F^T_{\omega})))$ *is equivalent to the existence of a nilpotent fiber* $\mathcal{F}^{(2i)}_{R\times L}$ *in the smooth principal bundle* $\mathrm{H}^{2i}(\mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})), \mathrm{Repsp}(D_i(F^T_{\overline{\omega}} \times F^T_{\omega})), p_{G\to \mathrm{Z}\,[G]})$ *which implies that:*

$$H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_R} \otimes M^{2i}_{T_L}) \simeq \mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega}))$$

only if $\mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega}))$ *, which is the fibered space of the principal bundle* $\mathrm{H}^{2i}(\cdot,\cdot,\cdot)$ *, is an irreducible representation space of* $\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})$.

Proof: as the connection $f_{R\times L}$ can be extended to the homomorphism $f'_{R\times L}$ given in definition 3.3.8 and generates the Chern-Weil homomorphism $h_{R\times L}$, it appears that the knowledge of $\mathrm{Repsp}(D_i(F^T_{\overline{\omega}} \times F^T_{\omega}))$, together with the connection $f_{R\times L}$ is sufficient to know the cohomology $H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_R} \otimes M^{2i}_{T_L})$ [Car]. Thus, the action of a connection on $\mathrm{Repsp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega}))$ is equivalent to the action of the nilpotent fiber $\mathcal{F}^{(2i)}_{R\times L}$ on $\mathrm{Repsp}(D_i(F^T_{\overline{\omega}} \times F^T_{\omega}))$. And, we have that the cohomology $H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_R} \otimes M^{2i}_{T_L})$, which is the cohomology of the fibered space, is in one-to-one correspondence with $\text{Repsp}(\text{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega}))$. ■

3.3.10 Proposition

Let $M^{2i}_{T_R} \otimes M^{2i}_{T_L}$ be an irreducible $\text{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})$-subbisemimodule of $(M_{T_R} \otimes M_{T_L})$, $i \le n$. Then, the $2i$-dimensional irreducible representation $\text{Irr}\, W^{(2i)}_{F_{R\times L}}(W^{ab}_{F_R} \times W^{ab}_{F_L})$ of the product of the global Weil groups is given by:

$$\begin{aligned}\text{Irr}\,\text{Rep}\, W^{(2i)}_{F_{R\times L}}(W^{ab}_{F_R} \times W^{ab}_{F_L}) &= G^{(2i)}(\widetilde{F}_{\overline{\omega}_\oplus} \times \widetilde{F}_{\omega_\oplus}) \\ \simeq \text{Repsp}(\text{GL}_i(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus}) &= H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{R_\oplus}} \otimes M^{2i}_{T_{L_\oplus}}) \ .\end{aligned}$$

Proof: indeed, according to proposition 3.1.13, we have that

$$H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{R_\oplus}} \otimes M^{2i}_{T_{L_\oplus}}) = \text{Repsp}(\text{GL}_i(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus})$$

and, according to proposition 2.4.7, we have that:

$$\text{Irr}\, W^{(2i)}_{F_{R\times L}}(W^{ab}_{F_R} \times W^{ab}_{F_L}) = G^{(2i)}(\widetilde{F}_{\overline{\omega}_\oplus} \times \widetilde{F}_{\omega_\oplus}) \ .$$

So, we are led to the isomorphism:

$$\text{Irr}\, W^{(2i)}_{F_{R\times L}}(W^{ab}_{F_R} \times W^{ab}_{R_L}) \simeq H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{R_\oplus}} \otimes M^{2i}_{T_{L_\oplus}})$$

because $\text{Irr}\, W^{(2i)}_{F_{R\times L}}(W^{ab}_{F_R} \times W^{ab}_{F_L})$ refers to not necessarily compact algebraic extensions while $H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_R} \otimes M^{2i}_{T_L})$ is the cohomology of the toroidal compactified bisemispace $\overline{S}^{P_n}_{K_n}$. ■

3.4 Langlands global correspondence on the irreducible algebraic bilinear semigroup $\mathbf{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$

3.4.1 Proposition

The cohomology $H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{R_\oplus}} \otimes M^{2i}_{T_{L_\oplus}})$ has a decomposition according to the equivalent representatives $g^{(2i)}_{T_{R\times L}}[j,m_j]$ of the conjugacy classes of the complete bilinear semigroup $G^{(2i)}(F^T_{\overline{\omega}} \times F^T_{\omega})$ according to:

$$H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{R_\oplus}} \otimes M^{2i}_{T_{L_\oplus}}) = \bigoplus_{j=1}^{r} \bigoplus_{m_j} g^{(2i)}_{T_{R\times L}}[j,m_j] \ .$$

Proof:

$$H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{R_\oplus}} \otimes M^{2i}_{T_{L_\oplus}}) = G^{(2i)}(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus})$$

where $G^{(2i)}(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus})$ decomposes according to the equivalent representatives $g^{(2i)}_{T_{R\times L}}[j,m_j]$ of $G^{(2i)}(F^T_{\overline{\omega}} \times F^T_{\omega})$. So, we get the thesis. ■

3.4.2 Corollary

Let

$$\overline{S}^{Z[P_n]}_{Z[K_n]} = D_n(F^T_{\overline{\omega}^1} \times F^T_{\omega^1}) \setminus D_n(F^T_R \times F^T_L) \Big/ D_n((\mathbb{Z} / N\ \mathbb{Z})^2)$$

be the diagonal compactified bisemispace and let $D^{(2i)}(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus})$ *be the diagonal bilinear semigroup decomposing into conjugacy classes having representatives* $g^{(2i)}_{T_{R\times L}}[j]$ *. Then, we have the following decomposition of the cohomology:*

$$H^{2i}(\overline{S}^{Z[P_n]}_{Z[K_n]}, M^{2i}_{T_{R_\oplus}}(D) \otimes M^{2i}_{T_{L_\oplus}}(D)) =\simeq \bigoplus_j g^{(2i)}_{T_{R\times L}}[j]$$

where $(M^{2i}_{T_{R_\oplus}}(D) \otimes M^{2i}_{T_{L_\oplus}}(D)) = \bigoplus_j (M^{2i}_{T_{\overline{\omega}_j}} \otimes M^{2i}_{T_{\omega_j}})$ *is a* $D_i(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus})$*-bisemimodule.*

Proof: referring to chapter 1, we note that the conjugacy classes of $D^{(2i)}(F^T_{\overline{\omega}} \times F^T_{\omega})$ have unique representatives $g^{(2i)}_{T_{R\times L}}[j]$, since there is no nilpotent action in the diagonal case, the nilpotent action being generated by $UT^t_i(F^T_{\overline{\omega}}) \times UT_i(F^T_{\omega}) \subset \mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})$. ■

3.4.3 Analytic development of the bilinear cohomology $H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_R} \otimes M^{2i}_{T_L})$

The objective consists now in proving that the decomposition of the bilinear cohomology $H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_R} \otimes M^{2i}_{T_L})$ corresponding to the representation space of the irreducible bilinear general semigroup $\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})$ has an analytic development given by the development in truncated Fourier series of the product of a "right" $2i$-dimensional cusp form by its left equivalent.

As the decomposition of the cohomology $H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_R} \otimes M^{2i}_{T_L})$ is given by the sum over j and m_j of equivalent representatives $g^{(2i)}_{T_{R\times L}}[j,m_j] = g^{(2i)}_{T_R}[j,m_j] \times g^{(2i)}_{T_L}[j,m_j]$ consisting in products, right by left, of i-dimensional complex semitori $T^{2i}_R[j,m_j] \times T^{2i}_L[j,m_j]$ according to proposition 3.3.3, an analytic development must be given to these semitori as follows:

3.4.4 Proposition

Let $\vec{z} = \sum\limits_{d=1}^{2i} z_d \vec{e}_d$ *be a vector of* $(F_{\omega_j})^i$ *and, more precisely, a point of* $G^{(2i)}(F_{\omega_j})$.

Then, every left (resp. right) i-dimensional complex semitorus has the analytic development:

$$\begin{aligned} T_L^{2i}[j,m_j] &\simeq \lambda^{\frac{1}{2}}(2i,j,m_j)e^{2\pi ijz} \\ (\text{resp.}\quad T_R^{2i}[j,m_j] &\simeq \lambda^{\frac{1}{2}}(2i,j,m_j)e^{-2\pi ijz}\ ) \end{aligned}$$

where

- $\lambda^{\frac{1}{2}}(2i,j,m_j) \simeq j^{2i}\cdot N^{2i}\cdot (m^{(j)})^{2i}$ *can be considered as a "global" Hecke character;*
- $z=\sum\limits_d z_d\ |\vec{e}_d|$.

Proof:

1. The cohomology $H^{2i}(\overline{S}^{P_n}_{K_n},M^{2i}_{T_R}\otimes M^{2i}_{T_L})$ corresponds to an endomorphism of $\overline{S}^{P_n}_{K_n}$ into itself. Its decomposition into classes " j " with representatives " m_j " refers to the cosets of $\mathrm{GL}_i(F_R^T\times F_L^T)/\mathrm{GL}_i((\mathbb{Z}/N\,\mathbb{Z})^2)$. So, the scalar $\lambda(2i,j,mj)$ will correspond to the eigenvalues of $g_i(\mathcal{O}_{F^T_{\overline{\omega}_{j,m_j}}}\times\mathcal{O}_{F^+_{\omega_{j,m_j}}})\in\mathrm{GL}_i((\mathbb{Z}/N\,\mathbb{Z})^2)$ which is the j-th coset representative $Uj_R\times U_{j_L}$ of the product $T_R(i;r)\otimes T_L(i;r)$ of the Hecke operators according to proposition 2.2.5.

   More precisely, let $\{\lambda_d(2i,j,m_j)\}_{d=1}^{2i}$ be the set of eigenvalues of $U_{j_R}\times U_{j_L}$ and let $\lambda(2i,j,m_j)=\prod\limits_{d=1}^{2i}\lambda_d(2i,j,m_j)$ be the product of these eigenvalues.

   According to definition 2.2.4, we have that

   $$\lambda(2i,j,m_j)=\prod_{d=1}^{2i}\lambda_d(2i,j,m_j)=\det(\alpha_{i^2;j^2}\times D_{j^2;m_{j^2}})_{ss}\simeq j^{2i}\cdot N^{2i}\cdot(m^{(j)})^{2i}$$

   where $D_{j^2;m_{j^2}}$ is the decomposition group of the j-th bisublattice with representative m_j and where $\alpha_{i^2;j^2}$ is the j-th split Cartan subgroup.

   So, it appears that the square root $\lambda^{\frac{1}{2}}(2i,j,m_j)$ of $\lambda(2i,j,m_j)$ can be considered as a global Hecke character having an inflation action on $e^{2\pi ijz}$.

2. On the other hand, as we are concerned with the j-th global or infinite complex place ω_j or $\overline{\omega}_j$, we have to consider the global Frobenius substitution for the left (resp. right) place ω_j (resp. $\overline{\omega}_j$) as given by the mapping:

   $$\begin{aligned} e^{2\pi iz} &\longrightarrow e^{2\pi ijz}\ , \\ (\text{resp.}\quad e^{-2\pi iz} &\longrightarrow e^{-2\pi ijz}\ ) \qquad\qquad z\in G^{(2i)}(F_{\omega_j})\equiv g_L^{(2i)}[j,m_j]\ , \end{aligned}$$

$e^{2\pi iz}$ being a "circular" function on the j_δ^{2i} completions of $g_L^{(2i)}[j,m_j]$ according to proposition 3.1.3.

So, the left (resp. right) semitorus corresponding to the equivalent representative $g_{T_L}^{(2i)}[j,m_j]$ (resp. $g_{T_R}^{(2i)}[j,m_j]$) has the following analytic development:

$$\begin{aligned} T_L^{2i}[j,m_j] &\simeq \lambda^{\frac{1}{2}}(2i,j,m_j)e^{2\pi ijz} \\ \text{(resp.}\quad T_R^{2i}[j,m_j] &\simeq \lambda^{\frac{1}{2}}(2i,j,m_j)e^{-2\pi ijz}\ ) \end{aligned}$$

taking into account that the n-dimensional complex left (resp. right) semitorus $T_L^{2i}[j,m_j]$ (resp. $T_R^{2i}[j,m_j]$) is diffeomorphic to the i-fold product

$$\begin{aligned} T_L^{2i}[j,m_j] &= T_L^2[j,m_j]\times\cdots\times T_L^2[j,m_j] \\ \text{(resp.}\quad T_R^{2i}[j,m_j] &= T_R^2[j,m_j]\times\cdots\times T_R^2[j,m_j]\ ) \end{aligned}$$

of 1-dimensional complex left (resp. right) semitori $T_L^2[j,m_j]$ (resp. $T_R^2[j,m_j]$). These semitori have the analytic development:

$$\begin{aligned} T_L^2[j,m_j] &= \lambda_{T_L^2}\ e^{2\pi ijz_d} \\ &= S_{d_1}^1[j,m_j]\times S_{d_2}^1[j,m_j] && (\ z_d\in\mathbb{C}\ ,\ ) \\ &= r_{S_{d_1}^1}\ e^{2\pi ijx_{d_1}}\times r_{S_{d_2}^1}\ e^{2\pi ij(iy_{d_2})} && (\ x_{d_1}\in\mathbb{R}\ ,\ y_{d_2}\in\mathbb{R}\ ) \\ &\simeq \lambda_{d_1}(1,j,m_j)\cdot\lambda_{d_2}(1,j,m_j)\ e^{2\pi ijx_{d_1}}\times e^{2\pi ij(iy_{d_2})} \end{aligned}$$

where

- $\lambda_{d_1}(1,j,m_j)\cdot\lambda_{d_2}(1,j,m_j)\simeq r_{S_{d_1}^1}\cdot r_{S_{d_2}^1}$ is a product of two eigenvalues of $U_{j_R}\times U_{j_L}$,
- $r_{S_{d_1}^1}$ and $r_{S_{d_2}^1}$ are radii.

Indeed, the left $1D$-complex semitorus $T_L^2[j,m_j]$ is diffeomorphic to the product of two circles

$$S_{d_1}^1[j,m_j]=r_{S_{d_1}^1}\ e^{2\pi ijx_{d_1}}$$

and

$$S_{d_2}^1[j,m_j]=r_{S_{d_2}^1}\ e^{2\pi ij(iy_{d_2})}$$

localized in perpendicular planes with $\cos(2\pi ijy_{d_2})$ and $\sin(2\pi(ijy_{d_2})$ of $e^{2\pi i(ijy_{d_2})}$ defined over $i\mathbb{R}$.

This is justified by the fact that a rotation of 90^0 of the circle $S_{d_2}^1[j,m_j]=r_{S_{d_2}^1}\ e^{2\pi ij(iy_{d_2})}$ over $i\mathbb{R}$ transforms it into the circle $S_{d_{2\perp}}^1[j,m_j]=r_{S_{d_2}^1}\ e^{2\pi ij(y_{d_2})}$ over

$I\!\!R$, localized in the same plane as the circle $S^1_{d_1}[j,m_j]$, according to:

$$\begin{array}{rrcl}
\mathrm{rot}(90^0): & S^1_{d_2}[j,m_j] & \longrightarrow & S^1_{d_{2_\perp}}[j,m_j] \\
& r_{S^1_{d_2}}\ e^{2\pi ij(iy_{d_2})}\big/ iI\!\!R & \longrightarrow & r_{S^1_{d_2}}\ e^{2\pi ij(y_{d_2})}\big/ I\!\!R
\end{array}$$

where:

- $S^1_{d_2}[j,m_j] = r_{S^1_{d_2}}[\cos(2\pi ijy_{d_2}) + i\sin(2\pi ijy_{d_2})]$,
- $S^1_{d_{2_\perp}}[j,m_j] = r_{S^1_{d_2}}[\cos(2\pi jy_{d_2}) + i\sin(2\pi jy_{d_2})]$,

Finally, recall that

$$e^{2\pi ijz} = e^{2\pi ijz_1} \times \cdots \times e^{2\pi ijz_d} \times \cdots \times e^{2\pi ijz_{2n}}$$

is a Laurent monomial in the variables $z_1, \cdots, z_{2n}$. ■

3.4.5 Proposition

The cohomology $H^{2i}(\overline{S}^{P_n}_{K_n}, \widehat{M}^{2i}_{T_{R_\oplus}} \otimes \widehat{M}^{2i}_{T_{L_\oplus}})$ *has the following analytic development:*

$$H^{2i}(\overline{S}^{P_n}_{K_n}, \widehat{M}^{2i}_{T_{R_\oplus}} \otimes \widehat{M}^{2i}_{T_{L_\oplus}}) = \sum_{j=1}^{r}\sum_{m_j}(\lambda^{\frac{1}{2}}(2i,j,m_j)e^{-2\pi ijz}) \otimes_{(D)} (\lambda^{\frac{1}{2}}(2i,j,m_j)e^{2\pi ijz})$$

where

$$\begin{array}{rl}
& \mathrm{EIS}_L(2i,j,m_j) = \sum\limits_{j=1}^{r}\sum\limits_{m_j}\lambda^{\frac{1}{2}}(2i,j,m_j)e^{2\pi ijz} \\
(\textit{resp.} & \mathrm{EIS}_R(2i,j,m_j) = \sum\limits_{j=1}^{r}\sum\limits_{m_j}\lambda^{\frac{1}{2}}(2i,j,m_j)e^{-2\pi ijz}\ )
\end{array}$$

is the (truncated) Fourier development of the equivalent of a normalized $2i$-dimensional left (resp. right) Eisenstein series of weight $k=2$ restricted to the upper (resp. lower) half space.

Proof:

1. First we recall that the Fourier development of a normalized Eisenstein series is:

$$G_k(y) \simeq \sum_{n=1}^{\infty} \sigma_{k-1}(n)\ q^n$$

with $y \in \mathbb{C}$, $q = e^{2\pi iy}$, $\sigma_{k-1}(n) = \sum\limits_{d|n} d^{k-1}$ and k the weight.

2. Referring to the development of the cohomology $H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_R}\otimes M^{2i}_{T_L})$ according to the sum over the products of equivalent representatives $g^{(2i)}_{T_{R\times L}}[j,m_j]=g^{(2i)}_{T_R}[j,m_j]\times g^{(2i)}_{T_L}[j,m_j]=T^{2i}_R[j,m_j]\times T^{2i}_L[j,m_j]$ of the complete bilinear semigroup $G^{(2i)}(F^T_{\overline{\omega}}\times F^+_{\omega})$ such that an i-dimensional left (resp. right) complex semitorus has the analytic development:

   $$\begin{aligned} T^{2i}_L[j,m_j] &\simeq \lambda^{\frac12}(2i,j,m_j)e^{2\pi ijz} \\ \text{(resp.}\quad T^{2i}_R[j,m_j] &\simeq \lambda^{\frac12}(2i,j,m_j)e^{-2\pi ijz}\ ), \end{aligned}$$

   we get the searched analytic decomposition of the cohomology in terms of the product $\mathrm{EIS}_R(2i,j,m_j)\times\mathrm{EIS}_L(2i,j,m_j)$ of the (truncated) Fourier development of the equivalent of a normalized $2i$-dimensional right Eisenstein series of weight two by its left equivalent.

   Remark that the sum $\sigma_{k-1}(n)=\sum\limits_{d|n}d^{k-1}$ in $G_k(y)$ corresponds in the $2i$-dimensional envisaged case to the sum $\sum\limits_{m_j}$ over the equivalent representatives m_j of the classes j . ■

3.4.6 Proposition

The analytic development $\mathrm{EIS}_R(2i,j,m_j)\otimes\mathrm{EIS}_L(2i,j,m_j)$ *of the bilinear cohomology* $H^{2i}(\overline{S}^{P_n}_{K_n},\widehat{M}^{2i}_{T_R}\otimes\widehat{M}^{2i}_{T_L})$ *is an eigenbifunction of the product of Hecke operators* $T_R(i;r)\otimes T_L(i;r)$ *whose j-th bicoset representative is*

$$U_{j_R}\times U_{j_L}=t^t_i(\mathcal{O}_{F_{\overline{\omega}_{j,m_j}}})\times t_i(\mathcal{O}_{F_{\omega_{j,m_j}}})\ .$$

Proof: the cohomology $H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_R}\otimes M^{2i}_{T_L})$ having coefficients in the $\mathrm{GL}_i(F^T_{\overline{\omega}}\times F^T_{\omega})$-bisemimodule $M^{2i}_{T_R}\otimes M^{2i}_{T_L}$ is in one-to-one correspondence with the functional representation space $\mathrm{FRepsp}(\mathrm{GL}_i(F^T_{\overline{\omega}}\times F^T_{\omega}))$ of the general bilinear semigroup $\mathrm{GL}_i(F^T_{\overline{\omega}}\times F^T_{\omega})$. Now, $\mathrm{GL}_i(F^T_{\overline{\omega}}\times F^T_{\omega})$ is a solvable bilinear semigroup such that we have a chain of embedded subsemigroups:

$$\begin{aligned} \mathrm{GL}_i(F^T_{\overline{\omega}_1}\times F^T_{\omega_1}) &\subset \mathrm{GL}_i(F^T_{\overline{\omega}_{1+2}}\times F^T_{\omega_{1+2}})\subset\cdots \\ &\qquad\subset \mathrm{GL}_i(F^T_{\overline{\omega}_{1+\cdots+j}}\times F^T_{\omega_{1+\cdots+j}}) \\ &\qquad\quad\subset \mathrm{GL}_i(F^T_{\overline{\omega}_{1+\cdots+j+\cdots+r}}\times F^T_{\omega_{1+\cdots+j+\cdots+r}}) \end{aligned}$$

corresponding to the sums of the bicosets of the quotient semigroup

$$\overline{X}_{S_{R\times L}}=\mathrm{GL}_i(F^T_R\times F^T_L)\Big/\mathrm{GL}_i((\mathbb{Z}/N\,\mathbb{Z})^2)=G^{(2i)}(F^T_{\overline{\omega}}\times F^T_{\omega})\ .$$

The conjugacy class representative $g^{(2i)}_{T_{R\times L}}[j,m_j]$ of $G^{(2i)}(F^T_{\overline{\omega}}\times F^T_\omega)$ is obtained throughout the action of the j-th bicoset representative $U_{j_R}\times U_{j_L}$ of the product of Hecke operators $T_R(i;r)\otimes T_L(i;r)$ according to proposition 2.2.5 since the Hecke bialgebra $\mathcal{H}_{R\times L}(i)$, generated by all the pseudo-ramified Hecke bioperators, has a representation in $\mathrm{GL}_i((\mathbb{Z}/N\,\mathbb{Z})^2)$.
According to propositions 3.4.1 and 3.4.5, 3.3.10 and 2.1.4, we have that:

$$\mathrm{FRepsp}(\mathrm{GL}_i(F^T_{\overline{\omega}}\times F^T_\omega))=G^{(2i)}(F^T_{\overline{\omega}_\oplus}\times F^T_{\omega_\oplus})=\mathrm{EIS}_R(2i,j,m_j)\otimes\mathrm{EIS}_L(2i,j,m_j)$$

for the upper value of $j=r\leq\infty$.
So, considering the quotient semigroup $\overline{X}_{S_{R\times L}}=\mathrm{GL}_n(F^T_R\times F^T_L)/\,\mathrm{GL}_n((\mathbb{Z}/N\ \mathbb{Z})^2)$ is equivalent to take into account the action of the Hecke bioperator $T_R(n;r)\otimes T_L(n;r)$ on $\mathrm{Repsp}(\mathrm{GL}_n(F^T_R\times F^T_L))$. It then results that $\mathrm{EIS}_R(2n,j,m_j)\otimes\mathrm{EIS}_L(2n,j,m_j)$ is an eigenbifunction of $T_R(n;r)\otimes T_L(n;r)$.
On the other hand, it is well-known that the Eisenstein series are eigenfunctions of the Hecke operators.
But, for more details on the algebraic spectral representations, see [Pie2] and [Clo]. ■

3.4.7 Corollary

The truncated (bi)series $\mathrm{EIS}_{R\times L}(2i,j,m_j)=\mathrm{EIS}_R(2i,j,m_j)\otimes\mathrm{EIS}_L(2i,j,m_j)$ *is:*

1. *a truncated supercuspidal (bi)form over* $\mathbb{C}^{\,i}$ *;*
2. *a solvable power (bi)series.*

Proof:

1. As $\mathrm{EIS}_L(2i,j,m_j)$ (resp. $\mathrm{EIS}_R(2i,j,m_j)$) constitutes the i-dimensional equivalent representative of the one-dimensional complex truncated Fourier development of a normalized Eisenstein series of weight two, which is a (quasi-) modular form, $\mathrm{EIS}_L(2i,j,m_j)$ (resp. $\mathrm{EIS}_R(2i,j,m_j)$) is a truncated (super)cuspidal form over $\mathbb{C}^{\,i}$.

2. As the general bilinear semigroup $\mathrm{GL}_i(F^T_{\overline{\omega}}\times F^T_\omega)$ is solvable and as we have that:

   $$\mathrm{FRepsp}(\mathrm{GL}_i(F^T_{\overline{\omega}_\oplus}\times F^T_{\omega_\oplus}))=\mathrm{EIS}_{R\times L}(2i,j,m_j)\ ,\quad F^T_{\omega_\oplus}=\bigoplus_j\bigoplus_{m_j}F^T_{\omega_{j,m_j}}\ ,$$

   according to propositions 3.4.6 and 2.1.4, the “Eisenstein” biseries $\mathrm{EIS}_{R\times L}(2i,j,m_j)$ will be said solvable.

   We thus have a chain of embedded (truncated) biseries:

$$\begin{aligned}\mathrm{EIS}_{R\times L}(2i,j^{\mathrm{up}}=1,m_j)\subset\cdots\subset\mathrm{EIS}_{R\times L}(2i,j^{\mathrm{up}}=j,m_j)\\ \subset\cdots\subset\mathrm{EIS}_{R\times L}(2i,j^{\mathrm{up}}=r,m_j)\end{aligned}$$

where j^{up} is the upper value of the integer " j " labeling the conjugacy classes.

This implies the injective mapping:

$$E_{j^{\mathrm{up}}}: \quad \mathrm{EIS}_{R\times L}(2i,j^{\mathrm{up}}=j,m_j) \longrightarrow \mathrm{EIS}_{R\times L}(2i,j^{\mathrm{up}}=j+1,m_j)$$

between two (truncated) biseries such that $\mathrm{EIS}_{R\times L}(2i,j^{\mathrm{up}}=j,m_j)$ is an eigenbifunction of the Hecke bioperator $T_R(i;j)\otimes T_L(i;j)$ while $\mathrm{EIS}_{R\times L}(2i,j^{\mathrm{up}}=j+1,m_j)$ is an eigenbifunction of $T_R(i;j+1)\otimes T_L(i;j+1)$. ■

3.4.8 Proposition

The cohomology $H^{2i}(\overline{S}^{\mathrm{Z}\,[P_n]}_{\mathrm{Z}\,[K_n]},\widehat{M}^{2i}_{T_{R_\oplus}}(D)\otimes\widehat{M}^{2i}_{T_{L_\oplus}}(D))$ *of the compactified diagonal bisemispace* $\overline{S}^{\mathrm{Z}\,[P_n]}_{\mathrm{Z}\,[K_n]}$ *has the analytic development:*

$$H^{2i}(\overline{S}^{\mathrm{Z}\,[P_n]}_{\mathrm{Z}\,[K_n]},\widehat{M}^{2i}_{T_{R_\oplus}}(D)\otimes\widehat{M}^{2i}_{T_{L_\oplus}}(D))=\sum_{j=1}^{r}(\lambda^{\frac12}(2i,j,1)\ e^{-2\pi ijz})\otimes(\lambda^{\frac12}(2i,j,1)\ e^{2\pi ijz})$$

where

$$\mathrm{EIS}_L(2i,j,0)=\sum_{j=1}^{r}\lambda^{\frac12}(2i,j,0)\ e^{2\pi ijz}$$

$$(\textit{resp.}\quad \mathrm{EIS}_R(2i,j,0)=\sum_{j=1}^{r}\lambda^{\frac12}(2i,j,0)\ e^{-2\pi ijz}\ )$$

is a solvable $2i$*-dimensional left (resp. right) cusp form of weight* $k=2$ *.*

Proof: this is a particular case of proposition 3.4.5 where the number of equivalent representatives of the left and right conjugacy classes is restricted to one. ■

3.4.9 Bialgebra of cusp forms $L^{1-1}_{\mathrm{cusp}}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega}))$

Let $L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}}\times F_{\omega}))$ be the bialgebra of complex-valued smooth continuous bifunctions $\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}})\otimes\phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}})$ on the pseudo-ramified bilinear semigroup $G^{(2n)}(F_{\overline{\omega}}\times F_{\omega})$ as developed in section 2.5.

Let

$$\gamma^C_{R\times L}: \quad G^{(2n)}(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_{\omega}) \longrightarrow G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega})$$

be the projective toroidal isomorphism of compactification of $G^{(2n)}(\widetilde{F}_{\overline{\omega}}\times \widetilde{F}_{\omega})$ mapping each conjugacy class representative $\tilde{g}^{(2n)}_{R\times L}[j,m_j]$ of $G^{(2n)}(\widetilde{F}_{\overline{\omega}}\times \widetilde{F}_{\omega})$ into its toroidal equivalent $g^{(2n)}_{T_{R\times L}}[j,m_j]$ according to proposition 3.3.3.
It is evident that $\gamma^{C}_{R\times L}$ generates $L^{1-1}_{\text{cusp}}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega}))$ in such a way that the bialgebra of smooth continuous bifunctions $\phi^{(2n)}_{G^T_{j_R}}(x_{g^T_{j_R}})\otimes\phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}})$ on the pseudo-ramified toroidal bilinear semigroup $G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega})$ is the bialgebra of cusp forms $L^{1-1}_{\text{cusp}}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega}))$.
On the other hand, the projective toroidal isomorphism maps each bifunction $\phi^{(2n)}_{G_{j_R}}(x_{g_{j_R}})\otimes \phi^{(2n)}_{G_{j_L}}(x_{g_{j_L}})\in L^{1-1}_{R\times L}(G^{(2n)}(F_{\overline{\omega}}\times F_{\omega}))$ on the conjugacy class representative $g^{(2n)}_{R\times L}[j,m_j]\in G^{(2n)}(F_{\overline{\omega}}\times F_{\omega})$ into its toroidal equivalent $\phi^{(2n)}_{G^T_{j_R}}(x_{g^T_{j_R}})\otimes\phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}})\in L^{1-1}_{\text{cusp}}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega}))$ on the conjugacy class representative $g^{(2n)}_{T_{R\times L}}[j,m_j]\in G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega})$ in such a way that:

1. $$\begin{aligned}\phi^{(2n)}_{G^T_{j_R}}(x_{g^T_{j_R}})\otimes_D\phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}}) &= T^{2n}_R[j,m_j]\otimes_D T^{2n}_L[j,m_j]\\ &= \lambda^{\frac12}(2n,j,m_j)\ e^{-2\pi ijz}\otimes_D\lambda^{\frac12}(2n,j,m_j)\ e^{2\pi ijz}\end{aligned}$$

   according to proposition 3.4.4.

2. $$\begin{aligned}\phi^{(2n)}_{G^T_{R}}(x_{g^T_{R}})\otimes_D\phi^{(2n)}_{G^T_{L}}(x_{g^T_{L}}) &= \mathop{\oplus}_{j=1}^{r}\mathop{\oplus}_{m_j}(\phi^{(2n)}_{G^T_{j_R}}(x_{g^T_{j_R}})\otimes_D\phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}}))\\ &= \mathop{\oplus}_{j=1}^{r}\mathop{\oplus}_{m_j}(\lambda^{\frac12}(2n,j,m_j)\ e^{-2\pi ijz}\otimes_D\lambda^{\frac12}(2n,j,m_j)\ e^{2\pi ijz}\\ &= \text{EIS}_R(2n,j,m_j)\otimes_{(D)}\text{EIS}_L(2n,j,m_j)\end{aligned}$$

   is the (truncated) supercuspidal biform on the pseudo-ramified toroidal bilinear semigroup $G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega})$ given by the product, right by left, of the equivalent of $2n$-dimensional Eisenstein series.

3.4.10 Proposition

Let $R_{G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega})}(\phi^{(2n)}_{G^T_R}\otimes\phi^{(2n)}_{G^T_L})$ denote the integral operator on cusp biforms $(\phi^{(2n)}_{G^T_R}\otimes\phi^{(2n)}_{G^T_L})$ over the pseudo-ramified toroidal bilinear semigroup $G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega})$.
Then, the trace formula of this integral operator is given by:

$$\text{tr}(R_{G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega})}(\phi^{(2n)}_{G^T_R}\otimes\phi^{(2n)}_{G^T_L})=\text{vol}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega}))\sum_j\sum_{m_j}\lambda(2n,j,m_j)$$

where $\lambda(2n,j,m_j)=\prod\limits_{d=1}^{2n}\lambda_d(2n,j,m_j)$ is the product of the eigenvalues $\lambda_d(n,j,m_j)$ of the j-th coset representative of the product, right by left, of Hecke operators.

Proof: Indeed, according to section 2.5.18, the trace of this integral operator can be developed as follows:

$$
\begin{aligned}
&\operatorname{tr}(R_{G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega})}(\phi^{(2n)}_{G^T_R}\otimes\phi^{(2n)}_{G^T_L})\\
&\quad=\bigoplus_{j,m_j}\operatorname{vol}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega}))\int_{G^{(2n)}}(\phi^{(2n)}_{G^T_{j_R}}(x_{g^T_{j_R}})\otimes_D\phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}})\;dx_{g^T_{j_R}}\;dx_{g^T_{j_L}}\\
&\quad=\bigoplus_{j,m_j}\operatorname{vol}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega}))\int_{G^{(2n)}}\lambda^{\frac12}(2n,j,m_j)\;e^{-2\pi ijz}\otimes_D\lambda^{\frac12}(2n,j,m_j)\;e^{2\pi ijz}\;dx_{g^T_{j_R}}\;dx_{g^T_{j_L}}\\
&\quad\simeq\operatorname{vol}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega}))(\mathrm{EIS}_R(2n,j,m_j),\mathrm{EIS}_L(2n,j,m_j))\\
&\quad=\operatorname{vol}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega}))\sum_{j,m_j}\lambda(2n,j,m_j)
\end{aligned}
$$

where $(\mathrm{EIS}_R(2n,j,m_j),\mathrm{EIS}_L(2n,j,m_j))$ is a bilinear form from $\operatorname{Irr}\operatorname{cusp}(\mathrm{GL}_n(F^T_{\overline{\omega}}\times F^T_{\omega}))=\mathrm{EIS}_{R\times L}(2n,j,m_j)$ to $\mathbb{C}$. ■

3.4.11 Proposition

Let $L^{1-1}_{R\times L}(P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1}))$ *be the bialgebra of complex-valued bifunctions on the bilinear parabolic semigroup* $P^{(2n)}(F_{\overline{\omega}^1}\times F_{\omega^1})$ *and let* $L^{1-1}_{R\times L}(P^{(2n)}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1}))$ *be its toroidal equivalent obtained by the projective toroidal isomorphism* $\gamma^C_{R\times L}$ *.*
Let $R_{P^{(2n)}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})}(\phi^{(2n)}_{P^T_R}\otimes\phi^{(2n)}_{P^T_L})$ *denote the integral operator on bifunctions* $\in L^{1-1}_{R\times L}(P^{(2n)}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1}))$ *.*
Then, the trace formula of this integral operator is given by:

$$
\begin{aligned}
&\operatorname{tr}(R_{P^{(2n)}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})}(\phi^{(2n)}_{P^T_R}\otimes\phi^{(2n)}_{P^T_L}))\\
&\quad=\operatorname{vol}(P^{(2n)}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})\sum_j\sum_{m_j}\int\chi^T_{j,m_{j_{R\times L}}}\;dx_{p_{j_R}}\;dx_{j_L}\\
&\quad=\operatorname{vol}(P^{(2n)}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1}))\sum_j\sum_{m_j}\lambda_I(2n,j,m_j)
\end{aligned}
$$

where

- $\chi^T_{j,m_{j_{R\times L}}}:x_{p_{j_R}}\times x_{p_{j_L}}\to\phi^{(2n)}_{P^T_{j_R}}(x_{p_{j_R}})\otimes_D\phi^{(2n)}_{P^T_{j_L}}(x_{p_{j_L}})$ *is a (bi)character on the irreducible equivalence class representative* $P^{(2n)}(F^T_{\overline{\omega}^1_{j,m_j}}\times F^T_{\omega^1_{j,m_j}})$ *;*
- $\lambda_I(2n,j,m_j)$ *is the "unitary" Hecke character restricted to* $P^{(2n)}(F^T_{\overline{\omega}^1_{j,m_j}}\times F^T_{\omega^1_{j,m_j}})$ *.*

Proof: This formula corresponds to the trace formula of proposition 2.5.13 submitted to the toroidal projective isomorphism $\gamma^C_{R\times L}$. ■

3.4.12 Proposition

Let

$$\operatorname{tr}(R_{G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega})}))(\phi^{(2n)}_{G^T_R}\otimes\phi^{(2n)}_{G^T_L})$$

$$=\operatorname{vol}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega})\sum_j\sum_{m_j}\lambda(2n,j,m_j)$$

be the trace of the integral operator on cusp biforms $(\phi^{(2n)}_{G^T_R}\otimes\phi^{(2n)}_{G^T_L})$ *over the pseudo-ramified toroidal bilinear semigroup* $G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega})$ *and let*

$$\operatorname{tr}(R_{P^{(2n)}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})}(\phi^{(2n)}_{P^T_R}\otimes\phi^{(2n)}_{P^T_L})$$

$$=\operatorname{vol}(P^{(2n)}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})\sum_j\sum_{m_j}\int\chi_{j,m_{j_{R\times L}}}\ dx_{p_{j_R}}\ dx_{p_{j_L}}$$

be the trace of the integral operator on bifunctions over the bilinear parabolic semigroup $P^{(2n)}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})$.
Then, the trace of the integral operator on pseudo-unramified cusp biforms $(\phi^{(2n)}_{G^{T,nr}_R}\otimes\phi^{(2n)}_{G^{T,nr}_L})$ *over the pseudo-unramified toroidal bilinear semigroup* $G^{(2n)}(F^{T,nr}_{\overline{\omega}_\oplus}\times F^{T,nr}_{\omega_\oplus})$ *is given by:*

$$\operatorname{tr}(R_{G^{(2n)}(F^{T,nr}_{\overline{\omega}}\times F^{T,nr}_{\omega})}))(\phi^{(2n)}_{G^{T,nr}_R}\otimes\phi^{(2n)}_{G^{T,nr}_L})$$

$$=\left(\operatorname{vol}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega})\Big/\operatorname{vol}(P^{(2n)}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})\right)\sum_j\sum_{m_j}\lambda(2n,j,m_j)\Big/\lambda_I(2n,j,m_j)\ .$$

Sketch of proof: This results from the action by convolution of the unitary operator $R(P^{(2n)}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1}))$ on the bifunctions of the pseudo-unramified bilinear semigroup $G^{(2n)}(F^{T,nr}_{\overline{\omega}}\times F^{T,nr}_{\omega}))$ as developed in proposition 2.5.12 and, more precisely, from the map:

$$G^{(2n)}_{F^T\to F^{T,nr}}:\quad G^{(2n)}(F^T_{\overline{\omega}}\times F^T_{\omega}))\longrightarrow G^{(2n)}(F^{T,nr}_{\overline{\omega}}\times F^{T,nr}_{\omega})$$

whose kernel is $\operatorname{Ker}(G^{(2n)}_{F^T\to F^{T,nr}})=P^{(2n)}(F^{T,nr}_{\overline{\omega}^1}\times F^{T,nr}_{\omega^1})$ as developed in section 2.4.1. ■

3.4.13 Proposition

Let

$$
\begin{aligned}
i_{G^T_{R\to L}}: \quad L^{1-1}_{\text{cusp}}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_\omega)) &\longrightarrow L^2_{\text{cusp}}(G^{(2n)}(F^T_\omega\times F^T_\omega))\;,\\
\phi^{(2n)}_{G^T_{j_R}}(x_{g^T_{j_R}})\otimes\phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}}) &\longrightarrow \phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}})\otimes\phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}})\;,
\end{aligned}
$$

be the involution map as introduced in section 2.5.4 and transforming the cusp biforms of $L^{1-1}_{\text{cusp}}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_\omega))$ *into the cusp biform* $\phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}})\otimes\phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}})$ *of* $L^2_{\text{cusp}}(G^{(2n)}(F^T_\omega\times F^T_\omega))$ *localized in the upper half space.*
Then, the trace formula

$$\operatorname{tr}(R_{G^{(2n)}(F^T_\omega\times F^T_\omega)}(\phi^{(2n)}_{G^T_L}\otimes\phi^{(2n)}_{G^T_L}))$$

of the integral operator on cusp biforms $(\phi^{(2n)}_{G^T_L}\otimes\phi^{(2n)}_{G^T_L})$ *leads to the Plancherel formula:*

$$\int_{G^{(2n)}}\left|\phi^{(2n)}_{G^T_L}(x_{g^T_L})\right|^2\;dx_{g^T_L}=\sum_j\sum_{m_j}\left|\widehat{\phi}^{(2n)}_{G^T_{j_L,m_{j_L}}}\right|^2$$

where

- $\widehat{\phi}^{(2n)}_{G^T_{j_L,m_{j_L}}}=\lambda^{\frac12}(2n,j,m_j)$ *can be considered as a global Hecke character;*
- *the sum over j and m_j runs over the conjugacy class representatives of the pseudo-ramified algebraic semigroup $G^{(2n)}(F^T_{\overline{\omega}}\times F^T_\omega)$ localized in the upper half space.*

Proof: According to proposition 3.4.10, we have from the following trace formula:

$$
\begin{aligned}
&\operatorname{tr}(R_{G^{(2n)}(F^T_{\overline{\omega}}\times F^T_\omega)}(\phi^{(2n)}_{G^T_L}\otimes\phi^{(2n)}_{G^T_L}))\\
&\quad=\bigoplus_{j,m_j}\operatorname{vol}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_\omega)\int_{G^{(2n)}}\left|(\phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}})\right|^2\;dx_{g^T_{j_L}}\\
&\quad=\bigoplus_{j,m_j}\operatorname{vol}(G^{(2n)}(F^T_{\overline{\omega}}\times F^T_\omega)\left|\lambda^{\frac12}(2n,j,m_j)\right|^2\;,
\end{aligned}
$$

the equality

$$\bigoplus_{j,m_j}\int_{G^{(2n)}}\left|(\phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}})\right|^2\;dx_{g^T_{j_L}}=\bigoplus_{j,m_j}\left|\lambda^{\frac12}(2n,j,m_j)\right|^2\;.$$

If we take into account that

1. $\int_{G^{(2n)}} \left|(\phi^{(2n)}_{G^T_L}(x_{g^T_L})\right|^2 \ dx_{g^T_L} = \underset{j,m_j}{\oplus} \int_{G^{(2n)}} \left|(\phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}})\right|^2 \ dx_{g^T_{j_L}}$ expressing the development of the integral of the square of the cusp biform $\phi^{(2n)}_{G^T_L}(x_{g^T_L})$ according to the squares of the cusp subbiforms $\phi^{(2n)}_{G^T_{j_L}}(x_{g^T_{j_L}})$ on the conjugacy class representatives $g^{(2n)}_{T_L}[,m_j]$ of the pseudo-ramified bilinear semigroup $G^{(2n)}(F^T_{\overline{\omega}} \times F^T_{\omega})$;

2. $$\begin{aligned} \widehat{\phi}^{(2n)}_{G^T_{j_L,m_{j_L}}} &= \int_{G^{(2n)}} \phi^{(2n)}_{G^T_L}(x_{g^T_L})\ e^{-2\pi ijz}\ dx_{g^T_L} \\ &= \lambda^{\frac{1}{2}}(2n,j,m_j)\ , \end{aligned}$$

   we get the searched Plancherel formula
$$\int_{G^{(2n)}} \left|(\phi^{(2n)}_{G^T_L}(x_{g^T_L})\right|^2 \ dx_{g^T_L} = \sum_j \sum_{m_j} \left|\widehat{\phi}^{(2n)}_{G^T_{j_L,m_{j_L}}}\right|^2 . \qquad \blacksquare$$

We can then consider the following Langlands global correspondence.

3.4.14 Proposition

On the compactified lattice bisemispace
$$\overline{X}_{S_{R\times L}} = \mathrm{GL}_n(F^T_R \times F^T_L))\Big/ \mathrm{GL}_n((\mathbb{Z}/N\ \mathbb{Z})^2)\ ,$$
there exists the **global Langlands correspondence***:*

$$\begin{tikzcd}
\mathrm{Irr\,Rep}^{(2i)}_{W_{F_{R\times L}}}(W^{ab}_{F_R} \times W^{ab}_{F_L}) \arrow[r] \arrow[d, equal] & \mathrm{Irr\,cusp}(\mathrm{GL}_i(F^T_{\overline{\omega}} \times F^T_{\omega})) \arrow[d, equal] \\
G^{(2i)}(\widetilde{F}_{\overline{\omega}_\oplus} \times \widetilde{F}_{\omega_\oplus}) \arrow[r] \arrow[d, "\wr"] & \mathrm{EIS}_{R\times L}(2i,j,m_j) \\
G^{(2i)}(F^T_{\overline{\omega}_\oplus} \times F^T_{\omega_\oplus}) \arrow[r] \arrow[d, equal] & \widehat{G}^{(2i)}(F^T_{\overline{\omega}} \times F^T_{\omega}) \arrow[u] \\
H^{2i}(\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{R_\oplus}} \otimes M^{2i}_{T_{L_\oplus}}) &
\end{tikzcd}$$

- *from the sum of the products, right by left, of the equivalence classes of the irreducible $2i$-dimensional Frobenius semisimple Weil-Deligne representation* $\mathrm{Irr}\,\mathrm{Rep}^{(2i)}_{W_{F_{R\times L}}}(W^{ab}_{F_R}\times W^{ab}_{F_L})$ *of the bilinear global Weil group* $(W^{ab}_{F_R}\times W^{ab}_{F_L})$ *given by the algebraic bilinear semigroup* $G^{(2i)}(\widetilde{F}_{\overline{\omega}_\oplus}\times \widetilde{F}_{\omega_\oplus}))$
- *to the sum of the products, right by left, of the equivalence classes of the irreducible cuspidal representation* $\mathrm{Irr}\,\mathrm{cusp}(\mathrm{GL}_i(F^T_{\overline{\omega}}\times F^T_{\omega}))$ *of* $\mathrm{GL}_i(F^T_{\overline{\omega}}\times F^T_{\omega})$ *given by $2i$-dimensional cusp biforms* $\mathrm{EIS}_{R\times L}(2i,j,m_j)$ *, in such a way that* $\mathrm{Irr}\,\mathrm{cusp}(\mathrm{GL}_i(F^T_{\overline{\omega}}\times F^T_{\omega}))=\mathrm{Irr}\,\mathrm{cusp}(\mathrm{GL}_i(F^T_{\overline{\omega}_\oplus}\times F^T_{\omega_\oplus}))$.

Proof: we start with $\mathrm{Irr}\,\mathrm{Rep}^{(2i)}_{W_{F_{R\times L}}}(W^{ab}_{F_R}\times W^{ab}_{F_L})$ which was proved to be equal to the algebraic bilinear semigroup $G^{(2i)}(\widetilde{F}_{\overline{\omega}_\oplus}\times \widetilde{F}_{\omega_\oplus})$ in proposition 3.3.10. Then, a toroidal compactification maps $G^{(2i)}(\widetilde{F}_{\overline{\omega}_\oplus}\times \widetilde{F}_{\omega_\oplus})$ into its compactified toroidal equivalent $G^{(2i)}(F^T_{\overline{\omega}_\oplus}\times F^T_{\omega_\oplus})$ according to sections 3.3.2 to 3.3.3. The analytic development of the compactified bilinear semigroup $G^{(2i)}(F^T_{\overline{\omega}_\oplus}\times F^T_{\omega_\oplus})$ is given by the product of the (truncated) Fourier development of a normalized $2i$-dimensional cusp form by its left equivalent: $\mathrm{EIS}_{R\times L}(2i,j,m_j)$. So, we get the searched bijection:

$$\mathrm{Irr}\,\mathrm{Rep}^{(2i)}_{W_{F_{R\times L}}}(W^{ab}_{F_R}\times W^{ab}_{F_L})\simeq \mathrm{Irr}\,\mathrm{cusp}(\mathrm{GL}_i(F^T_{\overline{\omega}}\times F^T_{\omega}))\ . \qquad \blacksquare$$

3.5 Langlands global correspondence on the boundary $\partial\overline{X}^{(\mathbb{R})}_{S_{R\times L}}$ of the irreducible compactified lattice bisemispace $\overline{X}_{S_{R\times L}}$

3.5.1 Definition: the boundary of the Borel-Serre compactification

Let $\gamma^c_{R\times L}: X_{S_{R\times L}}\to \overline{X}_{S_{R\times L}}$ be the projective emergent isomorphism, as introduced in proposition 3.3.2, which maps the lattice bisemispace $X_{S_{R\times L}}$ into its compactified toroidal correspondent $\overline{X}_{S_{R\times L}}$ such that $X_{S_{R\times L}}$ can be considered as the interior of $\overline{X}_{S_{R\times L}}$ in the context of the Borel-Serre compactification.
Then, the boundary $\partial\overline{X}_{S_{R\times L}}$ of $\overline{X}_{S_{R\times L}}$ will be defined as resulting from the inclusion morphism:

$$\gamma^\delta_{R\times L}:\quad \overline{X}_{S_{R\times L}}\hookrightarrow\partial\overline{X}_{S_{R\times L}}$$

sending $\overline{X}_{S_{R\times L}}=\mathrm{GL}_n(F^T_R\times F^T_L)/\mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)=GL_n(F^T_{\overline{\omega}}\times F^T_{\omega})$ to its real boundary $\partial\overline{X}^{(\mathbb{R})}_{S_{R\times L}}=\mathrm{GL}_n(F^{+,T}_R\times F^{+,T}_L)/\mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)=GL_n(F^{+,T}_{\overline{v}}\times F^{+,T}_v)$, where $F^{+,T}_v=\{F^{+,T}_{v_{1_\delta}},\cdots,F^{+,T}_{v_{j_\delta},m_{j_\delta}},\cdots,F^{+,T}_{v_{r_\delta}}\}$ covers its complex equivalent F^T_ω .

By this way, each complex diagonal conjugacy class representative $g^{(2n)}_{T_L}[j,m_j=0]$ (resp. $g^{(2n)}_{T_R}[j,m_j=0]$) of $\overline{X}_{S_{R\times L}}$ is covered by the set of m_{j_δ} real conjugacy class representatives $\{g^{(n)}_{T_L}[j_\delta,m_{j_\delta}]\}_{m_{j_\delta}}$ (resp. $\{g^{(n)}_{T_R}[j_\delta,m_{j_\delta}]\}_{m_{j_\delta}}$) of $\partial\overline{X}^{(I\!R)}_{S_{R\times L}}$.
The morphism $\gamma^\delta_{R\times L}: \quad \overline{X}_{S_{R\times L}}\to\partial\overline{X}_{S_{R\times L}}$ is such that there is a one-to-one correspondence between the left (resp. right) points $P^T_{a_{L[j,m_j]}}$ (resp. $P^T_{a_{R[j,m_j]}}$) of $\partial\overline{X}_{S_L}$ (resp. $\partial\overline{X}_{S_R}$) and their "real" equivalents on $\partial\overline{X}^{(I\!R)}_{S_L}$ (resp. $\partial\overline{X}^{(I\!R)}_{S_R}$).

3.5.2 Definition: the double coset decomposition of the equivalent of the Shimura bisemivariety $\partial\overline{S}^{P_n}_{K_n}$

The inclusion morphism $\gamma^\delta_{R\times L}$ has for result that the double coset decomposition

$$\overline{S}^{P_n}_{K_n}=P_n(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})\setminus\mathrm{GL}_n(F^T_R\times F^T_L)\big/\mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)$$

of $\mathrm{GL}_n(F^T_R\times F^T_L)$ corresponding to $\overline{X}_{S_{R\times L}}$ is transformed into the following double coset decomposition

$$\partial\overline{S}^{P_n}_{K_n}=P_n(F^{+,T}_{\overline{v}^1}\times F^{+,T}_{v^1})\setminus\mathrm{GL}_n(F^{+,T}_R\times F^{+,T}_L)\Big/\mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)$$

where $P_n(F^{+,T}_{v^1})$ (resp. $P_n(F^{+,T}_{\overline{v}^1})$) is the standard left (resp. right) parabolic subgroup over the set of real irreducible completions $F^{+,T}_{v^1_{j_\delta,m_{j_\delta}}}$ (resp. $F^{+,T}_{\overline{v}^1_{j_\delta,m_{j_\delta}}}$) (see section 2.4.1).
$\partial\overline{S}^{P_n}_{K_n}$ is the equivalent of a Shimura (bisemi)variety because [Del2], [Har1]:

- it is dependent on the morphism $\gamma^\delta_{R\times L}$ sending a "complex" bisemivariety $\overline{X}^{(I\!R)}_{S_{R\times L}}$ into a "real" bisemivariety $\partial\overline{X}^{(I\!R)}_{S_{R\times L}}$;
- it is defined with respect to an open compact subgroup $P_n(F^{+,T}_{\overline{v}^1}\times F^{+,T}_{v^1})$.

3.5.3 Proposition

Let $M^{2i}_{T_{\overline{v}_R}}\otimes M^{2i}_{T_{v_L}}$ *be a* $(B^{2i}_{\overline{v}^T}\otimes B^{2i}_{v^T})$*-bisemimodule where* $B^{2i}_{\overline{v}^T}\otimes B^{2i}_{v^T}\simeq T_i(F^{+,T}_{\overline{v}})^t\times T_i(F^{+,T}_v)$ *is a tensor product of division semialgebras and let* $M^{2i}_{T_{\overline{v}_{R_\oplus}}}\otimes M^{2i}_{T_{v_{L_\oplus}}}$ *be a* $\mathrm{GL}_i(F^{+,T}_{\overline{v}_\oplus}\times F^{+,T}_{v_\oplus})$*-bisemimodule where* $F^{+,T}_{v_\oplus}$ *is given by*

$$F^{+,T}_{v_\oplus}=\bigoplus_{j_\delta}F^{+,T}_{v_{j_\delta}}\bigoplus_{j_\delta,m_{j_\delta}}F^{+,T}_{v_{j_\delta,m_{j_\delta}}}\ .$$

Then, the bilinear cohomology of the equivalent of the Shimura bisemivariety $\partial\overline{S}^{P_n}_{K_n}$ *is the Eisenstein cohomology:*

$$H^{2i}(\partial\overline{S}^{P_n}_{K_n},M^{2i}_{T_{\overline{v}_{R_\oplus}}}\otimes M^{2i}_{T_{v_{L_\oplus}}})=G^{(2i)}(F^{+,T}_{\overline{v}_\oplus}\times F^{+,T}_{v_\oplus})$$

in bijection with the $2i$-dimensional real irreducible representation $\operatorname{Irr}\operatorname{Rep}^{(2i)}_{W_{F^+_{R\times L}}}(W^{ab}_{F^+_{\overline{v}}}\times W^{ab}_{F^+_v})$ *of the product, right by left, of global Weil groups.*

Proof:

- This proposition is a transposition of proposition 3.3.10 to $H^{2i}(\partial\overline{S}^{P_n}_{K_n},\cdot)$.
- The fact that $H^{2i}(\partial\overline{S}^{P_n}_{K_n},M^{2i}_{T_{\overline{v}_{R_\oplus}}}\otimes M^{2i}_{T_{v_{L_\oplus}}})$ is the Eisenstein (bilinear) cohomology results from the works of G. Harder and J. Shimura [Har2], [Sch].
- the global Weil groups $W^{ab}_{F^+_v}$ and $W^{ab}_{F^+_{\overline{v}}}$ referring to finite real algebraic extensions can be defined as they were introduced for complex algebraic extensions in definition 1.1.9.

Then, referring to proposition 3.3.10, we can state that:

$$\operatorname{Irr}\operatorname{Rep}^{(2i)}_{W_{F^+_{R\times L}}}(W^{ab}_{F^+_{\overline{v}}}\times W^{ab}_{F^+_v})=G^{(2i)}(\widetilde{F}^+_{\overline{v}_\oplus}\times\widetilde{F}^+_{v_\oplus})\qquad\blacksquare$$

3.5.4 Proposition

The cohomology $H^{2i}(\partial\overline{S}^{P_n}_{K_n},M^{2i}_{T_{\overline{v}_{R_\oplus}}}\otimes M^{2i}_{T_{v_{L_\oplus}}})$ *has a decomposition according to the equivalent representatives* $g^{(2i)}_{T_{R\times L}}[j_\delta,m_{j_\delta}]$ *of the conjugacy classes of the real bilinear semigroup* $G^{(2i)}(F^{+,T}_{\overline{v}}\times F^{+,T}_v)$ *according to:*

$$H^{2i}(\partial\overline{S}^{P_n}_{K_n},M^{2i}_{T_{\overline{v}_{R_\oplus}}}\otimes M^{2i}_{T_{v_{L_\oplus}}})=\bigoplus_{j_\delta}\bigoplus_{m_{j_\delta}}g^{(i)}_{T_{R\times L}}[j_\delta,m_{j_\delta}]\ .$$

Proof: this is an adaptation of the real case of proposition 3.4.1 since

$$H^{2i}(\partial\overline{S}^{P_n}_{K_n},M^{2i}_{T_{\overline{v}_{R_\oplus}}}\otimes M^{2i}_{T_{v_{L_\oplus}}})=G^{(2i)}(F^{+,T}_{\overline{v}}\times F^{+,T}_v)\ .\qquad\blacksquare$$

3.5.5 Proposition

The Eisenstein cohomology $H^{2i}(\partial\overline{S}^{P_n}_{K_n},\widehat{M}^{2i}_{T_{\overline{v}_{R_\oplus}}}\otimes\widehat{M}^{2i}_{T_{v_{L_\oplus}}})$ *has a power series representation given by the product of a right $2i$-dimensional global elliptic semimodule* $\mathrm{ELLIP}_R(2i,j_\delta,m_{j_\delta})$ *by its left equivalent* $\mathrm{ELLIP}_L(2i,j_\delta,m_{j_\delta})$ *according to:*

$$H^{2i}(\partial\overline{S}^{P_n}_{K_n},\widehat{M}^{2i}_{T_{\overline{v}_{R_\oplus}}}\otimes\widehat{M}^{2i}_{T_{v_{L_\oplus}}})=\mathrm{ELLIP}_R(2i,j_\delta,m_{j_\delta})\otimes_{(D)}\mathrm{ELLIP}_L(2i,j_\delta,m_{j_\delta})$$

where

$$\begin{aligned}\mathrm{ELLIP}_L(2i,j_\delta,m_{j_\delta}) &= \bigoplus_{j_\delta=1}^{t}\bigoplus_{m_{j_\delta}} \lambda^{\frac12}(i,j_\delta,m_{j_\delta})e^{2\pi i j_\delta x}\ ,\\ \mathrm{ELLIP}_R(2i,j_\delta,m_{j_\delta}) &= \bigoplus_{j_\delta=1}^{t}\bigoplus_{m_{j_\delta}} \lambda^{\frac12}(i,j_\delta,m_{j_\delta})e^{-2\pi i j_\delta x}\ ,\end{aligned}$$

with

- $\vec{x}=\sum\limits_{c=1}^{i} x_c\vec{e}_c$ *a vector of* $(F^+_{v_{j_\delta}})^i$ *and more precisely a point of* $G^{(i)}(F^+_{v_{j_\delta}}) \equiv g_L^{(i)}[j_\delta,mj_\delta]$ *and* $x=\sum\limits_{c=1}^{i} x_c\ |\vec{e}_c|$ *;*

- $\lambda(i,j_\delta,m_{j_\delta})=\prod\limits_{c=1}^{i}\lambda_c(i,j_\delta,m_{j_\delta})$ *a product of eigenvalues* $\lambda_c(i,j_\delta,m_{j_\delta})$ *of* $g_i(\mathcal{O}_{F^{+,T}_{\overline{v}_{j_\delta},m_{j_\delta}}} \times \mathcal{O}_{F^{+,T}_{v_{j_\delta},m_{j_\delta}}}) \in \mathrm{GL}_i((\mathbb{Z}/N\,\mathbb{Z})^2)$ *.*

Proof: since $g^{(i)}_{T_L}[j_\delta,m_{j_\delta}]$ (resp. $g^{(i)}_{T_R}[j_\delta,m_{j_\delta}]$) is a i-dimensional left (resp. right) real semitorus, it has the following analytical development:

$$\begin{aligned} T^i_L[j_\delta,m_{j_\delta}] &= \lambda^{\frac12}(i,j_\delta,m_{j_\delta})e^{2\pi i j_\delta x}\\ (\text{resp.}\quad T^i_R[j_\delta,m_{j_\delta}] &= \lambda^{\frac12}(i,j_\delta,m_{j_\delta})e^{-2\pi i j_\delta x}\ )\end{aligned}$$

where

$$\lambda(i,j_\delta,m_{j_\delta})=\prod_{c=1}^{i}\lambda_c(i,j_\delta,m_{j_\delta})=\det(\alpha_{i^2;(j_\delta)^2}\times D_{(j_\delta)^2;m^2_{j_\delta}})_{ss}\simeq j^{2i}\cdot(N)^{2i}$$

is a global Hecke character since the $\lambda_c(i,j_\delta,m_{j_\delta})$ are eigenvalues of $g_i(\mathcal{O}_{F^{+,T}_{\overline{v}_{j_\delta},m_{j_\delta}}}\times\mathcal{O}_{F^{+,T}_{v_{j_\delta},m_{j_\delta}}})$ which is the j_δ-th coset representative of the product of Hecke operators (see propositions 3.4.4 and 3.4.5).
$\lambda^{\frac12}(i,j_\delta,m_{j_\delta})$ has an inflation action on $e^{2\pi i j_\delta x}$ leading to the existence of a global elliptic **semimodule** ELLIP$(\boldsymbol{.},\boldsymbol{.},\boldsymbol{.})$.
On the other hand, $g^{(i)}_{T_L}[j_\delta,m_{j_\delta}]$ (resp. $g^{(i)}_{T_R}[j_\delta,m_{j_\delta}]$) is defined on the real left (resp. right) place v_{j_δ} (resp. $\overline{v}_{j_\delta}$) implying the global Frobenius substitution:

$$\begin{aligned} e^{2\pi i x} &\longrightarrow e^{2\pi i j_\delta x}\\ (\text{resp.}\quad e^{-2\pi i x} &\longrightarrow e^{-2\pi i j_\delta x}\ ).\end{aligned}$$

Note that these $2i$-dimensional left and right global elliptic semimodules ELLIP$_L(2i,j_\delta,m_{j_\delta})$ and ELLIP$_R(2i,j_\delta,m_{j_\delta})$ are the $2i$-dimensional equivalents of the 2-dimensional global elliptic semimodules introduced in [Pie3] (see also [Ande]). ■

3.5.6 Characters and extended bicharacters

The functions $x_c \to e^{2\pi i j_\delta x_c}$ from the real line, i.e. the irreducible central real completion $F^+_{v^1_{1_\delta}}$, to the complex numbers of modulus one, are characters if they satisfy the identity $\chi(x_c y_c) = \chi(x_c)\cdot\chi(y_c)$. In the bilinear context considered here, the characters are replaced by extended bicharacters defined as follows:

If $x_c^+ \cdot x_c^- \equiv -(x_c)^2$ denote a bipoint of the real irreducible completion $F^+_{\overline{v}^1_{1_\delta}} \times F^+_{v^1_{1_\delta}}$, i.e. the product of a point $x_c^- \equiv x_c$ by its symmetrical $x_c^+ \equiv -x_c$, $x_c \in F^+_{v^1_{1_\delta}}$, then an extended bicharacter $\chi_+(x_c^+)\cdot\chi_-(x_c^-)$ is a continuous bifunction

$$\begin{aligned}\chi_+\cdot\chi_- :\ & x_c^+\cdot x_c^- \\ & \longrightarrow \lambda^{\frac{1}{2}}(1,j_\delta,m_{j_\delta})\cdot e^{-2\pi i j_\delta x_c} \times \lambda^{\frac{1}{2}}(1,j_\delta,m_{j_\delta})\cdot e^{2\pi i j_\delta x_c} = \lambda(1,j_\delta,m_{j_\delta})\end{aligned}$$

from $F^+_{\overline{v}^1_{1_\delta}} \times F^+_{v^1_{1_\delta}}$ to the product of the complex numbers $\lambda^{\frac{1}{2}}(1,j_\delta,m_{j_\delta})\cdot e^{2\pi i j_\delta x_c}$ of modulus $\neq 1$ by their complex conjugates where $\lambda(1,j_\delta,m_{j_\delta})$ is generally an eigenvalue of a product $U_{j_{\delta_R}} \times U_{j_{\delta_L}}$ of Hecke operators.

3.5.7 Proposition

The $2i$-dimensional global elliptic bisemimodule $\mathrm{ELLIP}_R(2i,j_\delta,m_{j_\delta})\otimes_{(D)}\mathrm{ELLIP}_L(2i,j_\delta,m_{j_\delta})$ *is:*

1. *a solvable power biseries, constituting an irreducible (super)cuspidal representation of the (truncated) (super)cuspidal biform* $\mathrm{EIS}_{R\times L}(n,j,m_j=0)$ *;*
2. *an eigenbifunction of the Hecke bioperator* $T_R(2i;t)\otimes T_L(2i;t)$ *.*

Proof:

1. The fact that $\mathrm{ELLIP}_R(2i,j_\delta,m_{j_\delta})\otimes\mathrm{ELLIP}_L(2i,j_\delta,m_{j_\delta})$ is a solvable power biseries results from:

   $$\mathrm{Repsp}(\mathrm{GL}_i(F^{+,T}_{\overline{v}_\oplus}\times F^{+,T}_{v_\oplus})) = \mathrm{ELLIP}_{R\times L}(2i,j_\delta,m_{j_\delta})$$

   where $\mathrm{ELLIP}_{R\times L}\equiv\mathrm{ELLIP}_R\otimes\mathrm{ELLIP}_L$ (see proposition 3.4.7).

   $\mathrm{ELLIP}_R(2i,j_\delta,m_{j_\delta})\otimes\mathrm{ELLIP}_L(2i,j_\delta,m_{j_\delta})$ is an irreducible supercuspidal representation of $\mathrm{EIS}_{R\times L}(2i,j,m_j=0)$ due to the morphism $\gamma^\delta_{R\times L}:\overline{X}_{S_{R\times L}}\to\partial\overline{X}_{S_{R\times L}}$ related to the generation of the boundary of the Borel-Serre compactification according to definition 3.5.1.

2. The proof that $\text{ELLIP}_{R\times L}(2i,j_\delta,m_{j_\delta})$ is an eigenbifunction of the Hecke bioperator $T_R(2i;t)\otimes T_L(2i;t)$ can be handled as in proposition 3.4.6.

   Let us yet note that the j_δ-th bicoset representative of $T_R(2i;j_\delta)\otimes T_L(2i;j_\delta)$ is given, with evident notations, by :

   $$U_{j_{\delta_R}}\times U_{j_{\delta_L}}=t_i^t(\mathcal{O}_{F^{+,T}_{\overline{v}_{j_\delta,m_{j_\delta}}}})\times t_i(\mathcal{O}_{F^{+,T}_{v_{j_\delta,m_{j_\delta}}}})\ . \qquad \blacksquare$$

3.5.8 Bialgebra of elliptic semimodules $\boldsymbol{L^{1-1}_{\text{ELLIP}}(G^{(2i)}(F^{+,T}_{\overline{v}}\times F^{+,T}_{v}))}$

As in section 3.4.9, we can introduce a bialgebra $L^{1-1}_{\text{ELLIP}}(G^{(2i)}(F^{+,T}_{\overline{v}}\times F^{+,T}_{v}))$ of complex-valued smooth continuous bifunctions $\phi^{(i)}_{G^T_{j^\delta_R}}(x_{g^T_{j^\delta_R}})\otimes\phi^{(i)}_{G^T_{j^\delta_L}}(x_{g^T_{j^\delta_L}})$ on the pseudo-ramified toroidal bilinear semigroup $G^{(2i)}(F^{+,T}_{\overline{v}}\times F^{+,T}_{v})$ from the corresponding bialgebra $L^{1-1}_{R\times L}(G^{(2i)}(F^{+}_{\overline{v}}\times F^{+}_{v}))$ of smooth continuous bifunctions on the pseudo-ramified bilinear semigroup $G^{(2i)}(F^{+}_{\overline{v}}\times F^{+}_{v})$ under the action of a projective toroidal isomorphism.
Then, each bifunction on the conjugacy class representative $g^{(i)}_{T_{R\times L}}[j_\delta,m_{j_\delta}]\in G^{(i)}(F^{+,T}_{\overline{v}}\times F^{+,T}_{v})$ is given, according to proposition 3.4.5, by

$$\begin{aligned}\phi^{(i)}_{G^T_{j^\delta_R}}(x_{g^T_{j^\delta_R}})\otimes_D\phi^{(i)}_{G^T_{j^\delta_L}}(x_{g^T_{j^\delta_L}})&=T^i_R[j_\delta,m_{j_\delta}]\otimes_D T^i_L[j^\delta,m_{j^\delta}]\\&=\lambda^{\frac12}(i,j_\delta,m_{j_\delta})\;e^{-2\pi ij_\delta x}\otimes_{(D)}\lambda^{\frac12}(i,j_\delta,m_{j_\delta})\;e^{2\pi ij_\delta x}\end{aligned}$$

and the sum of all bifunctions on the conjugacy class representatives of $G^{(2i)}(F^{+,T}_{\overline{v}}\times F^{+,T}_{v})$:

$$\begin{aligned}\phi^{(2i)}_{G^T_{\delta_R}}(x_{g^T_R})\otimes_D\phi^{(2i)}_{G^T_{\delta_L}}(x_{g^T_L})&=\bigoplus_{j_\delta=1}^{t}\bigoplus_{m_{j_\delta}}\left(\phi^{(i)}_{G^T_{j^\delta_R}}(x_{g^T_{j^\delta_R}})\otimes_D\phi^{(i)}_{G^T_{j^\delta_L}}(x_{g^T_{j^\delta_L}})\right)\\&=\bigoplus_{j_\delta}\bigoplus_{m_{j_\delta}}(\lambda^{\frac12}(i,j_\delta,m_{j_\delta})\;e^{-2\pi ij_\delta x}\otimes_{(D)}\lambda^{\frac12}(i,j_\delta,m_{j_\delta})\;e^{2\pi ij_\delta x})\\&=\text{ELLIP}_R(2i,j_\delta,m_{j_\delta})\otimes_{(D)}\text{ELLIP}_L(2i,j_\delta,m_{j_\delta})\end{aligned}$$

is the (diagonal) product of a right $2i$-dimensional global elliptic semimodule $\text{ELLIP}_R(2i,j_\delta,m_{j_\delta})$ by its left equivalent $\text{ELLIP}_L(2i,j_\delta,m_{j_\delta})$.

3.5.9 Proposition

Let $R_{G^{(2i)}(F^{+,T}_{\overline{v}}\times F^{+,T}_{v})}(\phi^{(2i)}_{G^T_{\delta_R}}\otimes\phi^{(2i)}_{G^T_{\delta_L}})$ *be the integral operator on elliptic bisemimodules* $(\phi^{(2i)}_{G^T_{\delta_R}}\otimes\phi^{(2i)}_{G^T_{\delta_L}})$ *over the pseudo-ramified toroidal bilinear real semigroup* $G^{(2i)}(F^{+,T}_{\overline{v}_\oplus}\times F^{+,T}_{v_\oplus})$.

Then, the trace formula of this integral operator is given by:

$$\begin{aligned}
&\mathrm{tr}(R_{G^{(2i)}(F^{+,T}_{\overline{v}}\times F^{+,T}_{v})}(\phi^{(2i)}_{G^T_{\delta_R}} \otimes \phi^{(2i)}_{G^T_{\delta_L}})\\
&\quad = \mathrm{vol}(G^{(2i)}(F^{+,T}_{\overline{v}} \times F^{+,T}_{v}))\sum_{j^\delta}\sum_{m_{j^\delta}} \lambda(i, j_\delta, m_{j_\delta})\ .
\end{aligned}$$

Sketch of proof: This is an adaptation of the trace formula $\mathrm{tr}(R_{G^{(2i)}(F^T_{\overline{\omega}}\times F^T_{\omega})}(\phi^{(2i)}_{G^T_R}\otimes\phi^{(2i)}_{G^T_L})$ developed in proposition 3.4.10 to the real case. ■

3.5.10 Proposition

On the boundary $\partial\overline{X}^{(\mathbb{R})}_{S_{R\times L}} = \mathrm{GL}_i(F^{+,T}_R \times F^{+,T}_L)/\,\mathrm{GL}_i((\mathbb{Z}\,/N\,\mathbb{Z}\,)^2)$ *of the compactified lattice bisemispace* $\overline{X}_{S_{R\times L}}$ *, there exists the Langlands global correspondence:*

$$\begin{tikzcd}
\mathrm{Irr}\,\mathrm{Rep}^{(2i)}_{W^+_{F_{R\times L}}}(W^{ab}_{F^+_{\overline{v}}} \times W^{ab}_{F^+_{v}}) \arrow[r] \arrow[d, equal] & \mathrm{Irr}\,\mathrm{ELLIP}(\mathrm{GL}_i(F^{+,T}_{\overline{v}} \times F^{+,T}_{v})) \arrow[d, equal] \\
G^{(2i)}(\widetilde{F}^{+}_{\overline{v}_\oplus} \times \widetilde{F}^{+}_{v_\oplus}) \arrow[r] \arrow[d] & \mathrm{ELLIP}_{R\times L}(2i, j_\delta, m_{j_\delta}) \\
G^{(2i)}(F^{+,T}_{\overline{v}_\oplus} \times F^{+,T}_{v_\oplus}) \arrow[r] \arrow[d, equal] & \widehat{G}^{(2i)}(F^{+,T}_{\overline{v}} \times F^{+,T}_{v}) \arrow[u] \\
H^{2i}(\partial\overline{S}^{P_n}_{K_n}, M^{2i}_{T_{\overline{v}_{R_\oplus}}} \otimes M^{2i}_{T_{v_{L_\oplus}}}) &
\end{tikzcd}$$

- *from the sum of products, right by left, of the equivalence classes of the irreducible* $2i$*-dimensional Frobenius semisimple Weil-Deligne representation* $\mathrm{Irr}\,\mathrm{Rep}^{(2i)}_{W^+_{F_{R\times L}}}(W^{ab}_{F^+_{\overline{v}}} \times W^{ab}_{F^+_{v}})$ *of the bilinear global Weil group* $(W^{ab}_{F^+_{\overline{v}}} \times W^{ab}_{F^+_{v}})$ *given by the algebraic bilinear real semigroup* $G^{(2i)}(\widetilde{F}^{+}_{\overline{v}_\oplus} \times \widetilde{F}^{+}_{v_\oplus}))$
- *to the sum of the products, right by left, of the equivalence classes of the irreducible elliptic representation*$\mathrm{Irr}\,\mathrm{ELLIP}(\mathrm{GL}_i(F^{+,T}_{\overline{v}} \times F^{+,T}_{v}))$ *of* $\mathrm{GL}_i(F^{+,T}_{\overline{v}} \times F^{+,T}_{v})$ *given by the*

$2i$-dimensional solvable global elliptic bisemimodule $\mathrm{ELLIP}_{R\times L}(2i,j_\delta,m_{j_\delta})$ *in such a way that* $\mathrm{Irr}\,\mathrm{ELLIP}(\mathrm{GL}_i(F^{+,T}_{\overline{v}}\times F^{+,T}_{v}))=\mathrm{ELLIP}_{R\times L}(2i,j_\delta,m_{j_\delta})$.

3.5.11. The n-dimensional irreducible global Langlands correspondences can be summarized in the following diagram:

$$\begin{tikzcd}
\mathrm{Irr}\,\mathrm{Rep}^{(2n)}_{W_{F_{R\times L}}}(W^{ab}_{F_R}\times W^{ab}_{F_L})) \arrow[r] \arrow[d] & \mathrm{Irr}\,\mathrm{cusp}(\mathrm{GL}_n(F^T_{\overline{\omega}}\times F^T_{\omega})) \arrow[d] \\
\mathrm{Irr}\,\mathrm{Rep}^{(2n)}_{W_{F^+_{R\times L}}}(W^{ab}_{F^+_{\overline{v}}}\times W^{ab}_{F^+_{v}})) \arrow[r] & \mathrm{Irr}\,\mathrm{ELLIP}(\mathrm{GL}_n(F^{+,T}_{\overline{v}}\times F^{+,T}_{v}))
\end{tikzcd}$$

4 Langlands global correspondences for reducible representations of $\mathrm{GL}((2)n)$

4.1 Reducibility of $\mathrm{GL}_{(2)n}(F_{\overline{\omega}}\times F_{\omega})$

4.1.1 Definition: Partial reducibility of $\mathrm{GL}_n(F_{\overline{\omega}}\times F_{\omega})$

Another way of constructing admissible representations of the general bilinear semigroup $\mathrm{GL}_n(F_{\overline{\omega}}\times F_{\omega})$ is to consider a partition $n=n_1+n_2+\cdots+n_\ell+\cdots+n_s$ of the integer n , $n_\ell\in\mathbb{N}$, $n_\ell<n\in\mathbb{N}$, leading to the representations $\mathrm{Rep}(\mathrm{GL}_{n_\ell}(F_{\overline{\omega}}\times F_{\omega}))$ of $\mathrm{GL}_{n_\ell}(F_{\overline{\omega}}\times F_{\omega})$, for $1\le\ell\le s$, in such a way that $\mathrm{Rep}(\mathrm{GL}_{n_\ell}(F_{\overline{\omega}}\times F_{\omega}))$ constitutes an irreducible representation of $\mathrm{GL}_{n_\ell}(F_{\overline{\omega}}\times F_{\omega})$.

The tensor product

$$\mathrm{Rep}(\mathrm{GL}_{n_1}(F_{\overline{\omega}}\times F_{\omega}))\otimes\cdots\otimes\mathrm{Rep}(\mathrm{GL}_{n_\ell}(F_{\overline{\omega}}\times F_{\omega}))\otimes\cdots\otimes\mathrm{Rep}(\mathrm{GL}_{n_s}(F_{\overline{\omega}}\times F_{\omega}))$$

may not be irreducible but it has an irreducible quotient given by the formal sum:

$$\begin{aligned}&\mathrm{Rep}(\mathrm{GL}_{n=n_1+\cdots+n_s}(F_{\overline{\omega}}\times F_{\omega}))\\&\quad=\mathrm{Rep}(\mathrm{GL}_{n_1}(F_{\overline{\omega}}\times F_{\omega}))\boxplus\cdots\\&\qquad\boxplus\mathrm{Rep}(\mathrm{GL}_{n_\ell}(F_{\overline{\omega}}\times F_{\omega}))\boxplus\cdots\boxplus\mathrm{Rep}(\mathrm{GL}_{n_s}(F_{\overline{\omega}}\times F_{\omega}))\end{aligned}$$

which constitutes a partially reducible representation of $\mathrm{GL}_n(F_{\overline{\omega}}\times F_{\omega})$.

4.1.2 Definition: complete reducibility of $\mathrm{Rep}(\mathrm{GL}_{2n}(F_{\overline{\omega}}\times F_{\omega}))$

If the partition $2n=2_1+2_2+\cdots+2_\ell+\cdots+2_n$ of $2n$, $2_\ell=|2|$, is considered, then the tensor product

$$\mathrm{Rep}(\mathrm{GL}_{2_1}(F_{\overline{\omega}}\times F_{\omega}))\otimes\cdots\otimes\mathrm{Rep}(\mathrm{GL}_{2_n}(F_{\overline{\omega}}\times F_{\omega}))$$

of representations of $\mathrm{GL}_{2_\ell}(F_{\overline{\omega}}\times F_{\omega})$, $1\le\ell\le n$, must be taken into account. This tensor product has an irreducible quotient given by the formal sum:

$$\begin{aligned}&\mathrm{Rep}(\mathrm{GL}_{2n=2_1+\cdots+2_\ell+\cdots+2_n}(F_{\overline{\omega}}\times F_{\omega}))\\&\quad=\mathrm{Rep}(\mathrm{GL}_{2_1}(F_{\overline{\omega}}\times F_{\omega}))\boxplus\cdots\\&\qquad\boxplus\mathrm{Rep}(\mathrm{GL}_{2_\ell}(F_{\overline{\omega}}\times F_{\omega}))\boxplus\cdots\boxplus\mathrm{Rep}(\mathrm{GL}_{2_n}(F_{\overline{\omega}}\times F_{\omega}))\end{aligned}$$

which constitutes a completely reducible representation of $\mathrm{GL}_{2n}(F_{\overline{\omega}}\times F_{\omega}))$ decomposing into the direct sum of irreducible representations $\mathrm{Rep}(\mathrm{GL}_{2_\ell}(F_{\overline{\omega}}\times F_{\omega}))$, $1\le\ell\le n$.

4.1.3 Proposition

Let $2n_L = 2_{1_L} + 2_{2_L} + \cdots + 2_{k_L} + \cdots + 2_{\ell_L} + \cdots + 2_{n_L}$ *(resp.* $2n_R = 2_{1_R} + 2_{2_R} + \cdots + 2_{k_R} + \cdots + 2_{\ell_R} + \cdots + 2_{n_R}$ *) be a partition of* $2n_L$ *(resp.* $2n_R$ *) labeling the irreducible representations of* $T_{2n_L}(F_\omega)$ *(resp.* $T^t_{2n_R}(F_{\overline{\varpi}})$ *). Then:*

1. $\mathrm{Rep}(\mathrm{GL}_{2n=2_1+\cdots+2_\ell+\cdots+2_n}(F_{\overline{\varpi}}\times F_\omega)) = \boxplus_{\ell=1}^{n} \mathrm{Rep}(\mathrm{GL}_{2_\ell}(F_{\overline{\varpi}}\times F_\omega))$ *constitutes a completely reducible orthogonal bilinear representation of* $\mathrm{GL}_{2n}(F_{\overline{\varpi}}\times F_\omega)$ *.*

2. $\mathrm{Rep}(\mathrm{GL}_{2n_{R\times L}}(F_{\overline{\varpi}} \times F_\omega)) = \boxplus_{2_{\ell_R}\equiv 2_{\ell_L}=2}^{2n} \mathrm{Rep}(\mathrm{GL}_{2_{\ell_{R\times L}}}(F_{\overline{\varpi}} \times F_\omega)) \boxplus_{2_{k_R}\neq 2_{\ell_L}} \mathrm{Rep}(T^t_{2_{k_R}}(F_{\overline{\varpi}})\times T_{2_{\ell_L}}(F_\omega))$ *, where* $\mathrm{GL}_{2_{\ell_{R\times L}}}$ *is another notation for* GL_{2_ℓ} *, constitutes a completely reducible nonorthogonal bilinear representation of* $\mathrm{GL}_{2n}(F_{\overline{\varpi}}\times F_\omega))$ *.*

Proof: if we consider the decomposition of $\mathrm{GL}_{2n}(F_{\overline{\varpi}}\times F_\omega)$ into a product of trigonal semigroups according to:

$$\mathrm{GL}_{2n}(F_{\overline{\varpi}}\times F_\omega) = T^t_{2n_R}(F_{\overline{\varpi}})\times T_{2n_L}(F_\omega)$$

and the partitions of $2n_L$ and $2n_R$ as envisaged in this proposition, we have that:

$$\begin{aligned}
&\mathrm{Rep}(\mathrm{GL}_{2n_{R\times L}}(F_{\overline{\varpi}}\times F_\omega))\\
&= \left(\boxplus_{2_{\ell_R}=2}^{2n} \mathrm{Rep}(T^t_{2_{\ell_R}}(F_{\overline{\varpi}}))\right) \otimes \left(\boxplus_{2_{\ell_L}=2}^{2n} \mathrm{Rep}(T_{2_{\ell_L}}(F_\omega))\right)\\
&= \boxplus_{2_{\ell_R}\equiv 2_{\ell_L}=2}^{2n} \mathrm{Rep}(\mathrm{GL}_{2_\ell}(F_{\overline{\varpi}}\times F_\omega)) \boxplus_{2_{k_R}\neq 2_{\ell_L}} \mathrm{Rep}(T^t_{2_{k_R}}(F_{\overline{\varpi}})\times T_{2_{\ell_L}}(F_\omega))
\end{aligned}$$

where $\mathrm{GL}_{2_\ell}(F_{\overline{\varpi}}\times F_\omega) = T^t_{2_{\ell_R}}(F_{\overline{\varpi}})\times T_{2_{\ell_L}}(F_\omega)$.

$\mathrm{Rep}(\mathrm{GL}_{2n_{R\times L}}(F_{\overline{\varpi}}\times F_\omega))$ then decomposes diagonally according to the direct sum of irreducible representations $\mathrm{Rep}(\mathrm{GL}_{2_\ell}(F_{\overline{\varpi}}\times F_\omega))$ leading to a completely reducible orthogonal bilinear representation of $\mathrm{Rep}(\mathrm{GL}_{2n}(F_{\overline{\varpi}}\times F_\omega))$ if $\boxplus_{2_{k_R}\neq 2_{\ell_L}} \mathrm{Rep}(T^t_{2_{k_R}}(F_{\overline{\varpi}})\times T_{2_{\ell_L}}(F_\omega)) = 0$. ■

4.1.4 Corollary

1. *The representation of the general bilinear semigroup* $\mathrm{GL}_n(F_{\overline{\varpi}}\times F_\omega)$ *is* **partially reducible** *if it decomposes according to the direct sum of irreducible bilinear representations* $\mathrm{Rep}(\mathrm{GL}_{n_\ell}(F_{\overline{\varpi}}\times F_\omega))$ *.*

2. *The representation of the general bilinear semigroup* $\mathrm{GL}_{2n}(F_{\overline{\varpi}}\times F_\omega)$ *is* **orthogonally completely reducible** *if it decomposes diagonally according to the direct sum of irreducible bilinear representations* $\mathrm{Rep}(\mathrm{GL}_{2_\ell}(F_{\overline{\varpi}}\times F_\omega))$ *.*

3. *The representation of the general bilinear semigroup* $\mathrm{GL}_{2n}(F_{\overline{\omega}}\times F_\omega))$ *is* **nonorthogonally completely reducible** *if it decomposes diagonally according to the direct sum of irreducible bilinear representations* $\mathrm{Rep}(\mathrm{GL}_{2_\ell}(F_{\overline{\omega}}\times F_\omega))$ *and off diagonally according to the direct sum of irreducible bilinear representations* $\mathrm{Rep}(T^t_{2_{k_R}}(F_{\overline{\omega}})\times T_{2_{\ell_L}}(F_\omega))$ *.*

4.1.5 Lemma

Let $X^{2n}_{S_{R\times L}}=\mathrm{GL}_n(\widetilde F_R\times\widetilde F_L))\big/\mathrm{GL}_n((\mathbb{Z}/N\,\mathbb{Z})^2)$ *denote the lattice bisemispace as introduced in definition 2.3.1, and let*

$$\begin{aligned}X^{2n=2n_1+\cdots+2n_s}_{S_{R\times L}}&=\mathrm{GL}_{n=n_1+\cdots+n_s}(\widetilde F_R\times\widetilde F_L)\Big/\mathrm{GL}_{n=n_1+\cdots+n_s}((\mathbb{Z}/N\,\mathbb{Z})^2)\\&=\mathrm{Repsp}(\mathrm{GL}_{n=n_1+\cdots+n_s}(\widetilde F_{\overline{\omega}}\times\widetilde F_\omega)\end{aligned}$$

be the partially reducible lattice bisemispace.
Then, we have that the bilinear cohomology of $X^{2n=2n_1+\cdots+2n_s}_{S_{R\times L}}$ *decomposes according to:*

$$H^*(X^{2n=2n_1+\cdots+2n_s}_{S_{R\times L}})=\bigoplus_{2n_\ell=2n_1}^{2n_s}H^{2n_\ell}(X^{2n}_{S_{R\times L}},\widetilde M^{2n_\ell}_R\otimes\widetilde M^{2n_\ell}_L)$$

where $\widetilde M^{2n_\ell}_L$ *(resp.* $\widetilde M^{2n_\ell}_R$*)) is the left (resp. right)* $T_{n_\ell}(\widetilde F_\omega)$*-subsemimodule* $M^{2n_\ell}_L$ *(resp.* $T^t_{n_\ell}(\widetilde F_{\overline{\omega}})$*-subsemimodule* $M^{2n_\ell}_R$ *) of complex dimension* n_ℓ *.*

Proof: this results from the partial reducibility of $X^{2n=2n_1+\cdots+2n_s}_{S_{R\times L}}$ and from the equality:

$$H^{2n_\ell}(X^{2n}_{S_{R\times L}},\widetilde M^{2n_\ell}_R\otimes\widetilde M^{2n_\ell}_L)=\mathrm{Repsp}(\mathrm{GL}_{n_\ell}(\widetilde F_{\overline{\omega}}\times\widetilde F_\omega))$$

if we refer to section 3.2. ■

4.1.6 Proposition

Let

$$X^{2n=2_1+\cdots+2_n}_{S_{R\times L}}=\mathrm{GL}_{2n=2_1+\cdots+2_n}(\widetilde F_{\overline{\omega}}\times\widetilde F_\omega)\Big/\mathrm{GL}_{2n=2_1+\cdots+2_n}((\mathbb{Z}/N\,\mathbb{Z})^2)$$

be the orthogonal completely reducible lattice bisemispace and let $X^{2n_R\times 2n_L}_{S_{R\times L}}$ *be its nonorthogonal correspondent. Then, we have the following decompositions of the bilinear cohomologies:*

a) $H^{2n}(X^{2n}_{S_{R\times L}},\widetilde M^{2n}_R\otimes\widetilde M^{2n}_L)=\bigoplus_{2_\ell}H^2(X^{2n}_{S_{R\times L}},\widetilde M^{2_{\ell_R}}_R\otimes\widetilde M^{2_{\ell_L}}_L)$ *;*

b) $H^{2n}(X^{2n_{R\times L}}_{S_{R\times L}},\widetilde{M}^{2n_R}_R\otimes\widetilde{M}^{2n_L}_L)=\underset{2_\ell}{\oplus}H^2(X^{2_{n_{R\times L}}}_{S_{R\times L}},\widetilde{M}^{2_{\ell_R}}_R\otimes\widetilde{M}^{2_{\ell_L}}_L)$

$$\underset{2_{k_R},2_{\ell_L}}{\oplus}H^2(X^{2_{k_R-\ell_L}}_{S_{R\times L}},\widetilde{M}^{2_{k_R}}_R\otimes\widetilde{M}^{2_{\ell_L}}_L)$$

where

- $\widetilde{M}^{2n}_R\otimes\widetilde{M}^{2n}_L=\underset{2_{\ell_R}=2_{\ell_L}}{\oplus}(\widetilde{M}^{2_{\ell_R}}_R\otimes\widetilde{M}^{2_{\ell_L}}_L)$ *is the completely reducible orthogonal* $\mathrm{GL}_{2n=2_1+\cdots+2_n}(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_\omega)$*-bisemimodule* $\widetilde{M}^{2n}_R\otimes\widetilde{M}^{2n}_L$ *;*
- $\widetilde{M}^{2n_R}_R\otimes\widetilde{2}M^{n_L}_L=\underset{2_{\ell_R}=2_{\ell_L}}{\oplus}(\widetilde{M}^{2_{\ell_R}}_R\otimes\widetilde{M}^{2_{\ell_L}}_L)\underset{2_{k_R}\neq2_{\ell_L}}{\oplus}(\widetilde{M}^{2_{k_R}}_R\otimes\widetilde{M}^{2_{\ell_L}}_L)$ *is the completely reducible nonorthogonal* $\mathrm{GL}_{2n_{R\times L}}(\widetilde{F}_{\overline{\omega}}\times\widetilde{F}_\omega)$*-bisemimodule* $\widetilde{M}^{2n_R}_R\otimes\widetilde{M}^{2n_L}_L$ *.* ■

4.1.7 Proposition

Let $\mathrm{CY}^{2n_\ell}(X_R)$ *(resp.* $\mathrm{CY}^{2n_\ell}(X_L)$ *) be an algebraic semicycle of complex codimension* n_ℓ *over a right (resp. left) semischeme* X_R *(resp.* X_L *) isomorphic to the pseudo-ramified lattice semispace* $X^{2n}_{S_R}$ *(resp.* $X^{2n}_{S_L}$ *).*
Let

$$\begin{aligned}\mathrm{CY}^{2n_\ell}(X_R)\times\mathrm{CY}^{2n_\ell}(X_L)&=\underset{j}{\oplus}\underset{m_j}{\oplus}(\mathrm{CY}^{2n_\ell}(X_R[j,m_j])\times\mathrm{CY}^{2n_\ell}(X_L[j,m_j]))\\&\simeq\mathrm{Repsp}(\mathrm{GL}_{n_\ell}(\widetilde{F}_{\overline{\omega}_\oplus}\times\widetilde{F}_{\omega_\oplus}))\end{aligned}$$

be the product, right by left, of these algebraic semicycles decomposed according to their conjugacy classes (see section 1.3) in such a way that $\mathrm{CY}^{2n_\ell}(X_R)\times\mathrm{CY}^{2n_\ell}(X_L)$ *is isomorphic to an algebraic bilinear subsemigroup* $G^{(2n_\ell)}(\widetilde{F}_{\overline{\omega}_\oplus}\times\widetilde{F}_{\omega_\oplus})$ *.*
Then, we have the following decompositions of the bilinear cohomologies:

- $$\begin{aligned}H^*(X^{2n=2n_1+\cdots+2n_s}_{S_{R\times L}},\widetilde{M}^{2n}_{R_\oplus}\otimes\widetilde{M}^{2n}_{L_\oplus})&=\overset{2n_s}{\underset{2n_\ell=n_1}{\oplus}}H^{2n_\ell}(X^{2n}_{S_{R\times L}},\widetilde{M}^{2n_\ell}_{R_\oplus}\otimes\widetilde{M}^{2n_\ell}_{L_\oplus})\\&\simeq\underset{n_\ell}{\oplus}\underset{j}{\oplus}\underset{m_j}{\oplus}(\mathrm{CY}^{2n_\ell}(X_R[j,m_j])\times\mathrm{CY}^{2n_\ell}(X_L[j,m_j]))\end{aligned}$$
- $$\begin{aligned}H^{2n}(X^{2n}_{S_{R\times L}},\widetilde{M}^{2n}_{R_\oplus}\otimes\widetilde{M}^{2n}_{L_\oplus})&=\underset{2_\ell}{\oplus}H^2(X^{2n}_{S_{R\times L}},\widetilde{M}^{2_{\ell_R}}_{R_\oplus}\otimes M^{2_{\ell_L}}_{L_\oplus})\\&\simeq\underset{2_\ell}{\oplus}\underset{j}{\oplus}\underset{m_j}{\oplus}(\mathrm{CY}^{2_\ell}(X_R[j,m_j])\times\mathrm{CY}^{2_\ell}(X_L[j,m_j]))\end{aligned}$$
- $$\begin{aligned}H^{2n}(X^{2n_{R\times L}}_{S_{R\times L}},\widetilde{M}^{2n_R}_{R_\oplus}\otimes\widetilde{M}^{2n_L}_{L_\oplus})&=\underset{2_\ell}{\oplus}H^2(X^{2n_{R\times L}}_{S_{R\times L}},\widetilde{M}^{2_{\ell_R}}_{R_\oplus}\otimes\widetilde{M}^{2_{\ell_L}}_{L_\oplus})\\&\quad\underset{2_{k_R}\neq2_{\ell_L}}{\oplus}H^2(X^{2n_{R\times L}}_{S_{R\times L}},\widetilde{M}^{2_{k_R}}_{R_\oplus}\otimes\widetilde{M}^{2_{\ell_L}}_{L_\oplus})\end{aligned}$$

$$\begin{aligned}\simeq \bigoplus_{2_{\ell_{R\times L}}} \bigoplus_j \bigoplus_{m_j} & (\mathrm{CY}^{2_\ell}(X_R[j,m_j]) \times \mathrm{CY}^{2_\ell}(X_L[j,m_j])) \\ \bigoplus_{2_{k_R}\neq 2_{\ell_L}} \bigoplus_{j_R} \bigoplus_{m_{j_R}} \bigoplus_{j_L} \bigoplus_{m_{j_L}} & (\mathrm{CY}^{2_{k_R}}(X_R[j_R,m_{j_R}]) \times \mathrm{CY}^{2_{\ell_L}}(X_L[j_L,m_{j_L}]))\end{aligned}$$

being equal to the sums of products, right by left, of the conjugacy classes, counted with their multiplicities, of the irreducible semicycles.

Sketch of proof: indeed, $H^{2n_\ell}(X^{2n}_{S_{R\times L}}, \widetilde{M}^{2n_\ell}_{R_\oplus} \otimes \widetilde{M}^{2n_\ell}_{L_\oplus}) = \mathrm{Repsp}(\mathrm{GL}_{n_\ell}(\widetilde{F}_{\overline{\omega}_\oplus} \times \widetilde{F}_{\omega_\oplus})) = \bigoplus_j \bigoplus_{m_j} (\widetilde{M}^{2n_\ell}_{\overline{\omega}_{j,m_j}} \otimes \widetilde{M}^{2n_\ell}_{\omega_{j,m_j}})$ where $\widetilde{M}^{2n_\ell}_{\overline{\omega}_{j,m_j}}$ (resp. $\widetilde{M}^{2n_\ell}_{\omega_{j,m_j}}$) is the conjugacy class representative $g_R^{2n_\ell}[j,m_j]$ (resp. $g_L^{2n_\ell}[j,m_j]$) $\in G^{2n_\ell}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$. ■

4.2 Langlands global correspondences on the reducible compactified bisemispaces $\overline{X}^{(2)n}_{S_{R\times L}}$ and $\partial\overline{X}^{(2)n}_{S_{R\times L}}$

4.2.1 Definition: reducible compactified-lattice bisemispaces

Let

$$X^{2n=2n_1+\cdots+2n_s}_{S_{R\times L}}\ , \quad X^{2n=2_1+\cdots+2_n}_{S_{R\times L}} \quad \text{and} \quad X^{2n_R\times 2n_L}_{S_{R\times L}}$$

be respectively the partially reducible, orthogonal completely reducible and nonorthogonal completely reducible lattice bisemispaces as introduced in lemma 4.1.5 and proposition 4.1.6. Then, we define the projective emergent isomorphisms of toroidal compactifications:

$$\begin{aligned}\gamma^{c(2n)}_{R\times L} : \quad & X^{2n=2n_1+\cdots+2n_s}_{S_{R\times L}} \\ & \longrightarrow \overline{X}^{2n=2n_1+\cdots+2n_s}_{S_{R\times L}} = \mathrm{GL}_{n=n_1+\cdots+n_s}(F^T_R \times F^T_L) \Big/ \mathrm{GL}_{n=n_1+\cdots+n_s}((\mathbb{Z}/N\,\mathbb{Z})^2)\ ,\end{aligned}$$

$$\begin{aligned}\gamma^{c(2n)}_{R\times L} : \quad & X^{2n=2_1+\cdots+2_n}_{S_{R\times L}} \\ & \longrightarrow \overline{X}^{2n=2_1+\cdots+2_n}_{S_{R\times L}} = \mathrm{GL}_{2n=2_1+\cdots+2_n}(F^T_R \times F^T_L) \Big/ \mathrm{GL}_{2n=2_1+\cdots+2_n}((\mathbb{Z}/N\,\mathbb{Z})^2)\ ,\end{aligned}$$

$$\begin{aligned}\gamma^{c(2n_R\times 2n_L)}_{R\times L} : \quad & X^{2n_R\times 2n_L}_{S_{R\times L}} \\ & \longrightarrow \overline{X}^{2n_R\times 2n_L}_{S_{R\times L}} = \mathrm{GL}_{2n_R\times 2n_L}(F^T_R \times F^T_L) \Big/ \mathrm{GL}_{2n_R\times 2n_L}((\mathbb{Z}/N\,\mathbb{Z})^2)\ ,\end{aligned}$$

mapping these reducible lattice bisemispaces into their toroidal compactified equivalents in the sense of proposition 3.3.3 with the evident notations:

$$\overline{X}^{n=n_1+\cdots+n_s}_{S_{R\times L}} = \boxplus_{n_\ell=n_1}^{n_s} \overline{X}^{n_\ell}_{S_{R\times L}}\ .$$

4.2.2 Definition

The double coset decomposition of the reducible bilinear general semigroups are given by:

$$\begin{aligned}
&\overline{S}^{P_{n=n_1+\cdots+n_s}}_{K_{n=n_1+\cdots+n_s}}\\
&\quad = P_{n=n_1+\cdots+n_s}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})\setminus \mathrm{GL}_{n=n_1+\cdots+n_s}(F^T_R\times F^T_L)\Big/ \mathrm{GL}_{n=n_1+\cdots+n_s}((\mathbb{Z}/N\,\mathbb{Z})^2)\;,\\[2ex]
&\overline{S}^{P_{2n=2_1+\cdots+2_n}}_{K_{2n=2_1+\cdots+2_n}}\\
&\quad = P_{2n=2_1+\cdots+2_n}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})\setminus \mathrm{GL}_{2n=2_1+\cdots+2_n}(F^T_R\times F^T_L)\Big/ \mathrm{GL}_{2n=2_1+\cdots+2_n}((\mathbb{Z}/N\,\mathbb{Z})^2)\;,\\[2ex]
&\overline{S}^{P_{2n_R\times 2n_L}}_{K_{2n_R\times 2n_L}}\\
&\quad = P_{2n_R\times 2n_L}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})\setminus \mathrm{GL}_{2n_R\times 2n_L}(F^T_R\times F^T_L)\Big/ \mathrm{GL}_{2n_R\times 2n_L}((\mathbb{Z}/N\,\mathbb{Z})^2)\;,
\end{aligned}$$

with the notations of definition 3.3.4.

4.2.3 Definition: induction

If the double coset decompositions of the reducible bilinear general semigroups are considered, there are evident inductions respectively from the representations of the complex bilinear reducible parabolic subgroups:

$$P_{n=n_1+\cdots+n_s}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})\;,\quad P_{2n=2_1+\cdots+2_n}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})\quad\text{and}\quad P_{2n_R\times 2n_L}(F^T_{\overline{\omega}^1}\times F^T_{\omega^1})$$

and from the respective representations of the bilinear reducible arithmetic subgroups:

$$\mathrm{GL}_{n=n_1+\cdots+n_s}((\mathbb{Z}/N\,\mathbb{Z})^2)\;,\quad \mathrm{GL}_{2n=2_1+\cdots+2_n}((\mathbb{Z}/N\,\mathbb{Z})^2)\quad\text{and}\quad \mathrm{GL}_{2n_R\times 2n_L}((\mathbb{Z}/N\,\mathbb{Z})^2)$$

to the respective representations of the reducible bilinear general semigroups:

$$\mathrm{GL}_{n=n_1+\cdots+n_s}\;,\quad \mathrm{GL}_{2n=2_1+\cdots+2_n}\quad\text{and}\quad \mathrm{GL}_{2n_R\times 2n_L}\;.$$

4.2.4 Proposition

Let

$$\begin{aligned}
\gamma^{c(2n)}_{R\times L}:\quad &\mathrm{CY}^{2n=2n_1+\cdots+2n_s}(X_R)\times \mathrm{CY}^{2n=2n_1+\cdots+2n_s}(X_L)=\bigoplus_{2n_\ell=2n_1}^{2n_s}\mathrm{CY}^{2n_\ell}(X_R)\times \mathrm{CY}^{2n_\ell}(X_L)\\
&\longrightarrow\;\mathrm{CY}^{2n}_T(\overline{X}_R)\times \mathrm{CY}^{2n}_T(\overline{X}_L)=\bigoplus_{n_\ell}\mathrm{CY}^{2n_\ell}_T(\overline{X}_R)\times \mathrm{CY}^{2n_\ell}_T(\overline{X}_L)
\end{aligned}$$

be the projective emergent isomorphism of compactifications applied to a product, right by left, of $2n$*-dimensional partially reducible semicycles such that* $\mathrm{CY}_T^{2n_\ell}(\overline{X}_L)$ *(resp.* $\mathrm{CY}_T^{2n_\ell}(\overline{X}_R)$ *) be a* n_ℓ*-dimensional complex semitorus* $T_L^{2n_\ell}$ *(resp.* $T_R^{2n_\ell}$ *).*
Similarly, let

$$\begin{aligned}\gamma_{R\times L}^{c(2n_{R\times L})} : \quad & CY^{2n=2_1+\cdots+2_n}(X_R)\times \mathrm{CY}^{2n=2_1+\cdots+2_n}(X_L)\\ & \qquad\longrightarrow \mathrm{CY}_T^{2n=2_1+\cdots+2_n}(\overline{X}_R)\times \mathrm{CY}_T^{2n=2_1+\cdots+2_n}(\overline{X}_L)\end{aligned}$$

and

$$\begin{aligned}\gamma_{R\times L}^{c(2n_R\times 2n_L)} : \quad & CY^{2n_R}(X_R)\times \mathrm{CY}^{2n_L}(X_L)\\ & \qquad\longrightarrow \mathrm{CY}_T^{2n_R}(\overline{X}_R)\times \mathrm{CY}_T^{2n_L}(\overline{X}_L)\end{aligned}$$

be the two other considered projective emergent isomorphisms applied to products of completely reducible semicycles.
Then, we have the evident decompositions of the cohomologies:

- $$\begin{aligned} H^*(\overline{S}_{K_{n=n_1+\cdots+n_s}}^{P_{n=n_1+\cdots+n_s}}, \widehat{M}_{T_{R_\oplus}}^{2n}\otimes \widehat{M}_{T_{L_\oplus}}^{2n}) &= \bigoplus_{2n_\ell} H^{2n_\ell}(\overline{S}_{K_n}^{P_n}, \widehat{M}_{T_{R_\oplus}}^{2n_\ell}\otimes \widehat{M}_{T_{L_\oplus}}^{2n_\ell})\\ &\simeq \bigoplus_{n_\ell}\bigoplus_{j}\bigoplus_{m_j}(\mathrm{CY}_T^{2n_\ell}(\overline{X}_R[j,m_j])\times \mathrm{CY}_T^{2n_\ell}(\overline{X}_L[j,m_j]))\end{aligned}$$

- $$\begin{aligned} H^{2n}(\overline{S}_{K_{2n=2_1+\cdots+2_n}}^{P_{2n=2_1+\cdots+2_n}}, \widehat{M}_{T_{R_\oplus}}^{2n}\otimes \widehat{M}_{T_{L_\oplus}}^{2n}) &= \bigoplus_{2\ell_R=2\ell_L} H^{2}(\overline{S}_{K_{2n}}^{P_{2n}}, \widehat{M}_{T_{R_\oplus}}^{2\ell_R}\otimes \widehat{M}_{T_{L_\oplus}}^{2\ell_L})\\ &\simeq \bigoplus_{2\ell_R=2\ell_L}\bigoplus_{j}\bigoplus_{m_j}(\mathrm{CY}_T^{2\ell_R}(\overline{X}_R[j,m_j])\times \mathrm{CY}_T^{2\ell_L}(\overline{X}_L[j,m_j]))\end{aligned}$$

- $$\begin{aligned} H^{2n}(\overline{S}_{K_{2n_R\times 2n_L}}^{P_{2n_R\times 2n_L}}, \widehat{M}_{T_{R_\oplus}}^{2n_R}\otimes \widehat{M}_{T_{L_\oplus}}^{2n_L}) &= \bigoplus_{2\ell_R=2\ell_L} H^{2}(\overline{S}_{K_{2n_R\times 2n_L}}^{P_{2n_R\times 2n_L}}, \widehat{M}_{T_{R_\oplus}}^{2\ell_R}\otimes \widehat{M}_{T_{L_\oplus}}^{2\ell_L})\\ &\quad \bigoplus_{k_R\neq \ell_L} H^{2}(\overline{S}_{K_{2n_R\times 2\ell_L}}^{P_{2n_R\times 2\ell_L}}, \widehat{M}_{T_{R_\oplus}}^{2k_R}\otimes \widehat{M}_{T_{L_\oplus}}^{2\ell_L})\\ &\simeq \bigoplus_{\ell_R=\ell_L}\bigoplus_{j}\bigoplus_{m_j}(\mathrm{CY}_T^{2\ell_R}(\overline{X}_R[j,m_j])\times \mathrm{CY}_T^{2\ell_L}(\overline{X}_L[j,m_j]))\\ &\quad \bigoplus_{k_R\neq \ell_L}\bigoplus_{j_R}\bigoplus_{m_{j_R}}\bigoplus_{j_L}\bigoplus_{m_{j_L}}(\mathrm{CY}_T^{2k_R}(\overline{X}_R[j_R,m_{j_R}])\times \mathrm{CY}_T^{2\ell_L}(\overline{X}_L[j_L,m_{j_L}]))\end{aligned}$$

being equal to the sums of products, right by left, of the equivalence classes, counted with their multiplicities, of the irreducible semicycles.

4.2.5 Proposition

- *If* $\quad z_{n_\ell}=\sum\limits_{\alpha=1}^{2n_\ell} z_{n_\ell}(\alpha)\ |\vec e_\alpha| \in \mathbb{C}^{\,n_\ell}$ *, then every left (resp. right) n_ℓ-dimensional complex semitorus has the analytic development:*

$$\begin{aligned} &T^{2n_\ell}_L[j_L,m_{j_L}] \simeq \lambda^{\frac12}(2n_\ell,j_L,m_{j_L})\ e^{2\pi i j_L z_{n_\ell}} \\ (\text{resp.}\quad &T^{2n_\ell}_R[j_R,m_{j_R}] \simeq \lambda^{\frac12}(2n_\ell,j_R,m_{j_R})\ e^{-2\pi i j_R z_{n_\ell}}\ ). \end{aligned}$$

- *On the other hand, let*

$$\begin{aligned} &\mathrm{EIS}_L(2n_\ell,j_L,m_{j_L}) = \sum_{j_L=1}^{r}\sum_{m_{j_L}} \lambda^{\frac12}(2n_\ell,j_L,m_{j_L})\ e^{2\pi i j_L z_{n_\ell}} \\ (\text{resp.}\quad &\mathrm{EIS}_R(2n_\ell,j_R,m_{j_R}) = \sum_{j_R=1}^{r}\sum_{m_{j_R}} \lambda^{\frac12}(2n_\ell,j_R,m_{j_R})\ e^{-2\pi i j_L z_{n_\ell}}\ ). \end{aligned}$$

  be the (truncated) Fourier development of a $2n_\ell$-dimensional left (resp. right) cusp form of weight $k=2$ restricted to the upper (resp. lower) half space.

Then, we have the following analytical developments of the bilinear cohomologies:

- $H^*(\overline{S}^{P_{n=n_1+\cdots+n_s}}_{K_{n=n_1+\cdots+n_s}},\widehat{M}^{2n}_{T_{R_\oplus}}\otimes\widehat{M}^{2n}_{T_{L_\oplus}}) = \bigoplus\limits_{n_\ell}(\mathrm{EIS}_R(2n_\ell,j_R,m_{j_R})\otimes\mathrm{EIS}_L(2n_\ell,j_L,m_{j_L}))$;

- $H^{2n}(\overline{S}^{P_{2n=2_1+\cdots+2_n}}_{K_{2n=2_1+\cdots+2_n}},\widehat{M}^{2n}_{T_{R_\oplus}}\otimes\widehat{M}^{2n}_{T_{L_\oplus}}) = \bigoplus\limits_{\ell_R=\ell_L}(\mathrm{EIS}_R(2_{\ell_R},j_R,m_{j_R})\otimes\mathrm{EIS}_L(2_{\ell_L},j_L,m_{j_L}))$;

- $$\begin{aligned} H^{2n}(\overline{S}^{P_{2n_R\times 2n_L}}_{K_{2n_R\times 2n_L}},\widehat{M}^{2n_R}_{T_{R_\oplus}}\otimes\widehat{M}^{2n_L}_{T_{L_\oplus}}) &= \bigoplus_{\ell_R=\ell_L}(\mathrm{EIS}_R(2_{\ell_R},j_R,m_{j_R})\otimes\mathrm{EIS}_L(2_{\ell_L},j_L,m_{j_L})) \\ &\bigoplus_{k_R\neq\ell_L}(\mathrm{EIS}_R(2_{k_R},j_R,m_{j_R})\otimes\mathrm{EIS}_L(2_{\ell_L},j_L,m_{j_L})) \end{aligned}$$

where $\mathrm{EIS}_L(2_{\ell_L},j_L,m_{j_L})$ *corresponds to the (truncated) Fourier development of a cusp form of weight $k=2$.*

4.2.6 Reducible cusp biforms

- Let $L^{1-1}_{\mathrm{cusp}}(G^{(2n=2n_1+\cdots+2n_s)}(F^T_{\overline\omega}\times F^T_{\overline\omega}))$ denote the bialgebra of cusp biforms on the partially reducible pseudo-ramified toroidal bilinear semigroup $G^{(2n=2n_1+\cdots+2n_s)}(F^T_{\overline\omega}\times F^T_{\overline\omega})$.

Referring to section 3.4.10, a partially reducible cusp biform on this "diagonal" bialgebra is given by:

$$\begin{aligned}
&\phi^{(2n=2n_1+\cdots+2n_s)}_{G^T_R}(x_{g^T_R}) \otimes_D \phi^{(2n=2n_1+\cdots+2n_s)}_{G^T_L}(x_{g^T_L}) \\
&\quad = \bigoplus_{n_\ell=n_1}^{n_s} \phi^{(2n_\ell)}_{G^T_R} \otimes_D \phi^{(2n_\ell)}_{G^T_L} \\
&\quad = \bigoplus_{n_\ell=n_1}^{n_s} \bigoplus_{j=1}^{r} \bigoplus_{m_j} \big(\phi^{(2n_\ell)}_{G^T_{j_R}}(x_{g^T_{j_R}}) \otimes_D \phi^{(2n_\ell)}_{G^T_{j_L}}(x_{g^T_{j_L}})\big) \\
&\quad = \bigoplus_{n_\ell} (\mathrm{EIS}_R(2n_\ell, j_R, m_{j_R}) \otimes_{(D)} \mathrm{EIS}_L(2n_\ell, j_L, m_{j_L}))\ .
\end{aligned}$$

- Similarly, let $L^{1-1}_{\text{cusp}}(G^{(2n=2_1+\cdots+2_n)}(F^T_{\overline{\omega}} \times F^T_{\overline{\omega}}))$ denote the bialgebra of orthogonally completely reducible cusp biforms on $G^{(2n=2_1+\cdots+2_n)}(F^T_{\overline{\omega}} \times F^T_{\overline{\omega}})$ given by:

$$\begin{aligned}
&\phi^{(2n=2_1+\cdots+2_n)}_{G^T_R}(x_{g^T_R}) \otimes_D \phi^{(2n=2_1+\cdots+2_n)}_{G^T_L}(x_{g^T_L}) \\
&\quad = \bigoplus_{\ell_R=\ell_L} \phi^{2_{\ell_R}}_{G^T_R} \otimes_D \phi^{2_{\ell_L}}_{G^T_L} \\
&\quad = \bigoplus_{\ell_R=\ell_L} (\mathrm{EIS}_R(2_{\ell_R}, j_R, m_{j_R}) \otimes_{(D)} \mathrm{EIS}_L(2_{\ell_L}, j_L, m_{j_L}))\ .
\end{aligned}$$

- And, let $L^{1-1}_{\text{cusp}}(G^{(2n_R\times 2n_L)}(F^T_{\overline{\omega}} \times F^T_{\omega}))$ be the bialgebra of nonorthogonally completely reducible cusp biforms on $G^{(2n_R\times 2n_L)}(F^T_{\overline{\omega}} \times F^T_{\omega})$ given diagonally by:

$$\begin{aligned}
&\phi^{(2n_R=2_{1_R}+\cdots+2_{n_R})}_{G^T_R}(x_{g^T_R}) \otimes_D \phi^{(2n_L=2_{1_L}+\cdots+2_{n_L})}_{G^T_L}(x_{g^T_L}) \\
&\quad = \bigoplus_{\ell_R=\ell_L} (\phi^{2_{\ell_R}}_{G^T_R} \otimes_D \phi^{2_{\ell_L}}_{G^T_L}) \bigoplus_{k_R\neq\ell_L} (\phi^{2_{k_R}}_{G^T_R} \otimes_D \phi^{2_{\ell_L}}_{G^T_L}) \\
&\quad = \bigoplus_{\ell_R=\ell_L} (\mathrm{EIS}_R(2_{\ell_R}, j_R, m_{j_R}) \otimes_{(D)} \mathrm{EIS}_L(2_{\ell_L}, j_L, m_{j_L})) \\
&\qquad\qquad \bigoplus_{k_R\neq\ell_L} (\mathrm{EIS}_R(2_{k_R}, j_R, m_{j_R}) \otimes_{(D)} \mathrm{EIS}_L(2_{\ell_L}, j_L, m_{j_L}))\ .
\end{aligned}$$

4.2.7 Proposition

- *Let*

$$R_{G^{(2n=2n_1+\cdots+2n_s)}(F^T_{\overline{\omega}}\times F^T_{\omega})}(\phi^{(2n=2n_1+\cdots+2n_s)}_{G^T_R}\otimes\phi^{(2n=2n_1+\cdots+2n_s)}_{G^T_L})\ ,$$

$$R_{G^{(2n=2_1+\cdots+2_n)}(F^T_{\overline{\omega}}\times F^T_{\omega})}(\phi^{(2n=2_1+\cdots+2_n)}_{G^T_R}\otimes\phi^{(2n=2_1+\cdots+2_n)}_{G^T_L})\ ,$$

$$R_{G^{(2n_R\times 2n_L)}(F^T_{\overline{\omega}}\times F^T_{\omega})}(\phi^{2n_R}_{G^T_R}\otimes\phi^{2n_L}_{G^T_L})$$

be the integral operators on the respective cusp biforms introduced in sections 4.2.6 and 3.4.10.

- *Then, the trace formulas of these integral operators are respectively given by:*

$$\begin{aligned}
&\operatorname{tr}(R_{G^{(2n=2n_1+\cdots+2n_s)}(F^T_{\overline{\omega}}\times F^T_{\omega})}(\phi^{(2n=2n_1+\cdots+2n_s)}_{G^T_R}\otimes\phi^{(2n=2n_1+\cdots+2n_s)}_{G^T_L})\\
&\quad=\operatorname{vol}(G^{(2n=2n_1+\cdots+2n_s)}(F^T_{\overline{\omega}}\times F^T_{\omega}))\bigoplus_{n_\ell}(\mathrm{EIS}_R(2n_\ell,j_R,m_{j_R}),\mathrm{EIS}_L(2n_\ell,j_L,m_{j_L}))\\
&\quad=\operatorname{vol}(G^{(2n=2n_1+\cdots+2n_s)}(F^T_{\overline{\omega}}\times F^T_{\omega}))\bigoplus_{n_\ell}\bigoplus_{j}\bigoplus_{m_j}\lambda(2n_\ell,j,m_j)
\end{aligned}$$

where $(\mathrm{EIS}_R(2n_\ell,j_R,m_{j_R}),\mathrm{EIS}_L(2n_\ell,j_L,m_{j_L}))$ *is a diagonal bilinear form from* $\mathrm{Red\,cusp}(\mathrm{GL}_{(2n=2n_1+\cdots+2n_s)}(F^T_{\overline{\omega}}\times F^T_{\omega}))$ *to* $\mathbb{C}$.

$$\begin{aligned}
&\operatorname{tr}(R_{G^{(2n=2_1+\cdots+2_n)}(F^T_{\overline{\omega}}\times F^T_{\omega})}(\phi^{(2n=2_1+\cdots+2_n)}_{G^T_R}\otimes\phi^{(2n=2_1+\cdots+2_n)}_{G^T_L})\\
&\quad=\operatorname{vol}(G^{(2n=2_1+\cdots+2_n)}(F^T_{\overline{\omega}}\times F^T_{\omega}))\bigoplus_{\ell_R=\ell_L}(\mathrm{EIS}_R(2_{\ell_R},j_R,m_{j_R}),\mathrm{EIS}_L(2_{\ell_L},j_L,m_{j_L}))\\
&\quad=\operatorname{vol}(G^{(2n=2_1+\cdots+2_n)}(F^T_{\overline{\omega}}\times F^T_{\omega}))\bigoplus_{\ell_R=\ell_L}\bigoplus_{j}\bigoplus_{m_j}\lambda(2_\ell,j,m_j)
\end{aligned}$$

and finally

$$
\begin{aligned}
&\mathrm{tr}(R_{G^{(2n_R\times 2n_L)}(F^T_{\overline{\omega}}\times F^T_\omega)}(\phi^{(2n_R)}_{G^T_R}\otimes\phi^{(2n_L)}_{G^T_L})\\
&\quad=\mathrm{vol}(G^{(2n_R\times 2n_L)}(F^T_{\overline{\omega}}\times F^T_\omega))\Bigg[\bigoplus_{\ell_R=\ell_L}(\mathrm{EIS}_R(2_{\ell_R},j_R,m_{j_R}),\mathrm{EIS}_L(2_{\ell_L},j_L,m_{j_L}))\\
&\qquad+\bigoplus_{k_R=\ell_L}(\mathrm{EIS}_R(2_{k_R},j_R,m_{j_R}),\mathrm{EIS}_L(2_{\ell_L},j_L,m_{j_L}))\Bigg]\\
&\quad=\mathrm{vol}(G^{(2n_R\times 2n_L)}(F^T_{\overline{\omega}}\times F^T_\omega))\Bigg(\bigoplus_{\ell_R=\ell_L}\bigoplus_j\bigoplus_{m_j}\lambda(2_\ell,j,m_j)\\
&\qquad+\bigoplus_{k_R\neq\ell_L}\bigoplus_{j_R}\bigoplus_{j_L}\bigoplus_{m_{j_R}}\bigoplus_{m_{j_L}}\lambda^{\frac12}(2_{k_R},j_R,m_{j_R})\cdot\lambda^{\frac12}(2_{\ell_L},j_L,m_{j_L})\Bigg)
\end{aligned}
$$

Proof:

1. The proof is clear from the preceding developments and, especially from proposition 3.4.10.
2. These trace formulas are complete in the sense that they refer to a decomposition of these traces according to the direct sums of the irreducible representations of the bilinear algebraic semigroups, as in the classical case for the linear groups. ■

4.2.8 Proposition: Langlands global reducible correspondences

- *Let*

$$
\begin{aligned}
\mathrm{Red\,Rep}\,W^{2n=2n_1+\cdots+2n_s}_{F_{R\times L}}:\quad & W^{2n=2n_1+\cdots+2n_s}_{F_{\overline{\omega}}}\times W^{2n=2n_1+\cdots+2n_s}_{F_\omega}\\
&\quad\longrightarrow\ \mathrm{GL}_{n=n_1+\cdots+n_s}(\widetilde F_{\overline{\omega}_\oplus}\times\widetilde F_{\omega_\oplus})\ ,\\
\mathrm{Red\,Rep}\,W^{2n=2_1+\cdots+2_n}_{F_{R\times L}}:\quad & W^{2n=2_1+\cdots+2_n}_{F_{\overline{\omega}}}\times W^{2n=2_1+\cdots+2_n}_{F_\omega}\\
&\quad\longrightarrow\ \mathrm{GL}_{2n=2_1+\cdots+2_n}(\widetilde F_{\overline{\omega}_\oplus}\times\widetilde F_{\omega_\oplus})\ ,\\
\mathrm{Red\,Rep}\,W^{2n_R\times 2n_L}_{F_{R\times L}}:\quad & W^{2n_R}_{F_{\overline{\omega}}}\times W^{2n_L}_{F_\omega}\\
&\quad\longrightarrow\ \mathrm{GL}_{2n_R\times 2n_L}(\widetilde F_{\overline{\omega}_\oplus}\times\widetilde F_{\omega_\oplus})
\end{aligned}
$$

be respectively the partially reducible, orthogonal completely reducible and nonorthogonal completely reducible representations of the bilinear global Weil groups

$$W^{2n=2n_1+\cdots+2n_s}_{F_{\overline{\omega}}}\times W^{2n=2n_1+\cdots+2n_s}_{F_\omega}\ ,\quad W^{2n=2_1+\cdots+2_n}_{F_{\overline{\omega}}}\times W^{2n=2_1+\cdots+2_n}_{F_\omega}$$

$$\textit{and} \quad W^{2n_R}_{F_{\overline{\omega}}} \times W^{2n_L}_{F_{\omega}} \ .$$

- *Let*

$$\begin{aligned}
&\mathrm{Red\,cusp}(\mathrm{GL}_{n=n_1+\cdots+n_s}(F^T_{\overline{\omega}} \times F^T_{\omega})) \\
&\qquad = \bigoplus_{n_\ell}(\mathrm{EIS}_R(2n_\ell, j_R, m_{j_R}) \otimes \mathrm{EIS}_L(2n_\ell, j_L, m_{j_L}))\ , \\
&\mathrm{Red\,cusp}(\mathrm{GL}_{2n=2_1+\cdots+2_n}(F^T_{\overline{\omega}} \times F^T_{\omega}) \\
&\qquad = \bigoplus_{\ell_R=\ell_L}(\mathrm{EIS}_R(2_{\ell_R}, j_R, m_{j_R}) \otimes \mathrm{EIS}_L(2_{\ell_L}, j_L, m_{j_L}))\ , \\
&\mathrm{Red\,cusp}(\mathrm{GL}_{2n_R\times 2n_L}(F^T_{\overline{\omega}} \times F^T_{\omega}) \\
&\qquad = \bigoplus_{\ell_R=\ell_L}(\mathrm{EIS}_R(2_{\ell_R}, j_R, m_{j_R}) \otimes \mathrm{EIS}_L(2_{\ell_L}, j_L, m_{j_L})) \\
&\qquad\qquad \bigoplus_{k_R\neq\ell_L}(\mathrm{EIS}_R(2_{k_R}, j_R, m_{j_R}) \otimes \mathrm{EIS}_L(2_{\ell_L}, j_L, m_{j_L}))\ ,
\end{aligned}$$

be respectively the sums of the products, right by left, of the equivalence classes of the partially reducible, orthogonal completely reducible and nonorthogonal completely reducible cuspidal representations of the corresponding reducible bilinear semigroups.

- *Let*

$$\begin{gathered}
\mathrm{CY}^{2n=2n_1+\cdots+2n_s}_{T}(\overline{X}_R) \times \mathrm{CY}^{2n=2n_1+\cdots+2n_s}_{T}(\overline{X}_L)\ , \\
\mathrm{CY}^{2n=2_1+\cdots+2_n}_{T}(\overline{X}_R) \times \mathrm{CY}^{2n=2_1+\cdots+2_n}_{T}(\overline{X}_L)\ , \\
\textit{and} \quad \mathrm{CY}^{2n_R}_{T}(\overline{X}_R) \times \mathrm{CY}^{2n_L}_{T}(\overline{X}_L)\ ,
\end{gathered}$$

be respectively the sums of the products, right by left, of the equivalence classes of the partially reducible, orthogonal completely reducible and nonorthogonal completely reducible 2n-dimensional toroidal compactified cycles.

- *Then, we have the following* **Langlands global reducible correspondences***:*

$$\begin{tikzcd}
\mathrm{Red\,Rep}^{(2n)}_{W_{F_{R\times L}}}(W^{2n=2n_1+\cdots+2n_s}_{F_{\overline{\omega}}} \times W^{2n=2n_1+\cdots+2n_s}_{F_{\omega}}) \arrow[rr] \arrow[dr] & & \mathrm{Red\,cusp}(\mathrm{GL}_{n=n_1+\cdots+n_s}(F^T_{\overline{\omega}} \times F^T_{\omega})) \\
& \mathrm{CY}^{2n=2n_1+\cdots+2n_s}_{T}(\overline{X}_R) \times \mathrm{CY}^{2n=2n_1+\cdots+2n_s}_{T}(\overline{X}_L) \arrow[ur] &
\end{tikzcd}$$

$$\begin{tikzcd}
\mathrm{Red}\,\mathrm{Rep}^{(2n)}_{W_{F_{R\times L}}}(W^{2n=2_1+\cdots+2_n}_{F_{\overline{\omega}}}\times W^{2n=2_1+\cdots+2_n}_{F_{\omega}}) \arrow[rr] \arrow[dr] & & \mathrm{Red}\,\mathrm{cusp}(\mathrm{GL}_{2n=2_1+\cdots+2_n}(F^T_{\overline{\omega}}\times F^T_{\omega})) \\
& \mathrm{CY}^{2n=2_1+\cdots+2_n}_T(\overline{X}_R)\times\mathrm{CY}^{2n=2_1+\cdots+2_n}_T(\overline{X}_L) \arrow[ur] &
\end{tikzcd}$$

$$\begin{tikzcd}
\mathrm{Red}\,\mathrm{Rep}^{(2n)}_{W_{F_{R\times L}}}(W^{2n_R}_{F_{\overline{\omega}}}\times W^{2n_L}_{F_{\omega}}) \arrow[rr] \arrow[dr] & & \mathrm{Red}\,\mathrm{cusp}(\mathrm{GL}_{2n_R\times 2n_L}(F^T_{\overline{\omega}}\times F^T_{\omega})) \\
& \mathrm{CY}^{2n_R}_T(\overline{X}_R)\times\mathrm{CY}^{2n_L}_T(\overline{X}_L) \arrow[ur] &
\end{tikzcd}$$

Proof: this proposition is an adaptation of proposition 3.4.14 to the reducible case by noticing that:

- $\mathrm{CY}^{2n=2n_1+\cdots+2n_s}_T(\overline{X}_R)\times\mathrm{CY}^{2n=2n_1+\cdots+2n_s}_T(\overline{X}_L)=\widehat{G}^{2n=2n_1+\cdots+2n_s)}(F^T_{\overline{\omega}_\oplus}\times F^T_{\omega_\oplus})=\bigoplus\limits_{n_\ell}(\mathrm{EIS}_R(2n_\ell,j_R,m_{j_R})\otimes\mathrm{EIS}_L((2n_\ell,j_L,m_{j_L}))$ according to proposition 4.1.7.
- $\mathrm{Red}\,\mathrm{Rep}^{(2n)}_{W_{F_{R\times L}}}(W^{2n=2n_1+\cdots+2n_s}_{F_{\overline{\omega}}}\times W^{2n=2n_1+\cdots+2n_s}_{F_{\omega}})=\widehat{G}^{(n=n_1+\cdots+n_s)}(F_{\overline{\omega}_\oplus}\times F_{\omega_\oplus})$ is isomorphic to its toroidal equivalent $\widehat{G}^{2n=2n_1+\cdots+2n_s)}(F^T_{\overline{\omega}_\oplus}\times F^T_{\omega_\oplus})=\bigoplus\limits_{n_\ell}(\mathrm{EIS}_R(2n_\ell,j_R,m_{j_R})\times\mathrm{EIS}_L(2n_\ell,j_L,m_{j_L}))$. $\blacksquare$

4.2.9 Definition

The real boundaries of the reducible compactified bisemispaces $\overline{X}^{2n=2n_1+\cdots+2n_s}_{S_{R\times L}}$, $\overline{X}^{2n=2_1+\cdots+2_n}_{S_{R\times L}}$ and $\overline{X}^{2n_R\times 2n_L}_{S_{R\times L}}$ result from the surjective morphisms:

$$\begin{aligned}
\gamma^{\delta(2n)}_{R\times L}:\quad & \overline{X}^{2n=2n_1+\cdots+2n_s}_{S_{R\times L}}\\
&\longrightarrow \partial\overline{X}^{2n=2n_1+\cdots+2n_s}_{S_{R\times L}}=\mathrm{GL}_{n=n_1+\cdots+n_s}(F^{+,T}_R\times F^{+,T}_L)\Big/\mathrm{GL}_{n=n_1+\cdots+n_s}((\mathbb{Z}/N\,\mathbb{Z})^2)\ ,\\
\gamma^{\delta(2n)}_{R\times L}:\quad & \overline{X}^{2n=2_1+\cdots+2_n}_{S_{R\times L}}\\
&\longrightarrow \partial\overline{X}^{2n=2_1+\cdots+2_n}_{S_{R\times L}}=\mathrm{GL}_{2n=2_1+\cdots+2_n}(F^{+,T}_R\times F^{+,T}_L)\Big/\mathrm{GL}_{2n=2_1+\cdots+2_n}((\mathbb{Z}/N\,\mathbb{Z})^2)\ ,\\
\gamma^{\delta(2n_R\times 2n_L)}_{R\times L}:\quad & \overline{X}^{2n_R\times 2n_L}_{S_{R\times L}}\\
&\longrightarrow \partial\overline{X}^{2n_R\times 2n_L}_{S_{R\times L}}=\mathrm{GL}_{2n_R\times 2n_L}(F^{+,T}_R\times F^{+,T}_L)\Big/\mathrm{GL}_{2n_R\times 2n_L}((\mathbb{Z}/N\,\mathbb{Z})^2)\ ,
\end{aligned}$$

as developed in section 3.5.1.

4.2.10 Definition: the reducible Shimura bisemivarieties

The double coset decompositions of the reducible bilinear general semigroups over $(F_R^{+,T} \times F_L^{+,T})$ are given by:

$$\begin{aligned}
&\partial\overline{S}^{P_{n=n_1+\dots+n_s}}_{K_{n=n_1+\dots+n_s}} \\
&\qquad = P_{n=n_1+\dots+n_s}(F^{+,T}_{\overline{v}^1} \times F^{+,T}_{v^1}) \setminus \mathrm{GL}_{n=n_1+\dots+n_s}(F_R^{+,T} \times F_L^{+,T}) \\
&\qquad\qquad \Big/ \mathrm{GL}_{n=n_1+\dots+n_s}((\mathbb{Z}/N\,\mathbb{Z})^2)\;, \\
&\partial\overline{S}^{P_{2n=2_1+\dots+2_n}}_{K_{2n=2_1+\dots+2_n}} \\
&\qquad = P_{2n=2_1+\dots+2_n}(F^{+,T}_{\overline{v}^1} \times F^{+,T}_{v^1}) \setminus \mathrm{GL}_{2n=2_1+\dots+2_n}(F_R^{+,T} \times F_L^{+,T}) \\
&\qquad\qquad \Big/ \mathrm{GL}_{2n=2_1+\dots+2_n}((\mathbb{Z}/N\,\mathbb{Z})^2)\;, \\
&\partial\overline{S}^{P_{2n_R\times 2n_L}}_{K_{2n_R\times 2n_L}} \\
&\qquad = P_{2n_R\times 2n_L}(F^{+,T}_{\overline{v}^1} \times F^{+,T}_{v^1}) \setminus \mathrm{GL}_{2n_R\times 2n_L}(F_R^{+,T} \times F_L^{+,T}) \\
&\qquad\qquad \Big/ \mathrm{GL}_{2n_R\times 2n_L}((\mathbb{Z}/N\,\mathbb{Z})^2)\;;
\end{aligned}$$

they correspond to reducible Shimura bisemivarieties according to definition 3.5.2.

4.2.11 Proposition

The following Eisenstein cohomologies develop according to:

- $$\begin{aligned} H^*(\partial\overline{S}^{P_{n=n_1+\dots+n_s}}_{K_{n=n_1+\dots+n_s}}, \widehat{M}^{2n}_{T_{v_{R_\oplus}}} \otimes \widehat{M}^{2n}_{T_{v_{L_\oplus}}}) &= \bigoplus_{n_\ell} H^{2n_\ell}(\partial\overline{S}^{P_n}_{K_n}, \widehat{M}^{2n_\ell}_{T_{v_{R_\oplus}}} \otimes \widehat{M}^{2n_\ell}_{T_{v_{L_\oplus}}}) \\ &\simeq \bigoplus_{n_\ell}\bigoplus_{j_\delta}\bigoplus_{m_{j_\delta}} (\mathrm{CY}^{n_\ell}_T(\partial\overline{X}_R[j_\delta, m_{j_\delta}]) \times \mathrm{CY}^{n_\ell}_T(\partial\overline{X}_L[j_\delta, m_{j_\delta}])) \end{aligned}$$

- $$\begin{aligned} H^{2n}(\partial\overline{S}^{P_{2n=2_1+\dots+2_n}}_{K_{2n=2_1+\dots+2_n}}, \widehat{M}^{2n}_{T_{v_{R_\oplus}}} \otimes \widehat{M}^{2n}_{T_{v_{L_\oplus}}}) &= \bigoplus_{\ell_R=\ell_L} H^{2\ell}(\partial\overline{S}^{P_{2n}}_{K_{2n}}, \widehat{M}^{2\ell_R}_{T_{v_{R_\oplus}}} \otimes \widehat{M}^{2\ell_L}_{T_{v_{L_\oplus}}}) \\ &\simeq \bigoplus_{\ell_R=\ell_L}\bigoplus_{j_\delta}\bigoplus_{m_{j_\delta}} (\mathrm{CY}^{1_{\ell_R}}_T(\partial\overline{X}_R[j_\delta, m_{j_\delta}]) \times \mathrm{CY}^{1_{\ell_L}}_T(\partial\overline{X}_L[j_\delta, m_{j_\delta}])) \end{aligned}$$

- $$\begin{aligned} H^{2n}(\partial\overline{S}^{P_{2n_R\times 2n_L}}_{K_{2n_R\times 2n_L}}, \widehat{M}^{2n_R}_{T_{v_{R_\oplus}}} \otimes \widehat{M}^{2n_L}_{T_{v_{L_\oplus}}}) &= \bigoplus_{\ell_R=\ell_L} H^{2\ell}(\partial\overline{S}^{P_{2n_R,2n_L}}_{K_{2n_R,2n_L}}, \widehat{M}^{2\ell_R}_{T_{v_{R_\oplus}}} \otimes \widehat{M}^{2\ell_L}_{T_{v_{L_\oplus}}}) \\ &\quad \bigoplus_{k_R\neq\ell_L} H^{2k_R,2\ell_L}(\partial\overline{S}^{P_{2n_R\times 2n_L}}_{K_{2n_R\times 2n_L}}, \widehat{M}^{2k_R}_{T_{v_{R_\oplus}}} \otimes \widehat{M}^{2\ell_L}_{T_{v_{L_\oplus}}}) \\ &\simeq \bigoplus_{\ell_R=\ell_L}\bigoplus_{j_\delta}\bigoplus_{m_{j_\delta}} (\mathrm{CY}^{1_{\ell_R}}_T(\partial\overline{X}_R[j_\delta, m_{j_\delta}]) \times \mathrm{CY}^{1_{\ell_L}}_T(\partial\overline{X}_L[j_\delta, m_{j_\delta}])) \\ &\quad \bigoplus_{k_R\neq\ell_L}\bigoplus_{j^\delta_R}\bigoplus_{m_{j^\delta_R}}\bigoplus_{j^\delta_L}\bigoplus_{m_{j^\delta_L}} (\mathrm{CY}^{1_{k_R}}_T(\partial\overline{X}_R[j^\delta_R, m_{j^\delta_R}]) \times \mathrm{CY}^{1_{\ell_L}}_T(\partial\overline{X}_L[j^\delta_L, m_{j^\delta_L}])) \end{aligned}$$

they are equal to the sums of products, right by left, of the equivalence classes, counted with their multiplicities, of the irreducible cycles.

Proof: these decompositions of cohomologies are the real equivalents of those considered in proposition 4.2.4. ■

4.2.12 Proposition

a) If $x_{n_\ell}=\sum\limits_{\beta=1}^{n_\ell} x_{n_\ell}(\beta)\ |\vec{e}_\beta|\in\mathbb{R}^{n_\ell}$,

then, every left (resp. right) n_ℓ*-dimensional real semitorus has the analytic development:*

$$\begin{aligned} T_L^{n_\ell}[j_L^\delta,m_{j_L^\delta}] &\simeq \lambda^{\frac12}(n_\ell,j_L^\delta,m_{j_L^\delta})\ e^{2\pi i j_L^\delta x_{n_\ell}} \\ (\textit{resp.}\quad T_R^{n_\ell}[j_R^\delta,m_{j_R^\delta}] &\simeq \lambda^{\frac12}(n_\ell,j_R^\delta,m_{j_R^\delta})\ e^{-2\pi i j_R^\delta x_{n_\ell}}\ ). \end{aligned}$$

b) On the other hand, let

$$\begin{aligned} \mathrm{ELLIP}_L(2n_\ell,j_L^\delta,m_{j_L^\delta}) &= \sum_{j_L^\delta=1}^{t}\sum_{m_{j_L^\delta}} \lambda^{\frac12}(n_\ell,j_L^\delta,m_{j_L^\delta})\ e^{2\pi i j_L^\delta x_{n_\ell}} \\ (\textit{resp.}\quad \mathrm{ELLIP}_R(2n_\ell,j_R^\delta,m_{j_R^\delta}) &= \sum_{j_R^\delta=1}^{t}\sum_{m_{j_R^\delta}} \lambda^{\frac12}(n_\ell,j_R^\delta,m_{j_R^\delta})\ e^{-2\pi i j_R^\delta x_{n_\ell}})\ ),\quad t\le\infty\ , \end{aligned}$$

be the (truncated) Fourier development of a left (resp. right) n_ℓ*-dimensional solvable global elliptic semimodule where*

- $\lambda(n_\ell,j_\delta,m_{j_\delta})=\prod\limits_{c=1}^{n_\ell}\lambda_c(n_\ell,j_\delta,m_{j_\delta})\approx j^{n_\ell}\cdot N^{n_\ell}$ *;*
- $e^{2\pi i x_{n_\ell}}\to e^{2\pi i j_L^\delta x_{n_\ell}}$ *(resp.* $e^{-2\pi i x_{n_\ell}}\to e^{-2\pi i j_R^\delta x_{n_\ell}}$ *) is a left (resp. right) global Frobenius substitution.*

Then, we have the following analytic developments of the Eisenstein bilinear cohomologies of the equivalents of the reducible Shimura bisemivarieties:

- $H^*(\partial\overline{S}{}^{P_{n=n_1+\dots+n_s}}_{K_{n=n_1+\dots+n_s}},\widehat{M}^{2n}_{T_{v_{R_\oplus}}}\otimes\widehat{M}^{2n}_{T_{v_{L_\oplus}}})$
  $$=\bigoplus_{2n_\ell}(\mathrm{ELLIP}_R(2n_\ell,j_R^\delta,m_{j_R^\delta})\otimes\mathrm{ELLIP}_L(2n_\ell,j_L^\delta,m_{j_L^\delta}))$$
- $H^{2n}(\partial\overline{S}{}^{P_{2n=2_1+\dots+2_n}}_{K_{2n=2_1+\dots+2_n}},\widehat{M}^{2n}_{T_{v_{R_\oplus}}}\otimes\widehat{M}^{2n}_{T_{v_{L_\oplus}}})$
  $$=\bigoplus_{\ell_R=\ell_L}(\mathrm{ELLIP}_R(2_{\ell_R},j_R^\delta,m_{j_R^\delta})\otimes\mathrm{ELLIP}_L(2_{\ell_L},j_L^\delta,m_{j_L^\delta}))$$

- $H^{2n}(\partial\overline{S}^{P_{2n_R\times 2n_L}}_{K_{2n_R\times 2n_L}}, \widehat{M}^{2n_R}_{T_{v_{R_\oplus}}} \otimes \widehat{M}^{2n_L}_{T_{v_{L_\oplus}}})$

$$\begin{aligned} &= \bigoplus_{\ell_R=\ell_L} (\mathrm{ELLIP}_R(2_{\ell_R}, j^\delta_R, m_{j^\delta_R}) \otimes \mathrm{ELLIP}_L(2_{\ell_L}, j^\delta_L, m_{j^\delta_L}))\\ &\qquad\qquad \bigoplus_{k_R\neq\ell_L} (\mathrm{ELLIP}_R(2_{k_R}, j^\delta_R, m_{j^\delta_R}) \otimes \mathrm{ELLIP}_L(2_{\ell_L}, j^\delta_L, m_{j^\delta_L}))\end{aligned}$$

Proof: referring to proposition 3.5.4 and 3.5.5, each equivalence class representative $\mathrm{CY}^{n_\ell}_T(\partial\overline{X}_R[j_\delta, m_{j_\delta}])$ (resp. $\mathrm{CY}^{n_\ell}_T(\partial\overline{X}_L[j_\delta, m_{j_\delta}])$) of a semicycle of codimension n_ℓ corresponds to the analytic representation of a n_ℓ-dimensional real semitorus $T^{n_\ell}_R[j_\delta, m_{j_\delta}]$ (resp. $T^{n_\ell}_L[j_\delta, m_{j_\delta}]$). ■

4.2.13 Remark

Trace formulas for the real reducible cases considered here can be developed similarly as it was done for the complex cases in sections 4.2.6 and 4.2.7.

4.2.14 Proposition: Langlands global correspondences on the equivalents of the reducible Shimura bisemivarieties

- *Let*

$$\begin{aligned} &\mathrm{Red\,Rep}^{(n)}_{W_{F^+_{R\times L}}}(W^{n=n_1+\cdots+n_s}_{F^+_{\overline{v}}} \times W^{n=n_1+\cdots+n_s}_{F^+_v})\\ &\qquad\longrightarrow H^*(\partial\overline{S}^{P_{n=n_1+\cdots+n_s}}_{K_{n=n_1+\cdots+n_s}}, \widehat{M}^{2n}_{T_{v_{R_\oplus}}} \otimes \widehat{M}^{2n}_{T_{v_{L_\oplus}}})\\ &\qquad\qquad = \mathrm{FRepsp}(\mathrm{GL}_{n=n_1+\cdots+n_s}(F^{+,T}_{\overline{v}_\oplus} \times F^{+,T}_{v_\oplus}))\ ,\end{aligned}$$

$$\begin{aligned} &\mathrm{Red\,Rep}^{(n)}_{W_{F^+_{R\times L}}}(W^{2n=2_1+\cdots+2_n}_{F^+_{\overline{v}}} \times W^{2n=2_1+\cdots+2_n}_{F^+_v})\\ &\qquad\longrightarrow H^{2n}(\partial\overline{S}^{P_{2n=2_1+\cdots+2_n}}_{K_{2n=2_1+\cdots+2_n}}, \widehat{M}^{2n^*}_{T_{v_{R_\oplus}}} \otimes \widehat{M}^{2n^*}_{T_{v_{L_\oplus}}})\\ &\qquad\qquad = \mathrm{FRepsp}(\mathrm{GL}_{2n=2_1+\cdots+2_n}(F^{+,T}_{\overline{v}_\oplus} \times F^{+,T}_{v_\oplus}))\ ,\end{aligned}$$

$$\begin{aligned} &\mathrm{Red\,Rep}^{(n)}_{W_{F^+_{R\times L}}}(W^{2n_R}_{F^+_{\overline{v}}} \times W^{2n_L}_{F^+_v})\\ &\qquad\longrightarrow H^{2n}(\partial\overline{S}^{P_{2n_R\times 2n_L}}_{K_{2n_R\times 2n_L}}, \widehat{M}^{2n_R}_{T_{v_{R_\oplus}}} \otimes \widehat{M}^{2n_L}_{T_{v_{L_\oplus}}})\\ &\qquad\qquad = \mathrm{FRepsp}(\mathrm{GL}_{2n_R\times 2n_L}(F^{+,T}_{\overline{v}_\oplus} \times F^{+,T}_{v_\oplus}))\ ,\end{aligned}$$

be respectively the sums of the products, right by left, of the equivalence classes of the partially reducible, orthogonal completely reducible and nonorthogonal completely reducible representations of the bilinear global Weil groups $(W^{n=n_1+\cdots+n_s}_{F^+_{\overline{v}}} \times W^{n=n_1+\cdots+n_s}_{F^+_v})$, $(W^{2n=2_1+\cdots+2_n}_{F^+_{\overline{v}}} \times W^{2n=2_1+\cdots+2_n}_{F^+_v})$ *and* $(W^{2n_R\times 2n_L}_{F^+_{\overline{v}}} \times W^{2n_R\times 2n_L}_{F^+_v})$.

- *Let*

$$\begin{aligned}
&\mathrm{Red}\,\mathrm{ELLIP}(\mathrm{GL}_{n=n_1+\cdots+n_s}(F^{+,T}_{\overline{v}}\times F^{+,T}_{v})\\
&\qquad=\bigoplus_{2n_\ell}(\mathrm{ELLIP}_R(2n_\ell,j^\delta_R,m_{j^\delta_R})\otimes\mathrm{ELLIP}_L(n_\ell,j^\delta_L,m_{j^\delta_L}))\ ,\\[2ex]
&\mathrm{Red}\,\mathrm{ELLIP}(\mathrm{GL}_{2n=2_1+\cdots+2_n}(F^{+,T}_{\overline{v}}\times F^{+,T}_{v})\\
&\qquad=\bigoplus_{\ell_R=\ell_L}(\mathrm{ELLIP}_R(2_{\ell_R},j^\delta_R,m_{j_R})\otimes\mathrm{ELLIP}_L(2_{\ell_L},j^\delta_L,m_{j^\delta_L}))\ ,\\[2ex]
&\mathrm{Red}\,\mathrm{ELLIP}(\mathrm{GL}_{2n_R\times 2n_L}(F^{+,T}_{\overline{v}}\times F^{+,T}_{v})\\
&\qquad=\bigoplus_{\ell_R=\ell_L}(\mathrm{ELLIP}_R(2_{\ell_R},j^\delta_R,m_{j^\delta_R})\otimes\mathrm{ELLIP}_L(2_{\ell_L},j^\delta_L,m_{j^\delta_L}))\\
&\qquad\quad\bigoplus_{k_R\neq\ell_L}(\mathrm{ELLIP}_R(2_{k_R},j^\delta_R,m_{j^\delta_R})\otimes\mathrm{ELLIP}_L(2_{\ell_L},j^\delta_L,m_{j^\delta_L}))\ ,
\end{aligned}$$

  be respectively the sums of the products, right by left, of the equivalence classes of the partially reducible, orthogonal completely reducible and nonorthogonal completely reducible global elliptic representations of the corresponding reducible bilinear semigroups.

- *Let finally*

$$\mathrm{CY}^{n=n_1+\cdots+n_s}_{T}(\partial\overline{X}_R)\times\mathrm{CY}^{n=n_1+\cdots+n_s}_{T}(\partial\overline{X}_L)\ ,$$

$$\mathrm{CY}^{n=1_1+\cdots+1_n}_{T}(\partial\overline{X}_R)\times\mathrm{CY}^{n=1_1+\cdots+1_n}_{T}(\partial\overline{X}_L)\ ,$$

$$\textit{and}\quad\mathrm{CY}^{n_R}_{T}(\partial\overline{X}_R)\times\mathrm{CY}^{n_L}_{T}(\partial\overline{X}_L)\ ,$$

  be respectively the sums of the products, right by left, of the equivalence classes of the partially reducible, orthogonal completely reducible and nonorthogonal completely reducible n-dimensional toroidal compactified cycles over $\partial\overline{X}_R$ and $\partial\overline{X}_L$.

- *Then, we have the following **Langlands global reducible correspondences** on the equivalents of the reducible Shimura bisemivarieties:*

$$\begin{tikzcd}
\mathrm{Red}\,\mathrm{Rep}^{(n)}_{W_{F^+_{R\times L}}}(W^{n=n_1+\cdots+n_s}_{F^+_{\overline{v}}}\times W^{n=n_1+\cdots+n_s}_{F^+_{v}}) \arrow[rr] \arrow[dr] & & \mathrm{Red}\,\mathrm{ELLIP}(\mathrm{GL}_{n=n_1+\cdots+n_s}(F^{+,T}_{\overline{v}}\times F^{+,T}_{v})) \\
& \mathrm{CY}^{n=n_1+\cdots+n_s}_{T}(\partial\overline{X}_R)\times\mathrm{CY}^{n=n_1+\cdots+n_s}_{T}(\partial\overline{X}_L) \arrow[ur] &
\end{tikzcd}$$

$$\begin{tikzcd}
\mathrm{Red\,Rep}^{(n)}_{W_{F^+_{R\times L}}}(W^{2n=2_1+\cdots+2_n}_{F^+_{\overline{v}}} \times W^{2n=2_1+\cdots+2_n}_{F^+_{v}}) \arrow[rr] \arrow[dr] & & \mathrm{Red\,ELLIP}(\mathrm{GL}_{2n=2_1+\cdots+2_n}(F^{+,T}_{\overline{v}} \times F^{+,T}_{v})) \\
& \mathrm{CY}^{n=1_1+\cdots+1_n}_{T}(\partial\overline{X}_R) \times \mathrm{CY}^{n=1_1+\cdots+1_n}_{T}(\partial\overline{X}_L) \arrow[ur] &
\end{tikzcd}$$

$$\begin{tikzcd}
\mathrm{Red\,Rep}^{(n)}_{W_{F^+_{R\times L}}}(W^{2n_R}_{F^+_{\overline{v}}} \times W^{2n_L}_{F^+_{v}}) \arrow[rr] \arrow[dr] & & \mathrm{Red\,ELLIP}(\mathrm{GL}_{2n_R\times 2n_L}(F^{+,T}_{\overline{v}} \times F^{+,T}_{v})) \\
& \mathrm{CY}^{n_R}_{T}(\partial\overline{X}_R) \times \mathrm{CY}^{n_L}_{T}(\partial\overline{X}_L) \arrow[ur] &
\end{tikzcd}$$

Proof: this is an adaptation to the real case of proposition 4.2.8 (see also proposition 3.4.10). ■

Appendix

1 Philosophy of bisemiobjects

Instead of working with classical mathematical objects, the developments of this paper deal with the products of semiobjects. In this respect, a mathematical symmetric object " O " will be cut into two semiobjects, labeled left and right, in such a way that the left semiobject O_L be localized in (or refers to) the upper half space while the right symmetric semiobject O_R is localized in (or refers to) the lower half space. Then, informations concerning the internal mathematical structure of our object " O " can be more advantageously extracted from the product $O_R\times O_L$ of the two semiobjects O_R and O_L . Indeed, the endomorphism of our object " O " can be decomposed into the product $E_R\times E_L$ of the endomorphisms $E_L:O_L\to O_L$ acting on the semiobject O_L by the opposite endomorphism $E_R:O_R\to O_R$ acting on O_R such that $E_R=E_L^{-1}$. By this way, the action of E_L is neutralized by the coaction of E_R .

A classical analogue of this endormophism $E_R\times E_L$ on $O_R\times O_L$ is, for example, the normal endomorphism E_N of a group G given by:

$$a\ E_N(b)\ a^{-1}=E_N(a\ b\ a^{-1})\quad \text{for } a,b\in G\ .$$

(In the philosophy of bisemiobjects, the element b would have to be decomposed into the product $b_R\times b_L$ of two semielements b_R and b_L , if mathematically feasible.)

2 Semistructures

So, semistructures will be recalled or introduced in the following. The notation L,R means "left" or "right".

- We are concerned with a **semigroup** $G_{L,R}$, called left of right, which is a nonempty set of left (resp. right) elements, localized or referring to the upper (resp. lower) half space, together with a binary operation on $G_{L,R}$, i.e. a function $G_{L,R}\times G_{L,R}\to G_{L,R}$, or $G_{L,R}\times G_{L,L}\to G_{L,R}$.
- A left (resp. right) **monoid** is a left (resp. right) semigroup $G_{L,R}$ which contains an identity element $e_{L,R}\in G_{L,R}$ such that:

$$e_L\cdot a_L=a_L\quad (\text{resp. } a_R\cdot e_R=a_R\ ),\qquad \forall\ a_{L,R}\in G_{L,R}\ .$$

- A left (resp. right) **semiring** is a nonempty set $R_{L,R}$ together with two binary operations (addition and multiplication) such that:

a) $(R_{L,R},+)$ is an abelian left (resp. right) semigroup;

b) $(a_{L,R}\ b_{L,R})\ c_{L,R} = a_{L,R}\ (b_{L,R}\ c_{L,R}) \quad \forall\ a_{L,R}, b_{L,R}, c_{L,R} \in R_{L,R}$ (associative multiplication);

c) $a_{L,R}\ (b_{L,R} + c_{L,R}) = a_{L,R}\ b_{L,R} + a_{L,R}\ c_{L,R}$ and $(a_{L,R} + b_{L,R})\ c_{L,R} = a_{L,R}\ c_{L,R} + b_{L,R}\ c_{L,R}$ (left and right distribution).

- If $R_{L,R}$ is a commutative semiring with identity $\mathbb{1}_{R_{L,R}}$ and no zero divisors, it will be called a left (resp. right) integral semidomain.

  Furthermore, if every element of $R_{L,R}$ is a unit (i.e. left and right invertible), $R_{L,R}$ is a **division semiring**.

  And, a left (resp. right) **semifield** is a commutative division semiring.

- A **left (resp. right) adele semiring** is the product of the completions of the left (resp. right) considered semifield.

- Let $R_{L,R}$ be a left (resp. right) semiring. A left (resp. right) $\boldsymbol{R_{L,R}}$**-semimodule** is an additive abelian left (resp. right) semigroup $M_{L,R}$ together with a function $R_{L,R} \times M_L \to M_L$ (resp. $M_R \times R_{R,L} \to M_R$) such that:

a) $$r_L\ (a_L + b_L) = r_L\ a_L + r_L\ b_L\ , \qquad \forall\ r_L \in R_L\ ,\ a_L, b_L \in M_L$$

$$\text{(resp.}\quad (a_R + b_R)\ r_R = a_R\ r_R + b_R\ r_R)\ , \quad \forall\ r_R \in R_R\ ,\ a_R, b_R \in M_R\ );$$

b) $$(r_L + s_L)\ a_L = r_L\ a_L + s_L\ a_L\ , \quad \forall\ s_L \in R_L$$

$$\text{(resp.}\quad a_R\ (r_R + s_R) = a_R\ r_R + a_R\ s_R\ , \quad \forall\ s_R \in R_R\ );$$

c) $$r_L\ (s_L\ a_L) = (r_L\ s_L)\ a_L$$

$$\text{(resp.}\quad (a_R\ s_R)\ r_R = a_R\ (s_R\ r_R)\ ).$$

If $R_{L,R}$ has an identity element $\mathbb{1}_{L,R}$ such that $\mathbb{1}_L \cdot a_L = a_L$ (resp. $a_R \cdot \mathbb{1}_R = a_R$), $M_{L,R}$ is a left (resp. right) unitary $R_{L,R}$-semimodule.

If $R_{L,R}$ is a left (resp. right) division semiring, then the unitary left (resp. right) unitary $R_{L,R}$-semimodule, is a left (resp. right) **vector semispace**.

- If $R_{L,R}$ is a commutative semiring with identity, a $\boldsymbol{R_{L,R}}$**-semialgebra** $\boldsymbol{A_{L,R}}$ is a semiring $A_{L,R}$ such that:

  a) $(A_{L,R},+)$ is a unitary left (resp. right) $R_{L,R}$-semimodule;

  b) $$r_L\ (a_L\ b_L)=(r_L\ a_L)\ b_L=a_L\ (r_L\ b_L)\ ,\quad \forall\ r_L\in R_L\ ,\ a_L,b_L\in A_L$$

  $$\text{(resp.}\quad (a_R\ b_R)\ r_R=a_R\ (b_R\ r_R)=b_R\ (a_R\ r_R)\ ,\ \ \forall\ r_R\in R_R\ ,\ a_R,b_R\in A_R\ ).$$

  If $A_{L,R}$ is a division semiring, then $A_{L,R}$ is called a division semialgebra.

3 Bisemistructures [Pie5]

Before recalling the salient features of the algebraic bilinear semigroups, it is useful to give a precise definition of a R_R-R_L-bisemimodule M_{R-L} and of a $R_R\times R_L$-bisemimodule $M_R\otimes M_L$.

- Let R_L (resp. R_R) be a left (resp. right) semiring.

  Then, a $\boldsymbol{R_R-R_L}$**-bisemimodule** is an abelian (semi)group M_{R-L} (under the addition) with the structure of both a left R_L-semimodule M_L and a right R_R-semimodule M_R and together with a bifunction $R_{L,R}\times M_{L-R}\times R_{R,L}\to M_{R-L}$ such that:

  a) $M_{L-R}\equiv M_{R-L}=M_R\oplus M_L$.

  b) $(r_L\ a_L)\ (a_R\ r_R)=r_L\ (a_L\ a_R\ r_R)=(r_L\ a_L\ a_R)\ r_R$, $\forall\ a_L\in M_L$, $a_R\in M_R$, $r_L\in r_L$, $r_R\in R_R$.

  c) the conditions, mentioned above for a left (resp. right) $R_{L,R}$-semimodule, are applied on the left and on the right of M_{L-R} .

- **A** $\boldsymbol{R_R\times R_L}$**-bisemimodule** $\boldsymbol{M_R\otimes M_L}$ is a bilinear semigroup given by the map : $f:M_R\times M_L\to M_R\otimes M_L$, where f sends pairs $(a_{R_i},a_{L_i})\in M_R\times M_L$ of symmetric elements to their products $(a_{R_i}\times a_{L_i})$ such that:

  a) $f(a_{R_1}+a_{R_2},a_{L_1})=f(a_{R_1},a_{L_1})+f(a_{R_2},a_{L_1})$;

  b) $f(a_{R_1},a_{L_1}+a_{L_2})=f(a_{R_1},a_{L_1})+f(a_{L_1},a_{L_2})$;

  c) $f(a_R\ r_R,r_L\ a_L)\neq f(r_L\ a_R,a_L\ r_L)$, $\forall\ a_{R_i}\in M_R$, $i=1,2$, $a_{L_i}\in M_L$.

- Taking into account the section 1 of the appendix and the introduction, we have to specify **the product $\boldsymbol{X_R \times X_L}$ of a right affine semigroup scheme $\boldsymbol{X_R}$ by its left symmetric equivalent $\boldsymbol{X_L}$** as occurring from:

$$(X_R \circ \phi_R^{\prime -1}) \times (X_L \circ \phi_L^{\prime -1})](\widetilde{F}_R \times \widetilde{F}_L) = \{(v_R \times v_L) \mid v_R \in V_R \ , \ v_L \in V_L\}$$

  where $(v_R \times v_L)$ is a bipoint of an affine bisemispace $(V_R \otimes V_L)$ resulting from the bihomomorphism

$$X_R \times X_L : \quad Q_R \times Q_L \longrightarrow V_R \otimes V_L$$

  which sends the product of the quotient k-algebras Q_R and Q_L into $(V_R \otimes V_L)$.

- It was seen in chapter 2 that $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$ is an **algebraic bilinear semigroup** of matrices in the sense that its algebraic representation space is an $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$-bisemimodule $\widetilde{M}_R \otimes \widetilde{M}_L$ defined as follows:

  Let $\widetilde{M}_L$ (resp. $\widetilde{M}_R$) be an n-dimensional $T_n(\widetilde{F}_\omega)$- (resp. $T_n^t(\widetilde{F}_{\overline{\omega}})$)-semimodule. Then, the canonical bilinear map $\widetilde{M}_R \times \widetilde{M}_L \to \widetilde{M}_R \otimes \widetilde{M}_L$ defines the bisemimodule $\widetilde{M}_R \otimes \widetilde{M}_L$, provided that $\widetilde{M}_R \otimes \widetilde{M}_L$ is the bilinear tensor product of a left $T_n(\widetilde{F}_\omega)$-semimodule by a right $T_n^t(\widetilde{F}_{\overline{\omega}})$-semimodule.

- The **complete algebraic bilinear semigroup** $G^{(n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega)$ is generated by the bilinear algebraic semigroup of matrices $\mathrm{GL}_n(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_\omega) \equiv T_n^t(\widetilde{F}_{\overline{\omega}}) \times \widetilde{T}_n(F_\omega)$ where $T_n(\widetilde{F}_\omega)$ (resp. $T_n^t(\widetilde{F}_{\overline{\omega}})$) is the group of upper (resp. lower) triangular matrices.

- The complete algebraic bilinear semigroup $G^{(n)}(F_{\overline{\omega}} \times F_\omega)$ can be viewed as **a bilinear Lie semigroup** defined as follows:

  1. $\boldsymbol{G^{(n)}(F_{\overline{\omega}} \times F_\omega)}$ **is a topological bilinear semigroup**, that is to say that:
     (a) it is a **bilinear semigroup** $G_{R\times L}$:
        α) given by a function:

$$\begin{aligned} G_R \times G_L &\longrightarrow G_{R\times L} \\ (g_{r_i}, g_{L_i}) &\longrightarrow (g_{r_i} \times g_{L_i}) \end{aligned}$$

        sending pairs (g_{r_i}, g_{L_i}) of symmetric elements, localized in (or referring to) the upper and lower half spaces, to their products $(g_{r_i} \times g_{L_i})$, $i \in \mathbb{N}$.

        β) submitted to a binary operation $\underline{\times}$

$$G_{R\times L} \underline{\times} G_{R\times L} \longrightarrow G_{R\times L}$$

defined by

$$(g_{r_i} \times g_{L_i}) \underline{\times} (g_{r_j} \times g_{L_j}) \longrightarrow (g_{r_i} + g_{r_j}) \times (g_{L_i} + g_{L_j})$$

in such a way that cross products $(g_{r_i} \times g_{L_j})$ could be generated.

  (b) $\boldsymbol{G^{(n)}(F_{\overline{\omega}} \times F_{\omega})}$ **is a topological bisemispace**, product of a right topological semispace, restricted to the lower half space by its left symmetric equivalent.

2. $\boldsymbol{G^{(n)}(F_{\overline{\omega}} \times F_{\omega})}$ **is an abstract bisemivariety**, i.e. the product of a left semivariety restricted to the upper half space by the symmetric right semivariety restricted to the lower half space.

- Let $P^{(n)}(F_{\overline{\omega}^1} \times F_{\omega^1}) = P^{(n)}(F_{\overline{\omega}^1}) \times P^{(n)}(F_{\omega^1})$ be a bilinear subgroup of $G^{(n)}(F_{\overline{\omega}} \times F_{\omega})$ such that F_{ω^1} (resp. $F_{\overline{\omega}^1}$) denotes the set of left (resp. right) irreducible completions (see section 2.4.1).

  Let $a_{R,L}, b_{R,L} \in G^{(n)}(F_{\overline{\omega}} \times F_{\omega})$.

  a_R is right congruent to b_R modulo $P^{(n)}(F_{\overline{\omega}^1})$, denoted $a_R \equiv b_R \operatorname{mod}(P^{(n)}(F_{\overline{\omega}^1})$ if $a_R\, b_R^{-1} \in P^{(n)}(F_{\overline{\omega}^1})$.

  a_L is left congruent to b_L modulo $P^{(n)}(F_{\omega^1})$, denoted $a_L \equiv b_L \operatorname{mod}(P^{(n)}(F_{\omega^1})$ if $a_L\, b_L^{-1} \in P^{(n)}(F_{\omega^1})$.

  Then, $P^{(n)}(F_{\overline{\omega}^1} \times F_{\omega^1})$ is a **bilinear normal subgroup** of $G^{(n)}(F_{\overline{\omega}} \times F_{\omega})$ if left congruence modulo $P^{(n)}(F_{\omega^1})$ correspond to right congruence modulo $P^{(n)}(F_{\overline{\omega}^1})$.

- The **endomorphism** of an algebraic bilinear semigroup $G^{(n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$ is a homomorphism

  $$E_{R\times L} : \quad G^{(n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega}) \longrightarrow G^{(n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega})$$

  which can be decomposed into:

  $$\begin{aligned} E_R : &\quad G^{(n)}(\widetilde{F}_{\overline{\omega}}) \longrightarrow G^{(n)}(\widetilde{F}_{\overline{\omega}}) \; , \\ E_L : &\quad G^{(n)}(\widetilde{F}_{\omega}) \longrightarrow G^{(n)}(\widetilde{F}_{\omega}) \; , \end{aligned}$$

  for $G^{(n)}(\widetilde{F}_{\overline{\omega}} \times \widetilde{F}_{\omega}) = G^{(n)}(\widetilde{F}_{\overline{\omega}}) \times G^{(n)}(\widetilde{F}_{\omega})$ such that:

  1. $E_R(a_R\; b_R) = E_R(a_R) \cdot E_R(b_R)$ for all $a_R, b_R \in G^{(n)}(\widetilde{F}_{\overline{\omega}})$;

  2. $E_L(a_L\; b_L) = E_L(a_L) \cdot E_L(b_L)$ for all $a_L, b_L \in G^{(n)}(\widetilde{F}_{\omega})$;

  3. $E_{R\times L}(a_R\; b_R\; a_L\; b_L) = E_{R\times L}(a_R \cdot a_L) \cdot E_{R\times L}(b_R \cdot b_L)$.

If $E_R = E_L^{-1}$, then $E_R(a_R\ b_R) = E_L^{-1}(a_L^{-1}\ b_L^{-1})$ where a_L^{-1}, b_L^{-1} denote the opposites with respect to the addition or the complex conjugates of a_L, b_L .

So,

$$\begin{aligned} & E_{R\times L}(a_R\ b_R\ a_L\ a_L) = E_L^{-1} \times E_L(a_L^{-1}\ b_L^{-1}\ a_L\ b_L) \\ \text{or} \quad & E_{R\times L}(b_R\ a_R\ a_L\ b_L) = E_L^{-1} \times E_L(b_L^{-1}\ a_L^{-1}\ a_L\ b_L)\ . \end{aligned}$$

References

[A-M] A. Alekseev, E. Meinrenken, The non commutative Weil algebra, *Invent. Math.*, **139** (2000), 135–172.

[And] Y. Andre, Pour une théorie inconditionnelle des motifs, *Publ. Math. IHES*, **83** (1996), 5–49.

[Ande] G.W. Anderson, t-Motives, *Duke Math. J.*, **53** (1986), 457–502.

[Ar-Ma] M. Artin, B. Mazur, Formal groups arising from algebraic varieties, *Ann. Scient. Ec. Norm. Sup.*, **10** (1977), 87–132.

[Art1] J. Arthur, Eisenstein series and trace formula, *Proc. Symp. Pure Math.*, **33** (1979), 253–274.

[Art2] J. Arthur, A trace formula for reductive groups. I. Terms associated to classes in $G(\mathbb{Q})$, *Duke Math. J.*, **45** (1978), 911–953.

[Art3] J. Arthur, The principle of functoriality, *Bull. Amer. Math. Soc.*, **40** (2002), 39–53.

[Bor1] A. Borel, Introduction aux groupes arithmétiques, *Act. Scient. et Industr.*, **1341** (1969), Hermann.

[Bor2] A. Borel, Linear algebraic groups, *Proceed. Symp. Pure Math.*, **9** (1966), 3–19.

[B-S] A. Borel, J.P. Serre, Corners and arithmetic groups, *Comment. Math. Helv.*, **48** (1973), 436–491.

[B-Z] I. Bernstein, A. Zelevinsky, Induced representations of reducive p-adic groups, *Ann. Scient. Ec. Norm. Sup.*, **10** (1977), 441–472.

[Cara] H. Carayol, Preuve de la conjecture de Langlands locale pour GL_n : travaux de Harris-Taylor et Henniart, *Sém. Bourbaki*, **857** (1999), 1–52.

[Car] H. Cartan, Notions d'algèbre différentielle. In: H. Cartan, Oeuvres, *Collected Works*, Vol. 3 (1979), Springer, **86**, 1255–1237 & **87**, 1268–1282.

[Cart] P. Cartier, An introduction to zeta functions from number theory to physics (1990), 1–63, Springer.

[C-K-P-S] C.W. COGDELL, H. KIM, I. PIATETSKI-SHAPIRO, F. SHAKIDI, Functoriality for the classical groups, *Publ. Math. IHES*, **99** (2004), 163–233.

[Clo] L. CLOZEL, Représentations Galoisiennes associées aux représentations automorphes autoduales de $GL(n)$, *Publ. Math. IHES*, **73** (1991), 97–145.

[Dan] V.I.. DANILOV, The geometry of toric varieties, *Russ. Math. Surveys*, **33** (1978); 97–154.

[Del1] P. DELIGNE, Formes modulaires et représentations ℓ-adiques, *Lect. Notes Math.*, **179** (1972); *Sém. Bourbaki*, **355**, 139–172.

[Del2] P. DELIGNE, Travaux de Shimura, *Lect. Notes Math.*, **244** (1970); *Sém. Bourbaki*, **389**, 123–165.

[Del3] P. DELIGNE, Formes modulaires et représentations de GL(2) , *Lect. Notes Math.*, **349** (1973), 55–105.

[Del4] P. DELIGNE, J.S. MILNE, Tannakian categories, *Lect. Notes Math.*, **900** (1982), 101–228.

[D-M] C. DENINGER, J. MURRE, Motivic decomposition of abelian schemes and the Fourier transform, *J. Reine Angew. Math.*, **422** (1991), 201–219.

[D-M-O-S] P. DELIGNE, J. MILNE, A. OGUS, K. SHIH, Hodge cycles, motives and Shimura varieties, *Lect. Notes Math.*, **900**, 9–99 (1982), Springer.

[Duf] M. DUFLO, Opérateurs Différentiels bi-invariants sur un groupe de Lie, *Ann. Scient. Ec. Norm. Sup.*, **10** (1977), 265–288.

[D-F] D. DUMMIT, R. FOOTE, Abstract algebra, *Prentice Hall*.

[Frie] E. FRIEDLANDER, Motivic complexes of Suslin and Voevodsky, *Sém. Bourbaki*, **883** (1996-97).

[F-M] W. FULTON, R. MACPHERSON, A compactification of configuration spaces, *Annals of Math*, **139** (1994), 183–225.

[Gel] S. GELBART, An elementary introduction to the Langlands program, *Bull. Amer. Math. Soc.*, **10** (1984), 177–219.

[G-H-V] W. GREUB, S. HALPERIN, R. VANSTONE, Connections, curvature and cohomology, Vol. III (1976), Academic Press.

[G-MP] A. Goresky, R. Mac Pherson, Intersection cohomology theory, *Topology*, **19** (1980), 135–162.

[Gre] M. Green, Infinitesimal methods in Hodge theory. In: Algebraic cycles and Hodge theory, *Lect. Notes Math.*, **1594** (1999).

[Gro1] A. Grothendieck, Standard conjectures on algebraic cycles, algebraic geometry, *Bombay Coll.* (1968), Oxford, 193–199.

[Gro2] A. Grothendieck, Formule de Lefschetz et rationalité des fonctions L , *Sém. Bourbaki*, **279** (1964).

[Har1] G. Harder, Eisenstein Kohomologie and die Konstruction gemischter Motive, *Lect. Notes Math.*, **1562** (1993).

[Har2] G. Harder, Some results on the Eisenstein cohomology of arithmetic subgroups of GL_n , *Lect. Notes Math.*, **1447** (1990), 85–153.

[Hart] R. Hartshorne, Algebraic geometry, *Grad. Texts in Math.*, **52** (1977), Springer.

[Hen] G. Henniart, Une preuve simple des conjectures de Langlands pour $GL(n)$ sur un corps p-adique, *Invent. Math.*, **139** (2000), 439–455.

[H-T] M. Harris, R. Taylor, On the geometry and cohomology of some simple Shimura varieties, *Annals of Math. Stud.*, **151** (2002), Princeton Univ. Press.

[Jan1] U. Jannsen, Mixed motives, motivic cohomology and Ext-G-groups, *Proceed. Int. Congr. Math.* (1994), 667–679.

[Jan2] U. Jannsen, Motives, numerical equivalence and semisimplicity, *Invent. Math.*, **107** (1992), 447–452.

[K-K-M-S] G. Kempf, F. Knudsen, D. Mumford, B. Saint-Donat, Toroidal embeddings, *Lect. Notes Math.*, **339** (1979), Springer.

[Kle] S. Kleinman, The standard conjectures, *Proceed. Symp. Pure Math.*, **55** (1994), 1–20.

[Kna] A. Knapp, Introduction to the Langlands program, *Proceed. Symp. Pure Math.*, **561** (1997), 245–301.

[Kot1] R. Kottwitz, On the λ-adic representations associated to some simple Shimura varieties, *Invent Math.*, **108** (1992), 653–665.

[Kot2] R. Kottwitz, Involutions in Weyl groups, *Represent. Theory*, **4** (2000), 1–15.

[Laf] L. Lafforgue, Chtoucas de Drinfeld et correspondance de Langlands, *Invent Math.*, **147** (2002), 1–242.

[Lan1] R. Langlands, Problem in the theory of automorphic forms, *Lect. Mod. Analysis and Appl. III; Lect. Notes Math.*, **170** (1970), Springer.

[Lan2] R. Langlands, Where stands functoriality today?, *Proceed. Symp. Pure Math.*, **61** (1997), 457–471.

[Lan3] R. Langlands, Automorphic representations, Shimura varieties and motives, *Proceed. Symp. Pure Math.*, **33** (1977).

[Lang] S. Lang, Introduction to modular forms (1976), Springer.

[Lau] G. Laumon, La correspondance de Langlands sur les corps de fonctions, *Sém. Bourbaki*, **873**, (1999–2000).

[Mac] G. Mackey, Infinite dimensional group representations, *Bull. Amer. Math. Soc.*, **69** (1963), 618–686.

[Mor] F. Morel, Voevodsky's proof of Milnor's conjecture, *Bull. Amer. Math. Soc.*, **35** (1998), 123–143.

[Mur] J.P. Murre, Algebraic cycles and algebraic aspects of cohomology and K-theory, *Lect. Notes Math.*, **1594** (1994), 93–152.

[M-W] B. Mazur, A. Wiles, Class fields of abelian extensions of Q , *Invent Math.*, **76** (1984), 179–330.

[Neu] J. Neukirch, Algebraic number theory, *Grund. Math. Wiss.*, **322** (1999), Springer.

[Pha] F. Pham, Introduction à l'étude topologique des singularités de Landau, *Mém. Sci. Math.*, **164** (1967), Gauthiers-Villars.

[Pie1] C. Pierre, Endomorphism from Galois antiautomorphism, *Bull. Belg. Math. Soc.*, **2** (1995), 435–445.

[Pie2] C. Pierre, Algebraic representations of von Neumann algebras, Arxiv: Math-RT/609232 (2006).

[Pie3] C. Pierre, From global class field concepts and modular representations to the conjectures of Shimura-Taniyama-Weil, Birch-Swinnerton-Dyer and Riemann, preprint (2006), Arxiv: RT/0608084.

[Pie4] C. Pierre, Algebraic quantum theory, preprint (2004), Arxiv: Math-ph/0404024.

[Pie5] C. Pierre, Introducing bisemistructures, preprint (2006), Arxiv: GM/0607624.

[Pie6] C. Pierre, The Goldbach conjecture resulting from global-local cuspidal representations and deformations of Galois representations, Arxiv: 08120930 (2008).

[Ram] D. Ramakrishnan, Pure motives and automorphic forms, *Proceed. Symp. Pure Math.*, **55** (1994), 411–445.

[Sch] J. Schwermer, Cohomology of arithmetic groups, automorphic forms and L-functions, *Lect. Notes Math.*, **1447** (1990), 1–29.

[Sel] A. Selberg, Discontinuous groups and harmonic analysis, *Proc. Int. Congr. Math.* (1962), 177–189.

[Ser1] J.P. Serre, Cohomologie des groupes discrets, *Sém. Bourbaki*, **399**, in: Lect. Notes Math. **244**, (1971), 337–350.

[Ser2] J.P. Serre, Groupes de Galois sur Q , *Astérisque*, **161–162**; *Sém. Bourbaki*, **689** (1988), 73–85.

[Ser3] J.P. Serre, Local fields, *Grad. Texts in Math.*, **67** (1979), Springer.

[Shi] G. Shimura, Automorphic functions and number theory, *Lect. Notes Math.*, **54** (1968), Springer.

[Sol] R. Solomon, A brief history of the classification of the finite simple groups, *Bull. Amer. Math.*, **38** (2001), 315–352.

[Tat] J. Tate, Number theoretic background, *Proceed. Symp. Pure Math.*, **33** (1979), 3–26.

[Voe] V. Voevodsky, Triangulated categories of motives over a field, in cycles, transfers and motivic homology theories, *Ann. Math. Stud.*, **143** (2000), Princeton University Press.

[Wat] W. Waterhouse, Introduction to affine group schemes, *Grad. Texts in Math.*, **66** (1979), Springer.